

LUND UNIVERSITY

Online measurements of biological aerosols along the Greenland west coast

Alsved, Malin; Löndahl, Jakob; Finster, Kai; Šantl-Temkiv, Tina

2017

Link to publication

Citation for published version (APA): Alsved, M., Löndahl, J., Finster, K., & Šantl-Temkiv, T. (2017). Online measurements of biological aerosols along the Greenland west coast. Abstract from Nordic Society for Aerosol Research (NOSA) Aerosol Symposium 2017, Lund, Sweden.

Total number of authors: Δ

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights. • Users may download and print one copy of any publication from the public portal for the purpose of private study

- or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00

Online measurements of biological aerosols along the Greenland west coast

M. Alsved¹, J. Löndahl¹, K. Finster², T. Šantl-Temkiv^{2,3}

¹Department of Design Sciences, Lund University, SE-22100, Lund ²Department of Bioscience, Microbiology Section, Aarhus University, DK- 8000, Aarhus ³Department of Physics and Astronomy, Stellar Astrophysics Centre, Aarhus University, DK- 8000, Aarhus

Keywords: bioaerosol, atmosphere, Arctic, fluorescence, marine emissions

Introduction

Biological aerosols may have important effects on clouds and precipitation due to their ability to initiate ice formation at high subzero temperatures (-10°C to -2°C). The knowledge on bioaerosols emitted from ocean surfaces is very limited, especially in remote Arctic regions. The aim of this project was to investigate bioaerosols, and their cloud-active properties along the Greenland west coast. Here we present a first brief overview of micrometer sized aerosol particles and bioaerosol concentrations from measurements and microbial online cell concentrations of seawater.

Methods

Data were collected from a navy patrol vessel over an 11-day long period in August 2016 (Figure 1). Bioaerosols were measured with a BioTrak (Model 9510, TSI Inc.) with a 1 minute time resolution and a flow rate of 28.3 L/min. The instrument detects biological particles by their fluorescence in two bands: 405-500 nm and 500-650 nm. An algorithm provided by the manufacturer is used to interpret the data and avoid false positives. We used NovoCyte flow cytometer to evaluate microbial concentrations in sea water and sea surface microlayer.

Results and conclusion

The total and biological number concentrations along the route are shown in Figure 2 below. The average concentration of biological aerosol particles was about 100 m⁻³. Around 0.003% of the airborne particles detected in the range 0.5-10 μ m bioaerosols. However, the concentrations of biological particles may be underestimated because of threshold values used for the fluorescence detectors.





Figure 1. Ship route during the measurements.

By number, 70% of the biological particles were 1-2 µm and 20% around 3µm. The highest concentrations were found close to populated areas. There was a significant, but negligible, correlation between biological particles and total particle counts (Spearman's $\rho = 0.1$). Thus, the biological particles are likely to originate from specific sources. The seawater microbial concentrations were on average 7.7.10⁴ cells per mL of bulk water and $2.6 \cdot 10^5$ cells per mL in the sea surface microlayer. The low number of likely reflect low bioaerosols thus source concentrations of microorganisms in the ocean.

This work was supported by the Arctic Research Center and the Stellar Astrophysics Centre, Aarhus University, and the Swedish research council Formas. We greatly appreciate the support from the crew of Ejnar Mikkelsen patrol vessel.

