



LUND UNIVERSITY

A Multi-Method Analysis of the Cultural, Sociodemographic and Economic Elements in Osteoarthritis Care

Battista, Simone

2023

Document Version:

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Battista, S. (2023). *A Multi-Method Analysis of the Cultural, Sociodemographic and Economic Elements in Osteoarthritis Care*. [Doctoral Thesis (compilation), Department of Clinical Sciences, Lund]. Lund University, Faculty of Medicine.

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY


PO Box 117
221 00 Lund
+46 46-222 00 00



A Multi-Method Analysis of the Cultural, Sociodemographic and Economic Elements in Osteoarthritis Care

SIMONE BATTISTA

DINOGLI | SCHOOL OF MEDICAL AND PHARMACEUTICAL SCIENCES | UNIVERSITY OF GENOVA
ORTHOPAEDICS | CLINICAL SCIENCES | FACULTY OF MEDICINE | LUND UNIVERSITY



A Multi-Method Analysis of the Cultural, Sociodemographic and Economic Elements in Osteoarthritis Care

Joint Degree of *Doctor of Philosophy* (PhD) in:
Neurosciences – Motor and Sports Activities Sciences



**Università
di Genova**

Medical Science



LUND
UNIVERSITY

Doctoral Dissertation

By the permission of the School of Medical And Pharmaceutical Sciences,
University of Genova, Genova, Italy & Faculty of Medicine, Lund University, Lund, Sweden
To be defended at the Campus of Savona, Savona, Italy on 6 July 2023 at 3.00 PM.

Faculty Opponent

Professor Tanja Stamm

Institute of Outcomes Research, Medical University of Vienna, Vienna, Austria

PhD Candidate: Simone Battista

Main Supervisors: Dr Marco Testa

Dr Andrea Dell'Isola

Co-Supervisor: Prof Martin Englund

Organisation: University of Genova (Genova, Italy), Lund University (Lund, Sweden)

Document name: Doctoral Dissertation

Date of issue: 25 April 2023

Author: Simone Battista

Sponsoring organisations: University of Genova (Genova, Italy), Lund University (Lund, Sweden)

Title: A Multi-Method Analysis of the Cultural, Sociodemographic and Economic Elements in Osteoarthritis Care

Abstract:

Background: Osteoarthritis (OA) quality of care is influenced by various cultural, sociodemographic, and economic elements that shape the context in which first-line interventions are implemented. This thesis aims at providing insights to improve OA care quality by analysing the role of healthcare professionals' knowledge and adherence to clinical practice guidelines and patients' sociodemographic (e.g., sex and age) and economic factors (e.g., income), and their experience and beliefs of OA care, through four studies.

Methods: The four studies analysed: 1) physiotherapists' knowledge of and adherence to OA clinical practice guidelines (survey-based study in Italy); 2) patients' experience of the OA care process (qualitative study in Italy); 3) association between demographic, socioeconomic and disease-related factors and exercise adherence in OA (register-based study in Sweden); 4) income inequalities in people attending an OA first-line intervention and whether these inequalities changed after attending it (cohort study in Sweden).

Results: Study 1 revealed that Italian physiotherapists knew but did not always apply first-line interventions. Study 2 suggested that people with OA experience an uncertain care process due to the lack of clear explanations, resulting in a negative attitude towards first-line interventions. Study 3 found associations between exercise adherence and age, sex, socioeconomic status, disease severity and self-efficacy. All the investigated factors explained 1% of the exercise adherence variability. Study 4 indicated income inequalities among participants to an OA first-line intervention that widened after it.

Conclusion: Several areas in the OA care process require attention. Physiotherapists must bridge the gap between their knowledge of and adherence to OA clinical practice guidelines. Patients need to be guided throughout their care process to be more empowered. Further studies on the factors associated with OA exercise adherence are necessary. Finally, we need to address inequalities in OA care, prioritising upstream interventions.

Keywords: Osteoarthritis, Epidemiology, Qualitative Research, Physical Therapy Modalities, Surveys and Questionnaires, Health Policy, Quality Assurance, Health Care.

Language: English

ISSN: 1652-8220

ISBN: 978-91-8021-432-2

Number of pages: 97

I, the undersigned, being the copyright owner of the abstract of the abovementioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the abovementioned dissertation.

Signature



Date 2023-26-05

Paper 1 © by the Authors

Paper 2 © by the Authors

Paper 3 © by the Authors

Paper 4 © by the Authors (Unpublished manuscript)

University of Genova

School of Medical and Pharmaceutical Sciences

Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health

Lund University

Faculty of Medicine

Orthopaedics, Department of Clinical Sciences

ISSN 1652-8220

ISBN 978-91-8021-432-2

University of Genova, School of Medical and Pharmaceutical Sciences, Doctoral Dissertation

Lund University, Faculty of Medicine, Doctoral Dissertation Series 2023:92

Dedication

*“Let everything happen to you
Beauty and terror
Just keep going
No feeling is final”*

- Rainer Maria Rilke

Table of Contents

| | |
|--|----|
| Abstract | 7 |
| Popular Science Summary | 8 |
| List of Papers | 9 |
| Author’s Contribution to the Papers | 10 |
| Abbreviations | 11 |
| Preface | 12 |
| Background | 13 |
| Osteoarthritis Clinical Practice Guidelines..... | 13 |
| Symptoms and Comorbidities | 14 |
| Care Process | 15 |
| Aims | 16 |
| Cultural Elements | 17 |
| Study 1: Italian Physiotherapists’ Knowledge of and Adherence to Osteoarthritis Clinical Practice Guidelines: a Cross-Sectional Study | 18 |
| Background and Rationale..... | 18 |
| Study Design..... | 19 |
| Participants and Recruitment..... | 19 |
| Italian Physiotherapists’ Knowledge | 20 |
| Italian Physiotherapists’ Adherence | 23 |
| Study 2: Giving an Account of Patients’ Experience: A Qualitative Study on the Care Process of Hip and Knee Osteoarthritis | 25 |
| Background and Rationale..... | 25 |
| Study Design..... | 25 |
| Participants and Recruitment..... | 27 |
| Themes..... | 28 |
| Sociodemographic and Economic Elements | 36 |
| Study 3: Factors Associated with Adherence to a Supervised Exercise Intervention for Osteoarthritis: Data from the SOAR | 37 |
| Exercise Adherence: Why is it Fundamental in OA?..... | 37 |
| Study Design..... | 38 |
| Variables | 38 |
| Participants and Recruitment..... | 42 |
| Factors Associated with Exercise Adherence – Average Association | 45 |
| Factors Associated with Exercise Adherence – Variability Explanation | 51 |
| Study 4: Income-Related Health Inequality and Osteoarthritis Intervention Outcomes: a Cohort Study..... | 52 |

| | |
|---|-----------|
| Inequalities in OA Care | 52 |
| Income | 52 |
| Study Design..... | 53 |
| Variables | 53 |
| Participants and Recruitment | 54 |
| Income-Related Inequalities | 58 |
| Ethical Considerations | 64 |
| General Discussions | 65 |
| Studies' Limitations | 71 |
| Future Perspectives..... | 72 |
| Conclusions..... | 74 |
| References..... | 75 |
| Acknowledgements (English)..... | 93 |
| Acknowledgements (Italian) | 95 |
| Manuscripts..... | 97 |

Abstract

Background: Osteoarthritis (OA) quality of care is influenced by various cultural, sociodemographic, and economic elements that shape the context in which first-line interventions are implemented. This thesis aims at providing insights to improve OA care quality by analysing the role of healthcare professionals' knowledge and adherence to clinical practice guidelines and patients' sociodemographic (e.g., sex and age) and economic factors (e.g., income), and their experience and beliefs of OA care, through four studies.

Methods: The four studies analysed: 1) physiotherapists' knowledge of and adherence to OA clinical practice guidelines (survey-based study in Italy); 2) patients' experience of the OA care process (qualitative study in Italy); 3) association between demographic, socioeconomic and disease-related factors and exercise adherence in OA (register-based study in Sweden); 4) intervention generated income inequalities in people attending a Swedish OA first-line intervention (cohort study in Sweden).

Results: Study 1 revealed that Italian physiotherapists knew but did not always apply first-line interventions. Study 2 suggested that people with OA experience an uncertain care process due to the lack of clear explanations, resulting in a negative attitude towards first-line interventions. Study 3 found associations between exercise adherence and age, sex, socioeconomic status, disease severity and self-efficacy. All the investigated factors explained 1% of the exercise adherence variability. Study 4 indicated intervention-generated income inequalities among participants to an OA first-line intervention.

Conclusion: Several areas in the OA care process require attention. Physiotherapists must bridge the gap between their knowledge of and adherence to OA clinical practice guidelines. Patients need to be guided throughout their care process to be more empowered. Further studies on the factors associated with OA exercise adherence are necessary. Finally, we need to address inequalities in OA care, prioritising upstream interventions.

Popular Science Summary

Patients' quality of care does not depend only on healthcare professionals' knowledge and skills. It also depends on what the patient thinks and wants and the context they are in. Osteoarthritis is the most common joint disease worldwide. Still, only 40% of the patients receive first-line interventions (exercise, education on self-management strategies and diet). This poor care quality can result from cultural, sociodemographic and economic reasons belonging to the healthcare professionals and the patients.

Therefore, we investigated Italian physiotherapists' knowledge of and adherence to osteoarthritis evidence-based intervention with a web survey. Physiotherapists seemed to know the importance of adopting an evidence-based intervention, but they did not always use it in practice. Then, we interviewed patients to understand their beliefs about osteoarthritis. Patients reported exercise as a way to pass the time until they could have surgery. Moreover, they highlighted that healthcare professionals refrained from explaining the role of evidence-based interventions in osteoarthritis care.

Finally, we analysed the 'Swedish Osteoarthritis Registry' (SOAR), with data from approximately 200,000 people with osteoarthritis who attended a first-line intervention in different Swedish hospitals. From this register, we tried to understand if the demographic (e.g., sex and age), lifestyle (e.g., hours of physical activity per week), socioeconomic (e.g., income), and disease-related (e.g., pain intensity) factors could influence adherence to exercise. Furthermore, we tried to understand if people from lower socioeconomic backgrounds experienced more severe symptoms and less benefit from this Swedish osteoarthritis intervention. The factors we investigated in the SOAR influenced only minimally exercise adherence. On top of that, people from lower socioeconomic backgrounds seemed to experience less benefit from the intervention, resulting in a worsening of the inequality that was observed even before the intervention.

List of Papers

Paper I

Battista S, Salvioli S, Millotti S, Testa M, Dell’Isola A. Italian Physiotherapists’ Knowledge of and Adherence to Osteoarthritis Clinical Practice Guidelines: a Cross-Sectional Study. *BMC Musculoskelet Disord.* 2021. 22:380. doi: 10.1186/s12891-021-04250-4

Paper II

Battista S, Manoni M, Dell’Isola A, Englund M, Palese A, Testa M. Giving an account of patients’ experience: A qualitative study on the care process of hip and knee osteoarthritis. *Health Expect.* 2022, 1-17. doi: 10.1111/hex.13468

Paper III

Battista S, Kiadaliri A, Jönsson T, Gustafsson K, Englund M, Testa M, Dell’Isola A. Factors Associated with Adherence To a Supervised Exercise Intervention for Osteoarthritis: Data from the Swedish Osteoarthritis Registry. Accepted in *Arthritis Care Res.* 2023. doi: 10.1002/acr.25135

Paper IV

Battista S, Kiadaliri A, Jönsson T, Gustafsson K, Englund M, Testa M, Dell’Isola A. Income-Related Health Inequality and Osteoarthritis Intervention Outcomes: a Cohort Study. Under Review in *Archives of Physical Medicine and Rehabilitation.*

The full-texts of the papers are reported at the end of the thesis, after the acknowledgements.

Author's Contribution to the Papers

Paper I

Simone Battista made a substantial contribution to the concept or design of the article, and the acquisition, analysis, and interpretation of data for the article. He drafted the article or revised it critically for important intellectual content. He approved the version to be published.

Paper II

Simone Battista made a substantial contribution to the concept or design of the article, and the acquisition, analysis, and interpretation of data for the article. He drafted the article or revised it critically for important intellectual content. He approved the version to be published.

Paper III

Simone Battista made a substantial contribution to the concept or design of the article, analysis, and interpretation of data for the article. He drafted the article or revised it critically for important intellectual content. He approved the version to be published.

Paper IV

Simone Battista made a substantial contribution to the concept or design of the article, analysis, and interpretation of data for the article. He drafted the article or revised it critically for important intellectual content. He approved the version to be published.

Abbreviations

| | |
|--------|---|
| OA | Osteoarthritis |
| YLD | Years Lived with Disability |
| NSAIDs | Non-Steroidal Anti-Inflammatory Drugs |
| CPG | Clinical Practice Guidelines |
| SEP | Socioeconomic Position |
| BMI | Body Mass Index |
| HRQoL | Health-Related Quality of Life |
| RMD | Rheumatic and Musculoskeletal Diseases |
| NICE | National Institute for Health and Care Excellence |
| EULAR | European Alliance of Associations for Rheumatology |
| OARSI | Osteoarthritis Research Society International |
| STROBE | Strengthening the Reporting of Observational Studies in Epidemiology |
| COREQ | Consolidated Criteria for Reporting Qualitative Research |
| WHO | World Health Organisation |
| TJR | Total Joint Replacement |
| SOAR | Swedish Osteoarthritis Registry |
| LISA | Longitudinal Integration Database for Health Insurance and Labour Market Studies' |
| VAS | Visual Analogue Scale |
| RRR | Relative Risk Ratio |

Preface

Osteoarthritis (OA) is an inflammatory disease that can affect any joint in its whole [1–3]. It has a significant global impact [4–6], accounting for 2.2% of all years lived with disability worldwide (YLD) [5, 6]. Thus, OA is hastily becoming the third largest contributor to disability after diabetes and dementia [7]. As a result, high-income countries invest up to 2.5% of their gross domestic product in the care of OA — mostly in response to the growing demand for joint replacements — an expanse that will become unsustainable if we do not reverse this trend [8].

In this context, first-line interventions for OA — which include exercise, education and weight loss (when needed) — are safe and low-cost treatments that are at least as effective as analgesic and non-steroidal anti-inflammatory drugs (NSAIDs) devoid of their side effects [9–11]. Nonetheless, a marked discrepancy between what is known and what is translated into clinical practice remains, as only a limited proportion of people with OA receive appropriate care [12–14]. This disparity can be attributed to many factors that encompass cultural, sociodemographic, and economic influences affecting clinicians and patients. Healthcare providers' knowledge of and adherence to OA clinical practice guidelines (CPGs) are critical contributors to inadequate care [15–17]. Additionally, patient-related factors such as awareness, beliefs, and barriers to first-line interventions play a crucial role in determining the success of OA treatment [18, 19]. Further, sociodemographic and economic aspects can influence an individual's adherence to OA management strategies [20–24]. Finally, it is well established that individuals in lower socioeconomic positions (SEPs) often face inequalities in healthcare, regardless of the healthcare system [25–28].

Therefore, this thesis arises from the need to better understand how to improve the OA care process, with a focus on hip and knee, two of the most common forms of OA. To do so, we have combined different quantitative and qualitative methodologies to gain a comprehensive understanding of the cultural, sociodemographic, and economic factors that shape OA care.

Background

Osteoarthritis Clinical Practice Guidelines

To improve the quality of care of people with OA, different international CPGs have been developed [10, 11, 29]. All OA CPGs recommend exercise, education and diet (when necessary) as first-line interventions to reduce people's symptoms and levels of disability [10, 29, 30].

Exercise is considered a first-line intervention as it can improve people's symptoms and levels of functionality [31, 32]. Exercise has a positive local effect on the joint, improving muscle strength and balance, joint load capacity and stiffness perception [31, 32]. But exercise has shown benefits beyond joint health [33, 34]. It positively affects body weight, lipid metabolism, hyperglycaemia, mood and systemic inflammation [32, 35, 36]. However, people with OA struggle with meeting the recommended levels of exercise [37–39].

The role of diet in managing OA among those overweight is getting more and more attention [40, 41]. Elevated high body mass index (BMI) has been associated with low-grade systemic inflammation, higher biomechanical overload and increased symptoms of OA [40, 41]. Therefore, weight reduction can positively impact joint health by reducing the biomechanical load and restoring functional inflammation response, particularly when combined with exercise [42]. Additionally, OA is frequently accompanied by comorbidities such as metabolic syndromes and type-2 diabetes [41]. Thus, dietary management of these conditions (preventive or curative) is crucial [41]. However, maintaining optimal or losing weight can be challenging, especially for those with impaired mobility [43].

To achieve optimal health outcomes, people with OA must integrate exercise and diet into their lifestyles as part of their self-management approach. Moreover, first-line interventions are always recommended due to their positive impact on symptoms and OA-related comorbidities, regardless of the severity of the disease [44].

Symptoms and Comorbidities

In OA, pain is the most common symptom [45, 46]. People with OA tend to feel intermittent but severe or constant background pain [47, 48]. The former has the most significant impact on people's health-related quality of life (HRQoL), is typical of OA early stages and is triggered by specific activities [46]. The latter tends to be present in the middle stages of OA, typically at night [46]. People with advanced stages of OA feel a combination thereof compounded by episodes of unpredictable pain [49]. These wide varieties of pain responses suggest the presence of different pain phenotypes associated with OA [50].

The pathogenesis of pain seems to be multifactorial and stems from several mechanisms: 1) joint nociception (structural changes, inflammation, neovascularisation, new nerve growth) [51–53]; 2) peripheral sensitisation (neuronal hyperexcitability) [54, 55]; 3) central sensitisation (pain hypersensitivity in the spinal cord and brain, dysfunction of ascending and descending pain pathways) [56]; 4) types of pain mechanisms (nociceptive, inflammatory and neuropathic); 5) chemical mediators (cytokines, TRPV1, NGF, serotonin, TNF, substance P) [46, 57]; 6) contextual aspects (psychosocial, economic and cultural factors) [58, 59]. Nevertheless, the pain mechanisms of OA are still enigmatic, and further research is needed [60].

Additionally, people with OA may experience other physical symptoms such as stiffness, reduced function (e.g., walking difficulties), joint instability, reduced movement, swelling, deformity and crepitus [45]. One out of two people with knee or hip OA has other rheumatic and musculoskeletal diseases (RMD), such as low back pain, arthropathies, and synovial and tendon disorders [61–63]. Beyond physical functions, OA also has negative impacts on mental health. People with OA experience a broader array of symptoms that profoundly impacts their life, including low mood [64], depression [65], and poor-quality of sleep [66]. Finally, 67% of people with OA have at least one other chronic condition [61, 67]. In their systematic review, Hall *et al.* found that approximately 40% of people with OA concurrently have cardiovascular diseases [68]. Furthermore, metabolic conditions such as obesity, diabetes, hypertension, and dyslipidaemia are more prevalent in people with OA than those without it [69].

Therefore, the care process of hip and knee OA must encompass a holistic approach that considers the various aspects of an individual's life that this condition may impact. It is imperative to address not only the affected joints but also to take into account the various spheres of a person's life that may be affected, such as physical function, and overall health.

Care Process

By implementing a high-quality and holistic care process, healthcare providers can ensure that individuals with hip and knee OA receive the most effective treatments, achieving optimal outcomes and improved HRQoL [70]. However, Basedow et al. found the quality of OA care to be suboptimal in all treatment domains [12]. The authors analysed its quality through various indicators gathered from multiple sources such as medical records, patient questionnaires and interviews, and administrative databases and expressed with pass-rate scores [12]. First-line interventions (i.e., exercise, education and diet) achieved the lowest pass-rate scores [12].

Therefore, to improve the quality of the OA care process, it is essential to investigate and address the factors that can impact it. Other than the efficacy of the treatments *per se*, different cultural, sociodemographic, and economic factors that are specific to clinicians and people with OA might play a role in this matter (Figure 1).




Figure 1 Factors affecting the OA care process

As far as clinicians are concerned, their knowledge of and adherence to OA CPGs represent one of the critical points to providing people with OA with proper care. When it comes to patients, their awareness, beliefs and barriers to first-line interventions represent additional factors that can hinder or improve adherence to first-line interventions. As these treatments are lifestyle interventions, people with OA must be aware of their importance and integrate them into their daily routines. Also, people’s sociodemographic and economic elements may influence their adherence to OA first-line interventions such as exercise. Finally, people in lower SEP experience inequalities in healthcare, no matter the healthcare system (public or private). Healthcare interventions addressing behavioural changes (e.g., exercise) can even widen these inequalities (‘intervention-generated’ inequalities).

Aims

This PhD project aims at understanding how different cultural, sociodemographic and economic elements can influence OA care process through four papers. As per the cultural elements, **Study 1** analysed Italian physiotherapists' knowledge of and adherence to OA CPGs (cross-sectional survey-based study in Italy). **Study 2** explored the experience of OA care process giving an account of those who live it first-hand i.e., people with OA (qualitative study based on semi-structured interviews). As per the sociodemographic and economic elements, **Study 3** investigated the associations between demographic, socioeconomic and disease-related characteristics with exercise adherence in OA (register-based study in Sweden). **Study 4** investigated intervention-generated income-related inequalities in a Swedish OA first-line intervention (cohort study in Sweden).

Cultural Elements

Clinicians' knowledge of and adherence to OA CPGs represent one of the main issues in providing people with OA with proper care. Physiotherapists are one of the main healthcare professionals involved in OA care. However, implementing CPGs for RMD appears suboptimal among physiotherapists [71]. Therefore, Study 1 examined the knowledge of and adherence to OA CPGs in a cohort of Italian physiotherapists.

Patients' expectations, beliefs and emotions revolving around their care process represent other factors that can hinder or improve their adherence to first-line interventions [18, 19, 72–75]. People with OA must be aware of the importance of these interventions and learn how to integrate them into their daily routines. In OA, efforts to assess and understand patients' experiences of the care process could help policymakers foster improvements in healthcare providers' interpersonal aspects and patients' expectations of how healthcare should be delivered [74]. Therefore, study 2 explored people's experience of the OA care process.

Study 1: Italian Physiotherapists' Knowledge of and Adherence to Osteoarthritis Clinical Practice Guidelines: a Cross-Sectional Study

Background and Rationale

Clinical Practice Guidelines: Meaning and Application

CPGs are “statements that include recommendations intended to optimise patient care. These statements are informed by a systematic review of evidence and an assessment of the benefits and costs of alternative care options” [76]. In the last few decades, several OA CPGs have been released by the European Alliance of Association for Rheumatology (EULAR), Osteoarthritis Research Society International (OARSI) and the National Institute for Health and Care Excellence (NICE) [10, 11, 29]. These CPGs recommend exercise and education as first-line interventions, no matter the severity of the disease [77, 78]. However, OA is often under-treated and underdiagnosed, hindering its care process [12–14, 79]. Egerton et al. highlighted that clinicians who work with people with OA perceived themselves as under-prepared and unfamiliar with CPGs [80].

Physiotherapists' Attitudes to Clinical Practice Guidelines

Physiotherapists are paramount in the OA care process due to their unique training in exercise regimen prescription. However, implementing CPGs for RMD appears suboptimal among physiotherapists [71]. RMD care is complex as clinicians are called to balance knowledge of the best evidence with patients' preferences and beliefs [81–83]. This issue is compounded by several barriers, such as clinical applicability, English language knowledge and lack of time to retrieve and read evidence [81–84]. Zooming in on OA CPGs, similar results were found on the knowledge of and adherence to CPGs [85–89]. However, these studies are inconclusive. They only examined the knowledge of and adherence to CPGs in isolation [86–89] or solely focused on specific treatments (e.g. therapeutic exercise) [90]. Thus, analysing the knowledge of and adherence to CPGs in the same cohort would allow for a better understanding of the so-called *evidence-to-practice gap* [81], which, if unaddressed, can lead to suboptimal care for people with OA [91].

What Do We Know about OA Management in Italy?

Italy is a large southward Mediterranean country with the most significant proportion (21.4%) of older adults (aged>65) in Europe [92]. From 2001 to 2016, 812,639 total knee replacements were performed on people over 40 in Italy. The total number of surgeries increased by 262%, with an average annual growth rate of 6.6% and an increase of 45% in incidence rate is expected by 2050 [93]. This data suggests the urge

to understand how OA is treated in Italy. Moreover, gathering information from this country might allow for our results to be generalised to other Mediterranean countries, which appeared to have higher educational needs than the Northern-European ones regarding RMD management [94]. Thus, we conducted this study to explore the knowledge of and adherence to OA CPGs among Italian physiotherapists to identify the possible existence of an evidence-to-practice gap in OA interventions.

Study Design

We carried out a web-based cross-sectional study investigating Italian physiotherapists' knowledge of and adherence to OA CPGs. We decided to use a web survey as they are a valuable resource for understanding people's cultural contexts outside of an experimental setting in a real-world scenario [95]. We created the survey using the 'International Handbook of Survey Methodology', one of the most central guides for survey creation [96]. Finally, we reported our study following the 'Checklist for Reporting Results of Internet E-Survey' and the 'Strengthening the Reporting of Observational Studies in Epidemiology' (STROBE) recommendations for reporting observational studies [97, 98].

We based this survey on the EULAR, OARSI and NICE guidelines [10, 11, 29]. The survey was delivered in Italian and divided into two sections investigating (1) the knowledge of and (2) the adherence to OA CPGs, respectively. Before the first section, we included questions investigating the physiotherapists' sociodemographic factors (e.g. sex, age, years of experience) and whether they had read any OA CPG. The last draft of the questionnaire was tested for clarity on a sample of six physiotherapists specialised in RMD rehabilitation, and no significant changes were made.

We delivered the online survey through Microsoft 365 Forms, as it is a secure web application to build and manage online surveys and databases, respecting the European General Data Protection Regulations [99]. In the beginning, the questionnaire read a brief cover letter and the informed consent outlining the aim of the study. The cover letter emphasised that participation in the survey was voluntary and that anonymity and confidentiality were guaranteed. Our study followed the 'Declaration of Helsinki' and granted by the Ethics Committee for University Research (CERA: Comitato Etico per la Ricerca di Ateneo), University of Genova (approval date: 15/06/2020; CERA2020.07).

Participants and Recruitment

We delivered the online questionnaire through the Italian Association of Physiotherapists (AIFI) and the newsletter of the 'RMD rehabilitation' post-graduate degree of the University of Genova newsletters. Using the latter newsletter might have led to possible 'sampling error' as these physiotherapists received thorough training in RMD rehabilitation. To be included, physiotherapists had to treat at least one person with OA in the six months before the questionnaire. A preliminary question on the

survey assured this eligibility criterion. Those who flagged ‘No’ could not partake in the other part of the questionnaire. Hence, we reached 822 physiotherapists between June and July 2020 (Table 1).

Table 1 Participants’ demographic characteristics

| | |
|---|--------------|
| Age (years)(mean,(SD)) | 35.77 (13.3) |
| Assigned sex (at birth) (female) (N (%)): | 387 (47) |
| Years of practice (N (%)): | |
| Less than 1 year | 87 (11) |
| From 1 to 5 years | 319 (39) |
| From 6 to 10 years | 149 (18) |
| More than 10 years | 267 (32) |
| Post-graduate degrees (N (%))*: | |
| I Level Master Degree† | 382 (47) |
| Master of Science (MSc)/II Level Master‡ | 122 (15) |
| Doctor of Philosophy (PhD) | 36 (4) |
| Others | 282 (34) |

Legend: N, number; %, percentage; *, Percentages calculated based on the highest Post-Graduate Degrees achieved; †, Academic degree that can be gained after BSc (Italian education system); ‡, Academic degree that can be gained after MSc (Italian education system).

Italian Physiotherapists’ Knowledge

To test Italian physiotherapists’ knowledge, participants expressed agreement with 24 CPG statements through a 1 (completely disagree) to 5 (completely agree) Likert scale (Table 2). Participants who wholly or partially agreed (scores 4–5) were considered to agree with the statements. However, we added eleven reversed statements to avoid any possible acquiescence bias (i.e. the tendency to agree with all the survey statements) [100]. In this case, disagreement with reversed statements (scores 1–2) would indicate an agreement with the CPGs. Without a standard threshold, we defined a $\geq 70\%$ agreement with a statement as consensus [15, 84, 101, 102]. The frequencies of answers were calculated, and a visual representation through a bar chart graph was reported.

Table 2 Statements derived from OA CPGs

| Statements | Clinical Practice Guidelines |
|--|---|
| 1) Exercise can be effective on all patients, regardless of the pain severity. | NICE (1.2.5-1.4.1); EULAR (3-6-7); |
| 2) In an advanced stage of the disease, exercise can damage the joint (reverse statement). | NICE (1.2.5-1.4.1); EULAR (2-3-6-7); |
| 3) The rehabilitation programme must always include a part of education on the pathophysiology of osteoarthritis and self-management strategies. | NICE (1.3.1-1.3.2-1.3.3); EULAR (3-5); OARSI (tables 2-3) |

| | |
|--|---|
| 4) The rehabilitation programme should always include a part of manual treatment (reverse statement) | NICE (1.4.2); EULAR (-); OARSI (-) |
| 5) Exercise should only be undertaken after prescribing drug treatment to control pain (reverse statement). | NICE (1.2.5-1.4.1); EULAR (3-6-7); |
| 6) The use of topical anti-inflammatory drugs is effective for pain relief for knee osteoarthritis. | NICE (1.5.3); EULAR (-); OARSI (table 2) |
| 7) Radiographic findings are needed to express a functional diagnosis of osteoarthritis (reverse statement). | NICE (1.1.1); EULAR (1); OARSI (-) |
| 8) Radiographic findings are needed to plan the physiotherapy treatment (reverse statement). | NICE (1.1.1); EULAR (1); OARSI (-) |
| 9) Physical activity should be avoided because it can damage the joint (reverse statement). | NICE (1.2.5-1.4.1); EULAR (3-6-7); |
| 10) The use of topical anti-inflammatory drugs is effective for pain relief for hip osteoarthritis. | NICE (-); EULAR (-); OARSI (-) |
| 11) In case of severe joint degeneration, it is necessary to recommend rest from physical activity (reverse statement). | NICE (1.2.5-1.4.1); EULAR (2-3-6-7); |
| 12) In cases of severe pain (VAS \geq 6/10), arthroplasty surgery should be preferred to rehabilitation (reverse statement). | NICE (1.6); EULAR (-); OARSI (-) |
| 13) The use of TENS should be considered. | NICE (1.4.4); EULAR (-); OARSI (-) |
| 14) The use of physical therapies such as lasers, TECAR and ultrasound therapy should be considered (reverse statement). | NICE (1.4.4); EULAR (-); OARSI (-) |
| 15) In addition to the rehabilitation treatment, it is useful to recommend physical activity (for example, yoga, swimming, Nordic walking). | NICE (1.2.5-1.3.2-1.4.1); EULAR (-); OARSI (tables 2-3) |
| 16) It is important to recommend weight loss to overweight or obese patients. | NICE (1.2.5-1.4.3); EULAR (3-8); OARSI (tables 2-3) |
| 17) Age > 45, pain and absence of joint stiffness (or <30 min) in the morning are sufficient to diagnose osteoarthritis. | NICE (1.1.1); EULAR (-); OARSI (-) |
| 18) The use of comfortable footwear, braces or aids should be considered. | NICE (1.3.2-1.4.7-1.4.8-1.4.9); EULAR (3-9-10); OARSI (tables 2-3) |
| 19) It is advisable to refer the patient for arthroscopy surgery to reduce symptoms and start/continue treatment (reverse statement). | NICE (1.4.10); EULAR (-); OARSI (-) |
| 20) It is necessary to assess the impact of osteoarthritis on function, quality of life and disability. | NICE (1.2.1); EULAR (1); OARSI (-) |

| | |
|---|--|
| 21) At least 10-12 sessions are needed to ensure proper treatment for osteoarthritis. | NICE (1.4.1); EULAR (6); OARSI (-) |
| 22) In the treatment for osteoarthritis, the patient's adherence to the treatment must be motivated. | NICE (1.3.2-1.4.1-1.7.1); EULAR (-); OARSI (-) |
| 23) Joint hyaluronic acid and/or corticosteroid infiltrations should be considered. | NICE (1.5.12-1.5.13); EULAR (-); OARSI (table 2) |
| 24) The supplements of chondroitin and glucosamine should be considered (reverse statement). | NICE (1.4.5); EULAR (-); OARSI (-) |

A consensus was achieved for 13/24 statements (Figure 2). These statements addressed the role of clinical assessment, exercise, education, weight loss, and the effectiveness of physical therapies in OA. A consensus was not reached on the role of supplements, radiographic findings, manual therapy, topical non-steroidal drugs, TENS, the number of sessions and the criteria for clinical diagnosis.




Figure 2 Levels of agreement with the reported statements

* The black dashed line represent the line of agreement

Italian Physiotherapists' Adherence

To test Italian physiotherapists' adherence, participants were shown a clinical case with different interventions (Figure 3). Clinical vignettes are valuable for assessing physiotherapists' treatment adherence [103]. Participants were classified as 'Delivering' (all core treatments selected), 'Partially Delivering' (some core treatments missing) and 'Non-Delivering' (at least one selected non-core treatment), the recommended intervention, depending on chosen interventions.

| | | |
|---|---|---|
| Clinical Scenario: Maria, a 72-year-old housewife, lives with her husband, who is in good health. She cultivates the hobby of gardening. She has been suffering for ten years from knee pain which, in certain periods, forces her to take NSAIDs and to limit daily activities for a few days. Over the past two years, the pain has become increasingly frequent (VAS 5/10), so that she has decided to find some help with the housework and she is struggling to take care of the garden. She also suffers from diabetes and is overweight (BMI 28). She decides to consult her physician, who recommends her to do a visit to the physiotherapist. | | |
| Management: | | |
| Core Treatment | Partially Core Treatment | Non-Core Treatment |
| <ul style="list-style-type: none"> Evaluation and planning of the rehabilitation treatment; Weight loss advice. | <ul style="list-style-type: none"> Referral to the physician for drug therapy. | <ul style="list-style-type: none"> Referral to the physician for arthroscopic surgery (joint debridement); Referral to the physician for prosthetic intervention. |
| Assessment: | | |
| Core Treatment | | |
| <ul style="list-style-type: none"> Assessment of the quantity and quality of pain; Assessment of the function; Assessment of disability and participation. | | |
| Treatment: | | |
| Core Treatment | Partially Core Treatment | Non-Core Treatment |
| <ul style="list-style-type: none"> Specific exercise on the joint (muscle strengthening); Generic exercise (aerobic exercise or generic physical activity); Education on the pathophysiology of osteoarthritis. | <ul style="list-style-type: none"> Manual therapy (mobilisation and/or massage); TENS; Load reduction devices (braces, insoles or walking aids); Hyaluronic acid and corticosteroid injections. | <ul style="list-style-type: none"> Activity rest (reduce the load on the joint); Other physical therapies (Laser, Ultrasound etc.); Supplement integration: glucosamine and chondroitin. |
| For how many sessions would you treat this patient? | | |
| <ul style="list-style-type: none"> For less than 5 sessions; Between 5 and 10 sessions; For more than 10 sessions. | | |

Figure 3 Clinical case scenario

We classified the interventions as 'core treatment', 'partially core treatment', and 'non-core treatment' following OA CPG grading (Figure 4) [10, 11, 29, 104]. This grade is based on the level of evidence that CPGs attribute to each treatment combined with interventions' benefits and harms, patients' and clinicians' values and preferences and resource implications [105]. As we can see, exercise, education and weight loss are considered by all CPGs as 'strongly recommended' (core treatments).

However, not the same happened for the other treatments where disagreements are present. For these treatments, the most updated CPG was considered to develop the questionnaire.

| Treatments | OARSI (2019) | | NICE (2016) | | EULAR (2013) | |
|---------------------------------------|--------------|------|-------------|------|--------------|------|
| | Hip | Knee | Hip | Knee | Hip | Knee |
| Non-Pharmacological Treatments | | | | | | |
| Physical Activity | ○ | ○ | ○ | ○ | ○ | ○ |
| Patient Education | ○ | ○ | ○ | ○ | ○ | ○ |
| Weight Loss | ● | ○ | ○ | ○ | ○ | ○ |
| Manual Therapy | | | ● | ● | | |
| CBT | ● | | | | | |
| TENS | | | ● | ● | | |
| Orthosis and Aids | | | ● | ● | ● | ● |
| Tai-Chi | ○ | ○ | | | | |
| Yoga | ○ | ○ | | | | |
| Acupuncture | ● | ● | ● | ● | | |
| Pharmacological Treatments | | | | | | |
| Oral NSAIDs | ● | ● | ● | ● | ○ | ○ |
| Topic NSAIDs | | ● | | ● | ○ | ○ |
| Joint Supplements | | ● | ● | ● | | |
| Hyaluronic Acid Injection | ● | ● | ● | ● | | |
| Glucocorticoid Injection | ● | ● | ● | ● | | |
| Surgical Treatments | | | | | | |
| Total Joint Replacement | | | ● | ● | | |
| Debridment | | | ● | ● | | |

Figure 4 CPG position on OA interventions

Legend: OARSI, Osteoarthritis Research Society International; NICE, National Institute for Health and Care Excellence; EULAR: European Alliance of Association for Rheumatology; White dot: strongly recommended; Grey dot: partially recommended; Black dot: partially or strongly not recommended.

Based on their answers, 25% of the participants were classified as ‘Delivering’, 22% as ‘Partially Delivering’ and 53% as ‘Non-Delivering’. The ‘Delivering’ group provided the patient with all the CPGs’ recommended treatments. Only half delivered weight loss advice in the ‘Partially Delivering’ group. Most of the sample assessed functionality, disability, participation, and pain and offered muscle-strengthening exercises and education. About half of the group prescribed general exercise (e.g., aerobic exercise and general physical activity). Finally, in the ‘Non-Delivering Group’, most participants advised load reduction (rest) and other physical therapies (e.g., ultrasound and laser) not recommended by CPGs.

Study 2: Giving an Account of Patients' Experience: A Qualitative Study on the Care Process of Hip and Knee Osteoarthritis

Background and Rationale

The Care Process

The expression of care is a fundamental aspect of human ontology, as individuals are interdependent on one another [106, 107]. Care is rooted in human anthropology, and Benedetti views the care process as a “ritual” performed around the patient, encompassing their intricate psychosocial context [108]. Positive beliefs, expectations, and emotions experienced by patients around the care process can amplify the specific effect of treatments [59, 108, 109]. Conversely, once negative beliefs, expectations and emotions step in, they can hinder patients' progress.

Understanding Patients' Experience

More and more efforts have been put into assessing and understanding patients' experience of their care process, focusing on their expectations, beliefs and emotions around their healthcare [74]. Through this, researchers and health-policy makers can collect significant evidence on how to improve patients' care process, starting from the experience of those who live it first-hand. Qualitative studies are considered the optimal study design for investigating people's experiences and understanding their perspectives on a given topic, as they provide an in-depth exploration of the subject matter [110].

With regards to the Italian healthcare system, it affords its citizens comprehensive coverage with minimal fees. However, the lack of nationally recognised CPGs for the management of OA by the Italian Higher Institute of Health creates an absence of standardised care processes for this condition. As a result, healthcare professionals must resort to utilising international CPGs, which we have indicated in Study 1 to have limited adherence among Italian physiotherapists [15]. Hence, this qualitative study examines people's experiences of the OA care they received within the Italian healthcare system.

Study Design

A qualitative study was conducted with semi-structured interviews analysed through a descriptive phenomenological approach. The descriptive phenomenological inquiry aims at ensuring “direct explorations, analysis, and descriptions of particular phenomena [as in this case – the care process], as free as possible from unexamined presuppositions, maximising intuitive presentation” [110–114]. The underpinning intent of the phenomenological researcher is to give voice and power to people who experienced, or

are experiencing, first-hand the phenomenon of interest as, in this instance, the care process as lived by people with OA [113]. Although we had aimed at getting a deeper understanding of the participants' experience, we must acknowledge that most of the final retrieved themes remain on a surface level of significance.

The semi-structured interview protocol was created for this study based on prior literature on OA [10, 11, 15, 18, 29, 80, 104, 115–117] and input from a diverse group of experts, including physiotherapists, psychologists, nurses, and individuals with osteoarthritis (OA). The interview guide consisted of open-ended questions exploring various aspects of the OA care process, including experiential and emotional dimensions, expectations, and beliefs. Further exploration of participants' experiences was facilitated using follow-up questions such as "Can you give me an example?" or "Can you explain what you mean by that?" At the beginning of each interview, participants were asked to provide informed consent and demographic information, such as age, gender, nationality, and living location, as well as clinical information, such as height, weight, and joint(s) affected by OA. Only the interviewer and participant were present during the interview, and no follow-up interviews were conducted. The interviews were conducted remotely and recorded via Microsoft Teams, and saved in multimedia format on a University of Genova OneDrive folder, which only researchers had access to. Then, they were transcribed, and the transcriptions were anonymised in chronological order (P1, P2 etc.). Once the transcription was over, the recordings were deleted. Ethical approval was granted by the Ethics Committee for University Research (CERA: Comitato Etico per la Ricerca di Ateneo), University of Genova (approval date: 15/06/2020; CERA2020.07). The research was conducted in respect of the Declaration of Helsinki and reported following the Consolidated Criteria for Reporting Qualitative Research (COREQ) [118].

The interviews were conducted by SB. To ensure impartiality, participants were unaware of SB's professional background and had no prior close relationships with him. The interviews were performed through videoconferencing and were recorded for transcription. Both audio and visual recordings were transcribed *verbatim* and analysed by SB and MM. SB and MM are both PhD candidates, with SB trained in physiotherapy and MM in psychology. They both received training in qualitative research methodologies and identified themselves as men.

To ensure the study rigour and trustworthiness, we adopted multiple strategies. First, we documented and shared field notes ("Memos") after completing each interview to promote reflexivity [119]. Then, the research team met frequently to refine the themes and subthemes before agreeing on the final themes. Afterwards, an audit trail containing meeting notes, analysis discussions, and research decisions was continuously reorganised [119]. Lastly, a Synthesised Member Checking was performed to improve the credibility of the analysis [120]. Specifically, we provided the participants with a one-page summary highlighting the main themes and subthemes identified in the study and a plain explanation of the key findings. Thus, we asked the interviewees to send us

back any comments about the results and doubts and concerns about our findings. All participants agreed with what was retrieved, and we did not change the final themes and subthemes.

Participants and Recruitment

A purposeful sampling method was employed to attain a wide range of experiences and maximum variation [121]. Participants with hip, knee, or hip and knee OA residing in urban and suburban areas were contacted to encompass a diverse understanding of the phenomenon being studied. An alliance was formed with physicians specialised in orthopaedics, rheumatology, general practitioners, and other healthcare professionals, such as physiotherapists and nurses, to create a network for recruiting participants. Healthcare professionals were initially approached by the research team and informed about the study's objectives and data collection procedures. After obtaining their cooperation, eligible participants were informed about the study's aim, the interview process, and the confidentiality and anonymity of the data by the healthcare professionals in the network. Participants were free to withdraw from the study and encouraged to voice any questions or concerns. Only those who expressed interest in participating were personally contacted by the researcher to obtain informed consent and arrange the interview. A snowball sampling technique was also employed to reach individuals within the network of participants with OA who agreed to participate [121].

Eligible participants for this study were individuals with a physician diagnosis of hip or knee OA who could speak Italian and willing to participate. Patients reporting primary OA symptoms in joints other than the hip or knee were excluded from the eligibility criteria. The recruitment process was terminated once data saturation was achieved, as determined by the two authors (SB and MM) analysing of the interviews. The data saturation was evaluated using an inductive thematic saturation method, in which SB and MM continued to conduct interviews and analyse the data simultaneously until no new themes emerged [122]. Hence, we interviewed eleven Italian people from northern Italy (table 3).

Table 3 Interviewees' demographic and clinical characteristics

| Patient | Age | Gender | BMI | Retirement | Affected Joint(s) | Diagnosis |
|---------|-----|--------|------|------------|-------------------|---------------|
| P1 | 49 | M | 26.3 | No | Hip | X-Ray |
| P2 | 68 | W | 26.0 | Yes | Hip and Knee | MRI |
| P3 | 73 | W | 27.3 | Yes | Knee | X-Ray |
| P4 | 47 | W | 28.7 | No | Hip | MRI |
| P5 | 72 | W | 25.8 | No | Hip | X-Ray and CT |
| P6 | 55 | M | 34.0 | No | Hip and Knee | X-Ray |
| P7 | 45 | W | 25.1 | No | Hip | X-Ray and MRI |
| P8 | 66 | M | 28.7 | Yes | Knee | X-Ray |
| P9 | 73 | W | 28.7 | Yes | Knee | X-Ray |
| P10 | 65 | M | 24.7 | No | Hip | X-Ray |
| P11 | 56 | M | 22.2 | No | Hip and Knee | X-Ray |

Legend: P, person; M, man; W, woman; BMI, Body Mass Index; OA, Osteoarthritis; MRI, magnetic resonance Image; CT, computed tomography.

Themes

The transcriptions of the interviews were thoroughly analysed by SB and MM. The analysis of the transcripts involved an initial independent analysis by each author, followed by a joint review. This process involved the identification of subthemes, themes, concepts, and patterns through line-by-line examination of the data, with meaning units framed into codes representing essential aspects of the participants' experiences. The emerging codes were compared with previously identified codes to provide a deeper understanding of the phenomenon of interest and to generate focused codes. The focused codes were then synthesised and merged to extract final subthemes and themes. Representative quotes were selected for each theme and reported anonymously, with the themes derived from the data and not predetermined.

Analysis of the interview data revealed seven main themes (Figure 5) related to the OA care process: 1) Experiencing a sense of uncertainty; 2) Establishing challenging relationships with the self and the other; 3) Being stuck in one's own or the health professionals' beliefs about the disease management; understanding; 4) Dealing with one's own attitudes towards the disease; Understanding 5) the facilitators of and 6) the barriers to the adherence to therapeutic exercise; 7) Developing an uneasy relationship with food. The themes (and related subthemes) are discussed and explored hereafter.

| | | | | | |
|---|--|---|--|--|-------------------------------------|
| <i>Experiencing a sense of uncertainty</i> | Need of a straightforward treatment | Doubts (treatments to follow and pathology genesis) | Different opinions heard by various health professionals | Frustration and anger | |
| <i>Establishing Challenging Relationship with the Self and the Other</i> | Not being understood and the importance of empathy | Shame | Hopelessness | Use of metaphors to describe the pathology | |
| <i>Being Stuck in One's Own or the Health Professionals' Beliefs about the Disease Management</i> | Sealed faith (surgery) | OA as a pathology of the old adults | Necessity of radiographic findings (diagnose/treatment) | (Ab)use of passive therapies | Movement as dangerous for the joint |
| <i>Dealing with One's Own Attitudes Towards the Disease</i> | Fight, resignation and acceptance | Coping strategies (mostly passive) | | | |
| <i>Understanding the Facilitators of the Adherence to Therapeutic Exercise</i> | Importance of being active | Perceived exercises as concrete support to the cure | Mean to maintain functionality | Willingness to change life-habits | |
| <i>Understanding the Barriers to the Adherence to Therapeutic Exercise</i> | Cost and lack of time | Lack of clear indications | Lack of willpower and fatigue in changing life habits | Exercise perceived useful only after surgery | |
| <i>Developing an Uneasy Relationship with Food</i> | Diet as fatigue and deprivation | Diet seen as useful only to lose weight | | | |

Figure 5 Themes and subthemes in a glance

Theme 1: Experiencing a Sense of Uncertainty

The study participants shared a common perception of uncertainty regarding the care process for OA, specifically regarding the appropriate approach and treatments to take. Participants reported a lack of clear evidence-based recommendations compounded by healthcare professionals’ personal biases and attitudes in the OA decision-making process.

“There is an almost religious way of thinking about how to deal with the pathology. It is not an exact science; when you choose the physicians, you choose the treatment”
(P1, male, age 49)

The study participants generally lacked an understanding of the underlying causes of their disease. Despite this, they attributed the condition to factors such as overuse and poor posture, with a primarily biomechanical perspective. Additionally, the explanations given to them by healthcare professionals were perceived to be vague and unsatisfactory.

“I thought [OA] was a consequence of bad posture, as I’ve been using my leg wrongly after slipping on ice once.” (P6, male, 55)

“They tried to explain to me how OA works somehow, but I still don’t have a clear idea of how it works.” (P3, female, 73)

The participants reported feeling uncertain and apprehensive about managing OA due to a lack of clear and consistent indications from healthcare professionals and conflicting opinions from various sources.

“I was worried because we, as patients, hear different opinions coming from our friends and acquaintances that give us their personal point of view on how they take care of their disease.” (P9, female, 73)

As a result of their feelings of uncertainty, the interviewees sought information from multiple sources, including healthcare professionals, personal recommendations, and the Internet.

“In my experience, I’ve had to consult two or three physicians unless the first two agree.” (P5, female, 72)

“No, the doctors did not explain it [OA] to me. But, eventually, I looked it up on the Internet and found answers to my questions.” (P8, male, 66)

The participants experienced frustration and anger due to the absence of a precise treatment plan. The lack of specific indications and the presence of conflicting opinions among various sources of information led the interviewees to either neglect their condition or make decisions based on personal assumptions and instincts.

“It is very frustrating for a patient [not to have a precise indication] because you expect to have a disease, and a common one too, so the care process should be clear.” (P1, male, 49)

“Eventually, I did not do anything anymore, just nothing.” (P5, female, 72)

“... Yes, I would like to have a precise guideline, also regarding nutrition... It looks as if there are some things that are left to our intuition.” (P2, female, 68)

Theme 2: Establishing Challenging Relationships with the Self and the Other

Concerning their relationship with themselves and others, the interviewees shared everyday experiences of shame and hopelessness, articulated with different nuances. The former feeling was more prevalent when the participants were in the presence of loved ones. At the same time, the latter was more pronounced due to the sense of despair

conveyed by the healthcare professionals they consulted. Participants reported embarrassment stemming from exhibiting their conditions and limitations to others.

“I felt it [shame] recently. I went to the beach with my granddaughter [...] she wanted me to be involved in her games, and she said, “Grandma come, sit down next to me”. I had to kneel down to play in the sand with her... I felt, how can I say... erm... like a piece of wood, like someone who can no longer manage their body.” (P2, female, 68)

Additionally, the participants expressed a feeling of hopelessness regarding the prognosis they received and the belief among physicians that the condition was irreversible and would inevitably worsen.

“Erm.. Yes [I can only do surgery].. because I dragged it on for too long, and they told me that I have no other possibilities with other [non-surgical] interventions.” (P10, male, 65)

Both shame and hopelessness were closely tied to a perceived lack of empathy from their social network and healthcare providers. Specifically, the interviewees reported that their acquaintances failed to acknowledge certain aspects of their OA (such as limping), causing them to feel ashamed of their condition.

“What annoys me the most is when people that know my condition ask me, “What did you do? Why are you limping?” This makes me really upset because others see what I sometimes don’t even notice.” (P1, male, 49)

Additionally, they felt they were being treated as mere numbers rather than human beings by healthcare providers, contributing to their hopelessness.

“The orthopaedic surgeon did not give me much attention, and they told me that I have OA and that I have to live with it.” (P2, female, 68)

Finally, some participants expressed their self-into-the-pathology using relevant metaphors that helped them better understand their condition.

“I see it [the joint affected by OA] as a mountain which is crumbling.” (P3, female, 73)

Theme 3: Being Stuck in One's Own or the Health Professionals' Beliefs about the Disease Management

The interviewees held a widespread belief that resulted from their interactions with health professionals: surgery was an inevitable outcome for individuals with OA and represented a permanent and inescapable fate.

“It [OA surgery] is something you think about every day, something you try to resist, but that is your fate.” (P1, male, 49)

“And so when I went to see him [the physician], he said, “no madam, your joint is ruined... try and get on with it for as long as you can, but sooner or later you will have to do it [surgery].” (P9, female, 73)

Another widely held belief among the participants was that OA is an age-related condition. This perception made them feel like they were undergoing an abnormal ageing process, even if they still felt active and vibrant.

“The doctor told me: “You know that if I did not know that these x-rays belong to you, I would think that they belong to another person who is at least 30 years older than you”... but, I guess I did not feel as bad as he was describing me.” (P11, male, 56)

In fact, based on the narratives gathered, it was revealed that health professionals were taken aback by the presence of radiographic evidence of OA in younger participants. The reliance on radiographic findings to diagnose OA and formulate a care plan was perceived by the interviewees as a primary focus for physicians rather than the symptoms they reported.

“They told me: here we have the problem, and it is evident as we can see from the x-ray.” (P11, male, 56)

Two other prevalent perceptions held by the interviewees based on their experiences with healthcare professionals were related to the management of OA. These were the frequent utilisation of passive treatments to delay surgical interventions and the limited recommendation for physical activity, which was perceived as a risk factor for early surgery.

“And he [the doctor] told me that I was too young for surgery, and he recommended I do this therapy, to put some ice on my joint.” (P5, female, 72)

“The doctor told me: “You have to try to postpone surgery for as long as you can. So please stop [any physical exercise].” (P1, male, 49)

Theme 4: Dealing with One's Own Attitudes Towards the Disease

The interviewees displayed varying attitudes towards OA. Older patients perceived OA as an indication of surrendering to the ageing process, whereas younger patients viewed it as something they either must or can resist.

“Maybe I am accepting my becoming old, what can I say...” (P3, female, 73)

“From a certain perspective, I took it [OA] positively since it is something I have to fight against.” (P1, male, 49)

However, they all matured a sense of acceptance sooner or later, as if OA was something they could not change.

“Besides, I am also a fatalist, things happen in life, and when they do, you face them.” (P4, female, 47)

All interviewees adopted primarily passive coping strategies to manage their symptoms of OA. The predominant features of their care process were the utilisation of medication to relieve pain and the implementation of various physical therapies, with limited scope for active therapies.

“It's not an issue for me to take some pills not to feel any pain.” (P4, female, 47)

“I thought that by doing some thermal treatments [...] mud treatments [...] mesotherapy and other things ... I thought that with them I would sort the disease out.” (P5, female, 72)

Theme 5: Understanding the Facilitators of the Adherence to Therapeutic Exercise

According to the interviewees, various facilitators were identified in their exercise adherence, with the perceived benefit from the exercises being the most reported. In particular, the experience of being active was viewed as an essential requirement of the body to preserve functionality and provide tangible and immediate support to the care

process, in contrast to taking medication, which can take time to produce noticeable results.

“...The body has to move...” (P3, female, 73)

“It [physical exercise] is not like taking supplements with hyaluronic acid, those (supplements) you do not see what they do.” (P1, male, 49)

Additionally, the role of determination was emphasised by all participants as a crucial factor in adhering to therapeutic exercise. Their perception of willpower varied; some viewed willpower as a facilitator, and others as a hindrance. However, they all concurred that a solid resolution to alter their lifestyle was essential for sustained adherence to an exercise regimen, and that determination was a necessary component in making such life changes.

“...Determination and willpower [to change life-habits].” (P7, female, 45)

Theme 6: Understanding the Barriers to the Adherence to Therapeutic Exercise

The interviewees identified several barriers that hindered their adherence to therapeutic exercise, including the cost of therapy, the required time investment, unclear guidelines, a lack of motivation to modify their lifestyles, and the belief that exercise was only proper after surgery. They noted that participating in physical activity required a significant allocation of time, energy, and resources, often conflicting with their work and familial responsibilities.

“Yes... but also from an economic point of view [it is difficult to do supervised exercises].” (P2, female, 68)

“We are trapped into a spiral in which work, we can say, takes up a lot of energy and a lot of time, and then that time is taken away from us...” (P6, male, 55)

According to the interviewees, one of the major barriers to engaging in therapeutic exercise was the absence of explicit directions from health professionals regarding the specific exercises to be performed, their intensity and frequency. This lack of clear guidance caused confusion and uncertainty among the interviewees, who felt they were left to rely solely on their instincts.

“My doctor told me to “go for a walk”, or maybe to “move”, but never specifically, something like “it would be better in your case to do something more targeted.” (P11, male, 56)

The interviewees highlighted that the lack of determination and the hindrance in changing life habits were some of the main reasons behind not adhering to the exercise plan.

“I think so, for laziness... Because if you want to, you are able to find the time. So it is, therefore, laziness.” (P5, female, 72)

Finally, a few participants felt exercise to be valid only after surgery, therefore, not worth doing prior to total joint replacement.

“It is useless to start doing physiotherapy/exercise if I am undertaking surgery in a month.” (P9, female, 73)

In fact, they only saw exercise as applicable after surgery, with little to no utility in its own right.

“But I imagine that someone can do this... let’s call it preventive activity. Activity that can help with the recovery process following the intervention.” (P6, male, 55)

Theme 7: Developing an Uneasy Relationship with Food

Some of the participants had developed an unhealthy relationship with food. In fact, they perceived diet as a deprivation, a time-consuming and tiresome sacrifice. Moreover, some of our interviewees overeat “to eat their feelings”.

“To follow a diet is a mental fatigue [...] and eating is an easy outlet to manage the stress of daily life.” (P4, female, 47)

Furthermore, the participants considered following a diet only as a way to reduce weight on their joints without feeling its positive effects on inflammation. As a result, they tended to draw a direct association between weight and joint load.

“Of course, there is a relationship [between weight and OA]. The heavier the body, the more the knee suffers, it’s a matter of physics.” (P2, female, 68)

Sociodemographic and Economic Elements

Sociodemographic and economic elements (e.g., age, HRQoL, BMI, educational attainment etc.) may influence the OA care process. For instance, some of them (e.g., assigned sex (at birth), age, self-efficacy, educational attainment etc.) has been hypothesised to be associated with exercise adherence in conditions other than OA [8, 21, 123–126]. Poor adherence to exercise might limit its benefits [127]. Considering the rising prevalence [67] and the economic burden of OA [128], identifying factors associated with exercise adherence is fundamental to creating specific intervention programmes to improve it. Therefore, Study 3 explored the association between different lifestyle, demographic, socioeconomic and disease-related factors with adherence to the supervised exercise of an OA management programme delivered nationwide in Swedish primary care [129]. Besides, this study explored these factors' ability to explain exercise variability.

Preliminary evidence indicated that people with OA in low SEP generally experience inequalities in care [27, 28, 130, 131], and that healthcare interventions may inadvertently exacerbate pre-existing inequalities [132]. This phenomenon is known as 'intervention-generated inequalities', and it is more likely to happen with interventions targeting behavioural changes such as exercise [25, 133]. Given this, it is essential to examine the impact of the care process on inequalities to understand how to structure the management of OA. Hence, Study 4 shed some light on income-related intervention-generated inequality in the outcomes (i.e., pain intensity, arthritis-specific self-efficacy, the desire for surgery and the use of NSAIDs) of the abovementioned OA management programme.

Study 3: Factors Associated with Adherence to a Supervised Exercise Intervention for Osteoarthritis: Data from the SOAR

Exercise Adherence: Why is it Fundamental in OA?

The World Health Organisation (WHO) describes adherence as “the extent to which a person's behaviour—taking medication, following a diet and/or executing lifestyle changes [exercise], corresponds with agreed recommendations from a health care provider” [134]. In OA, poor adherence to exercise can hamper its benefits [127]. Therefore, it is fundamental to study which factors might be associated with exercise adherence to create specific interventions to improve it.

Several factors, including lifestyle, demographic characteristics, socioeconomic status, and disease-related factors, have been posited to influence exercise adherence [8, 21, 123–126]. Demographic and lifestyle factors such as female sex, higher age and BMI and low HRQoL appear to be associated with lower adherence [123, 124, 135]. Similarly, low SEPs (e.g., educational attainment and income) have been suggested to influence people's attitudes, experiences, and exposure to several health risk factors [125] and be associated with insufficient exercise levels [8, 125, 126]. Finally, different disease-related factors could impact exercise adherence. People with chronic pain seem to avoid exercise due to maladaptive coping strategies such as fear avoidance, kinesiophobia and low levels of self-efficacy [21, 136, 137].

However, all the evidence mentioned above is primarily derived from conditions other than OA, qualitative studies whose aims are not to generalise knowledge and studies with small sample sizes [8, 18, 19, 21, 123–126]. Besides, the WHO stated that adherence is influenced by the interplay of various factors, not just a single element [134]. Notwithstanding, prior studies concentrated on single factors and their average association with adherence, as evidenced by the use of measures of association such as odds ratio (OR), ignoring the variability of effects at the individual level [138]. Therefore, this study investigated the associations between lifestyle and demographic, socioeconomic and disease-related factors with adherence to the supervised exercise of an OA management programme delivered nationwide in Swedish primary care. Furthermore, we investigated these factors' ability to explain exercise adherence variability.

Study Design

This register-based study uses individual-level data from two Swedish registers, the ‘Swedish Osteoarthritis Registry’ (SOAR) and the ‘Longitudinal Integration Database for Health Insurance and Labour Market Studies’ (LISA). SOAR contains information on approximately 200,000 people with OA who participated in an OA management programme offered by the Swedish healthcare system [139]. The LISA is a valuable resource for health and labour market research. LISA is a crucial tool that allows for a better understanding of individuals' life situations in relation to the labour market, working life, and ill-health. The registry provides researchers with annual data that tracks important aspects of individuals' lives, such as educational attainment, income, occupation, and employment status by calendar year [140, 141]. The datasets were merged using unique personal identity numbers (PINs) assigned to all Swedish citizens. The research was conducted in respect of the Declaration of Helsinki and reported following the Strengthening the Reporting of Observational studies in Epidemiology (STROBE). Ethical approval was granted by the Swedish Ethics Committee (Dnr: 2019-02570).

The OA management programme in SOAR consists of two parts: education and exercise [142, 143]. The education component involves three sessions that focus on the pathophysiology of the disease and its self-management. The first two are mandatory and led by a physiotherapist, and the third session is optional and led by an OA communicator. The optional exercise component begins with a one-on-one session with a physiotherapist to tailor the programme to participants' needs and preferences and can be performed either at home or in supervised group sessions with a physiotherapist twice a week for 6-8 weeks for up to 12 sessions, following OA Swedish guidelines [144].

Variables

The ‘Levels of Adherence’ to the supervised exercise part, reported in the SOAR, is the dependent variable of this study. The collected independent variables are reported hereafter and divided as ‘Demographic and Lifestyle Characteristics’, ‘Socioeconomic Characteristics’ and ‘Disease-Related Characteristics’ (Table 4).

Table 4 Variables included in the study

| Variable | Type | Measured | Register |
|--|-------------|--|----------|
| Dependent variable | | | |
| Levels of adherence | Categorical | - Low levels of adherence (1-6 training sessions) - Medium Levels of Adherence (7-9 training sessions) - High Levels of Adherence (10-12 sessions) | SOAR |
| Independent variables | | | |
| Demographic and lifestyle factors | | | |
| Assigned sex (at birth) | Binary | - Male - Female | SOAR |
| Age | Continuous | | SOAR |
| BMI | Continuous | | SOAR |
| EQ5DVAS (HRQoL) | Continuous | VAS 0 (worst possible HRQoL) - 100 (best possible HRQoL) | SOAR |
| Socioeconomic factors | | | |
| Living Alone | Binary | - Living alone - Living with someone | LISA |
| Educational Attainment | Categorical | - Low (primary school [0–9 years]) - Medium (secondary school up to postsecondary education <3 years [10–14 years]) - High (postsecondary education [\geq 15 years]) | LISA |
| Employment | Binary | - Employed - Retired/unemployed | LISA |
| Residential area | Categorical | - Suburban - Urban | LISA |
| Individual yearly net income* | Categorical | - Lowest income quartile (< 146,500 SEK) - Second income quartile (146,501-198,100 SEK) - Third income quartile (198,111 – 278,800 SEK) - Highest income quartile (> 278,800 SEK) | LISA |
| Disease-related factors | | | |
| Index joint [139] | Binary | - Hip - Knee | SOAR |
| Numbers of painful joints [139] | Continuous | | SOAR |
| Desire for surgery [139] | Binary | - Yes - No | SOAR |

| | | | |
|--|-------------|--|------|
| Pain intensity in their index joint [145] | Continuous | VAS 0 (no pain) - 10 (unbearable pain) | SOAR |
| Pain frequency [139] | Binary | - Infrequent pain [less than every week] - Frequent pain [almost every day] | SOAR |
| Fear of movement [139] | Binary | - Yes - No | SOAR |
| Charnley score (disease that affects their walking) [146] | Categorical | - A = unilateral hip or knee OA - B = bilateral hip or knee OA - C = multiple joint OA or some other condition | SOAR |
| Self-efficacy [147, 148] | Continuous | Arthritis self-efficacy scale (ASES), 'self-efficacy pain' and 'other symptoms' subscales combined score 10 (low levels of self-efficacy) -100 (high levels of self-efficacy) | SOAR |

Legend: SOAR, Swedish Osteoarthritis Registry; VAS, Visual Analogue Scale; HRQoL; health-related quality of life; LISA, Longitudinal Integration Database for Health Insurance and Labour Market Studies; *The individual yearly net income was categorised into quartiles based on the sample income distribution.

Levels of Adherence

The 'Levels of Adherence' is a categorical variable recorded by the physiotherapists and stratified on the number of sessions the participants partook in (Low levels of adherence (1-6 training sessions) / Medium Levels of Adherence: (7-9 training sessions) / High Levels of Adherence: 10-12 sessions).

Demographic and Lifestyle Characteristics

Participants' demographic and lifestyle characteristics were reported by the participants at the baseline and recorded in the SOAR. These outcomes were 'assigned sex (at birth)' (binary variable – Male / Female), 'age' (continuous variable), body mass index ('BMI') computed from participants' height and weight (continuous variable), 'weekly physical activity' (continuous variable – hour) and 'HRQoL' (continuous variable - EuroQoL5-visual analogue scales, EQ5DVAS). The EQ5DVAS is part of the EQ-5D scale. The EQ-5D is an instrument that assesses HRQoL [149]. In the EQ5DVAS, the respondents report their perceived HRQoL on a VAS scale that scores from 0 (the worst possible HRQoL) to 100 (the best possible HRQoL).

Socioeconomic Factors

Each SEP indicator from the year before the enrolment to the SOAR register was considered for the analysis. In particular, the following SEP factors were retrieved: 'living alone' (binary variable - living alone / living with someone), 'educational attainment' (categorical variable - low (primary school [0–9 years]) / medium (secondary school up to postsecondary education <3 years [10–14 years]), / high (postsecondary education [\geq 15 years]), 'employment' (binary variable - employed / retired-unemployed), 'residential area' (categorical variable – rural / suburban / urban)

and the 'individual yearly net income' (see below for description). 'Residential area' is classified based on the 'Swedish Association of Local Authorities and Regions' (SALAR) classification of Swedish municipalities. Specifically, 'rural' areas are smaller towns/urban areas and rural municipalities, 'suburban' areas are medium-sized towns ($\geq 40,000$ inhabitants) and municipalities near medium-sized towns, 'urban' areas are large cities ($\geq 200,000$ inhabitants) and municipalities near large cities [150]. The individual yearly net income was categorised into quartiles based on the sample income distribution: lowest income quartile ($< 146,500$ SEK) / second income quartile ($146,501-198,100$ SEK) / third income quartile ($198,111 - 278,800$ SEK) / highest Income Quartile ($> 278,800$ SEK) [150].

Disease-Related Characteristics

The physiotherapists recorded the 'index joint' (categorical variable - hip or knee) [139], namely, the joint with OA. They assessed this variable based on the participant's medical history, symptoms, and clinical assessment. In the case of multiple joints with OA, the most symptomatic joint was considered the index joint for the treatment. The participants self-recorded the 'number of painful joints' (continuous variable), their 'desire for surgery' (binary variable - yes/no) that was assessed by asking them: "Are your knee/hip symptoms so severe that you wish to undergo surgery?" [139]; their 'pain intensity' (continuous variable 0-10, Numeric Rating Scale (NRS) [145]) in their 'index Joint'; their 'pain frequency' (binary variable – infrequent pain [less than every week] / frequent pain [almost every day]) that was assessed with the question: "How often do you have pain in your knee/hip" [139]; their 'fear of movement' (binary variable – yes / no) that was assessed with the question "Are you afraid your joints will be injured by physical training/activity?"; the 'Charnley score' (categorical variable – Charnley score; A = unilateral hip or knee OA / B = bilateral hip or knee OA / C = multiple joint OA or some other condition) that categorises people with OA into three classes based on the disease(s) that affect walking ability [146]; and the 'arthritis-specific self-efficacy' (continuous variable 10-100 – pain and symptom arthritis self-efficacy scale, ASES) the Swedish version of the scale was adopted [147]. The ASES scale is a reliable instrument that assesses patients' arthritis-specific self-efficacy, namely, their beliefs about their ability to perform a specific task and cope with OA [148]. The full version is composed of three subscales: 1) 'self-efficacy pain scale' (5 items); 2) 'function scale' (9 items); 3) 'other symptoms scale' (6 items). Participants indicate to what extent they feel confident they can do the different tasks reported in the items from 10 ('very uncertain') to 100 ('very certain'). In the SOAR, only 1) and 3) were adopted and combined as suggested in the scale instruction [148]. As per the psychometric properties of the Swedish ASES, the three factor structure was confirmed. Cronbach's alpha for internal consistency ranged between 0.82-0.91 and test-retest correlations between 0.81-0.91, similarly to the original version [148], showing that the Swedish ASES met satisfactorily psychometric standards. Similar results are reported in the two subscales adopted with a Cronbach's alpha of 0.92 for the 'self-efficacy pain scale' and 0.82 for the 'other symptoms' scale [147].

Participants and Recruitment

Our cohort gathered all the participants in the SOAR with a first registration (baseline) between 2012 and 2015. We included only those who started the exercise group sessions supervised by the physiotherapists after the initial encounter. We selected participants with knee or hip OA who were recorded in the SOAR only once.

Specifically, between 1st January 2012 and 31st December 2015, 46,905 people with OA were recorded in the SOAR. However, we excluded $n=7$ participants as they had joints other than the hip and knee as their first cause of pain, $n=27,147$ as they did not perform any supervised exercise session(s) and $n=1$ because they attended the programme more than once. Hence, among them, only 19,750 (73% Female sex; age: 67 (SD: 8.9)) participants with the knee (69%) or hip (31%) OA started the supervised exercise sessions. Figure 6 reports the participants' selection process.




Figure 6 Participants' selection

Table 5 presents the characteristics of the whole cohort and stratified by the levels of adherence. Specifically, 5,862 (30%) reached a low level of adherence, 3,947 (20%) a medium level and 9,941 (50%) a high level.

Table 5 Descriptive Statistics

| Variables | Total Sample (n=19,750) | Low Levels of Adherence (n=5,862) | Medium Levels of Adherence (n=3,947) | High Levels of Adherence (n=9,941) |
|--|----------------------------|---|--|--|
| Demographic and Lifestyle Characteristics | | | | |
| Assigned sex (at birth) | n=19,750 | n=5,862 | n=3,947 | n=9,941 |
| Male, n(%) | 5,421 (27.45) | 1,519 (25.91) | 925 (23.44) | 2,977 (29.95) |
| Female, n(%) | 14,329 (72.55) | 4,343 (74.09) | 3,022 (76.65) | 6,964 (70.05) |
| Age | n=19,750 | n=5,862 | n=3,947 | n=9,941 |
| Mean(SD) | 66.86 (8.94) | 65.87 (9.39) | 66.47 (9.01) | 67.60 (8.57) |
| BMI | n=19,381 | n=5,735 | n=3,867 | n=9,779 |
| Mean(SD) | 27.56 (4.76) | 27.73 (4.90) | 27.75 (4.89) | 27.43 (4.63) |
| HRQoL (EQ5DVAS, 0-100) | n=17,933 | n=5,317 | n=3,592 | n=9,024 |
| Mean(SD) | 65.82 (19.22) | 65.84 (19.37) | 65.74 (19.35) | 65.85 (19.07) |
| Weekly physical activity (hour) | n=18,050 | n=5,364 | n=3,606 | n=9,080 |
| Mean(SD) | 4.11 (2.53) | 4.14 (2.53) | 4.03 (2.49) | 4.13 (2.54) |
| Socioeconomic Characteristics | | | | |
| Educational attainment (n) | n=19,699 | n=5,862 | n=3,938 | n=9,918 |
| Low, n(%) | 4,331 (21.99) | 1,170 (20.02) | 795 (20.19) | 2,366 (23.86) |
| Medium, n(%) | 9,843 (49.97) | 2,962 (50.69) | 2,007 (50.96) | 4,874 (49.14) |
| High, n(%) | 5,525 (28.05) | 1,711 (29.28) | 1,136 (28.85) | 2,678 (27.00) |
| Income – quartile (n) | n=19,738 | n=5,858 | n=3,945 | n=9,935 |
| Lowest income quartile, n(%) | 4,942 (25.04) | 1,345 (22.96) | 1,022 (25.91) | 2,575 (25.92) |
| Second income quartile, n(%) | 4,936 (25.01) | 1,393 (23.78) | 982 (24.89) | 2,561 (25.78) |
| Third income quartile, n(%) | 4,929 (24.97) | 1,517 (25.90) | 976 (24.74) | 2,436 (24.52) |
| Highest income quartile, n(%) | 4,931 (24.98) | 1,603 (27.36) | 965 (24.46) | 2,363 (23.78) |
| Area of living (n) | n=19,738 | n=5,858 | n=3,945 | n=9,935 |
| Rural, n(%) | 6,047 (30.64) | 1,667 (28.46) | 1,180 (29.91) | 3,200 (32.21) |
| Suburban, n(%) | 8,252 (41.81) | 2,435 (41.57) | 1,708 (43.30) | 4,109 (41.36) |

| | | | | |
|--|----------------|---------------|---------------|---------------|
| Urban, n(%) | 5,439 (27.56) | 1,756 (29.98) | 1,057 (26.79) | 2,626 (26.43) |
| Employment (n) | n=19,738 | n=5,858 | n=3,945 | n=9,935 |
| Unemployed, n(%) | 12,244 (62.03) | 3,275 (55.91) | 2,394 (60.68) | 6,575 (66.18) |
| Employed, n(%) | 7,494 (37.97) | 2,583 (44.09) | 1,551 (39.32) | 3,360 (33.82) |
| Living alone (n) | n=19,738 | n=5,858 | n=3,945 | n=9,935 |
| Living alone, n(%) | 7,754 (39.28) | 2,411 (41.16) | 1,457 (36.93) | 3,886 (39.11) |
| Living with someone, n(%) | 11,984 (60.72) | 3,447 (58.84) | 2,488 (63.07) | 6,049 (60.89) |
| Disease-Related Characteristics | | | | |
| Worst joint (n) | | n=5,862 | n=3,947 | n=9,941 |
| Hip, n(%) | 6,049(30.63) | 1,708 (29.14) | 1,188 (30.10) | 3,153 (31.72) |
| Knee, n(%) | 13,701 (69.37) | 4,154 (70.86) | 2,759 (69.90) | 6,788 (68.28) |
| Pain intensity (NRS 0-10) | n=19,686 | n=5,843 | n=3,935 | n=9,908 |
| Mean(SD) | 5.25 (1.83) | 5.23 (1.85) | 5.24 (1.87) | 5.26 (1.80) |
| Pain frequency (n) | n=19,700 | n=5,842 | 3,940 | n=9,918 |
| Infrequent, n(%) | 3,436 (17.44) | 1,100 (18.83) | 723 (18.35) | 1,613 (16.26) |
| Frequent, n(%) | 16,264 (82.56) | 4,742 (81.17) | 3,217 (81.65) | 8,305 (87.34) |
| Number of painful joints | n=19,750 | n=5,862 | n=3,947 | n=9,941 |
| Mean(SD) | 1.94 (1.29) | 1.95 (1.28) | 2.00 (1.32) | 1.91 (1.27) |
| Charnley score (n) | n=19,735 | n=5,855 | n=3,946 | n=9,934 |
| A, n(%) | 6,814 (34.53) | 2,000 (34.16) | 1,340 (33.96) | 3,474 (34.97) |
| B, n(%) | 3,437 (17.42) | 1,009 (17.23) | 686 (17.38) | 1,742 (17.54) |
| C, n(%) | 9,484 (48.06) | 2,946 (48.61) | 1,920 (48.66) | 4,718 (47.49) |
| Walking difficulties (n) | n=19,651 | n=5,835 | n=3,932 | n=9,884 |
| No, n(%) | 3,472 (17.67) | 1,105 (18.94) | 731 (18.59) | 1,636 (16.55) |
| Yes, n(%) | 16,179 (82.33) | 4,730 (81.06) | 3,201 (81.41) | 8,248 (83.45) |
| Fear of movement (n) | n=19,651 | n=5,821 | n=3,928 | n=9,902 |
| No, n(%) | 16,562 (84.28) | 4,871 (83.68) | 3,303 (84.09) | 8,388 (84.71) |
| Yes, n(%) | 3,089 (15.72) | 950 (16.32) | 625 (15.91) | 1,514 (15.29) |
| Desire for surgery (n) | n= 19,558 | n=5,798 | n=3,906 | n=9,854 |

| | | | | |
|---------------------------------------|----------------|---------------|---------------|---------------|
| No, n(%) | 14,936 (76.37) | 4,441 (76.60) | 3,017 (77.24) | 7,478 (75.89) |
| Yes, n(%) | 4,622 (23.63) | 1,357 (23.40) | 889 (22.76) | 2,376 (24.11) |
| Self-efficacy (ASES 10-100) | n=19,149 | n=5,660 | n=3,834 | n=9,655 |
| Mean(SD) | 65.54 (16.43) | 65.44 (16.54) | 65.51 (16.62) | 65.61 (16.28) |

Legend: n, number; SD, standard deviation; HRQoL, health-related quality of life; EQ5DVAS, EuroQol-5D Health Visual Analogue Scale; IQ, income quartile; OA, osteoarthritis; NRS, numeric rating scale; ASES, Arthritis Self-Efficacy Scale.

Factors Associated with Exercise Adherence – Average Association

A multivariable exploratory analysis was performed to identify which independent variables were independently associated with exercise adherence in the SOAR [151]. Multivariable exploratory analyses detect patterns and identify relationships between independent variables and outcomes [151–153]. Stata 17 was used for the analysis.

Since the proportional odds assumption was not met, an ordered logistic regression could not be performed. Hence, we ran a multinomial logistic regression with a listwise deletion (stata function ‘mlogit’) to determine the association between the independent variables and the adherence to exercise. No missing data were reported in the outcome (adherence). Less than 1% of the data on 'socioeconomic' characteristics was missing, primarily due to an error during the data upload process in ‘LISA’. ‘Demographic and lifestyle’ and ‘disease-related’ characteristics missing data in the SOAR are most likely a result of a mistake by the physiotherapists responsible for uploading the data at the local unit. Hence, missing data in both registers could be considered missing completely at random, introducing no or minimal bias in our analysis. Table 6 reports the descriptive statistics of the total sample and the samples included in and excluded from the analysis, highlighting minimal differences between them.

Table 6 Descriptive statistics of the total sample and the samples included in and excluded from the analysis

| Variables | Total Sample (n=19,750) | Included in the Analysis (n=16,685) | Excluded from Analysis (n=3,065) |
|--|-----------------------------------|---|--|
| Demographic and Lifestyle Characteristics | | | |
| Assigned sex (at birth) | n=19,750 | n=16,685 | n=3,065 |
| Male, n(%) | 5,421 (27.45) | 4,662 (27.94) | 759 (24.76) |
| Female, n(%) | 14,329 (72.55) | 1,2023 (72.06) | 2,306 (75.24) |
| Age | n=19,750 | n=16,685 | n=3,065 |
| Mean(SD) | 66.86 (8.94) | 66.79 (8.90) | 67.24 (9.15) |
| BMI | n=19,381 | n=16,685 | n=2,696 |
| Mean(SD) | 27.56 (4.76) | 27.57 (4.76) | 27.50 (4.81) |
| HRQoL (EQ5DVAS, 0-100) | n=17,933 | n=16,685 | n=1,248 |
| Mean(SD) | 65.82 (19.22) | 65.87 (19.15) | 65.10 (20.02) |
| Weekly physical activity (hour) | n=18,050 | n=16,685 | n=1,365 |
| Mean(SD) | 4.11 (2.53) | 4.13 (2.52) | 3.87 (2.56) |
| Socioeconomic Characteristics | | | |
| Educational attainment (n) | n=19,699 | n=16,685 | n=3,014 |
| Low, n(%) | 4,331 (21.99) | 3,581 (21.46) | 750 (24.88) |
| Medium, n(%) | 9,843 (49.97) | 8,382 (50.24) | 1,461 (48.47) |
| High, n(%) | 5,525 (28.05) | 4,722 (28.30) | 803 (26.64) |
| Income – quartile (n) | n=19,738 | n=16,685 | n=3,053 |
| Lowest income quartile, n(%) | 4,942 (25.04) | 4,018 (24.08) | 924 (30.27) |
| Second income quartile, n(%) | 4,936 (25.01) | 4,163 (24.95) | 773 (25.32) |
| Third income quartile, n(%) | 4,929 (24.97) | 4,187 (25.09) | 742 (24.30) |
| Highest income quartile, n(%) | 4,931 (24.98) | 4,317 (25.87) | 614 (20.11) |
| Area of living (n) | n=19,738 | n=16,685 | n=3,053 |
| Rural, n(%) | 6,047 (30.64) | 5,150 (30.87) | 897 (29.38) |
| Suburban, n(%) | 8,252 (41.81) | 6,818 (40.86) | 1,434 (46.97) |
| Urban, n(%) | 5,439 (27.56) | 4,717 (28.27) | 722 (23.65) |
| Employment (n) | n=19,738 | n=16,685 | n=3,053 |
| Unemployed, n(%) | 12,244 (62.03) | 10,243 (61.39) | 2,001 (65.54) |
| Employed, n(%) | 7,494 (37.97) | 6,442 (38.61) | 1,052 (34.46) |
| Living alone (n) | n=19,738 | n=16,685 | n=3,053 |
| Living alone, n(%) | 7,754 (39.28) | 6,522 (39.09) | 1,232 (40.35) |
| Living with someone, n(%) | 11,984 (60.72) | 10,163 (60.91) | 1,821 (59.65) |
| Disease-Related Characteristics | | | |

| | | | |
|---|----------------|----------------|---------------|
| Worst joint (n) | n=19,759 | n=16,685 | n=3947 |
| Hip, n(%) | 6,049(30.63) | 5,129 (30.74) | 920 (30.02) |
| Knee, n(%) | 13,701 (69.37) | 11,556 (69.26) | 2,145 (69.98) |
| Pain intensity (NRS 0-10) | n=19,686 | n=16,685 | n=3,001 |
| Mean(SD) | 5.25 (1.83) | 5.25 (1.82) | 5.24 (1.86) |
| Pain frequency (n) | n=19,700 | n=16,685 | n=3,015 |
| Infrequent, n(%) | 3,436 (17.44) | 2,957 (17.72) | 479 (15.89) |
| Frequent, n(%) | 16,264 (82.56) | 13,728 (82.28) | 2,536 (84.11) |
| Number of painful joints | n=19,750 | n=16,685 | n=3,065 |
| Mean(SD) | 1.94 (1.29) | 2.11 (1.21) | 1.03 (1.32) |
| Charnley score (n) | n=19,735 | n=16,685 | n=3,050 |
| A, n(%) | 6,814 (34.53) | 5,745 (34.43) | 1,069 (35.05) |
| B, n(%) | 3,437 (17.42) | 2,705 (16.21) | 732 (24.00) |
| C, n(%) | 9,484 (48.06) | 8,235 (49.36) | 1,249 (40.95) |
| Walking difficulties (n) | n=19,651 | n=16,685 | n=2,966 |
| No, n(%) | 3,472 (17.67) | 2,942 (17.63) | 530 (17.87) |
| Yes, n(%) | 16,179 (82.33) | 13,743 (82.37) | 2,436 (82.13) |
| Fear of movement (n) | n=19,651 | n=16,685 | n=2,966 |
| No, n(%) | 16,562 (84.28) | 14,088 (84.44) | 2,474 (83.41) |
| Yes, n(%) | 3,089 (15.72) | 2,597 (15.56) | 492 (16.59) |
| Desire for surgery (n) | n= 19,558 | n=16,685 | n=2,873 |
| No, n(%) | 14,936 (76.37) | 12,738 (76.34) | 2,198 (76.51) |
| Yes, n(%) | 4,622 (23.63) | 3,947 (23.66) | 675 (23.49) |
| Arthritis-specific self-efficacy | | | |
| (ASES Pain and Symptoms, 0-100) | n=19,149 | n=16,685 | n=2,464 |
| Mean(SD) | 65.54 (16.43) | 65.74 (16.42) | 64.18 (16.38) |

Legend: n, number; SD, standard deviation; HRQoL, health-related quality of life; EQ5DVAS, EuroQol-5D Health Visual Analogue Scale; OA, osteoarthritis; NRS, numeric rating scale; ASES, Arthritis Self-Efficacy Scale.

Variable selection in the model was informed by previous literature on exercise adherence in other chronic pain conditions [8, 21, 123–126] and the evidence for action on adherence by the WHO [134]. Then, the variables were clustered in ‘demographic and lifestyle’, ‘socioeconomic and ‘disease-related’ following the dimensions proposed by the WHO [134]. The multicollinearity assumption between continuous variables was tested, and none of the continuous variables was highly correlated. The relative risk ratio (RRR) of being in ‘medium level of adherence’ or ‘high level of adherence’ with respect to ‘low level of adherence’ and 95% confidence intervals (CIs) were estimated for each covariate in the model. For the variables ‘HRQoL’ and ‘arthritis-specific self-efficacy’,

the RRR is presented as a 10-point change in these scales. Hence, After the listwise deletion, the multinomial logistic regression was run on n=16,685 (85%) with low levels of adherence as the reference category (Table 7).

Table 7 Association between exercise adherence and investigated variables

| Variables n=16,685 | p-value | Relative Risk Ratio | 95% C.I. for EXP(B) | |
|--|---------|------------------------|---------------------|-------|
| | | | Lower | Upper |
| Low Levels of Adherence | | | | |
| Medium Levels of Adherence | | | | |
| Assigned sex (at birth) | | | | |
| Male | | (Base Category) | | |
| Female | 0.03 | 1.13 | 1.02 | 1.27 |
| Age | 0.14 | 1.00 | 0.99 | 1.01 |
| BMI | 0.37 | 0.99 | 0.99 | 1.01 |
| HRQoL (EQ5DVAS, 0-100)* | 0.57 | 0.99 | 0.97 | 1.02 |
| Weekly physical activity (hour) | 0.02 | 0.98 | 0.96 | 0.99 |
| Educational attainment | | | | |
| Low | | (Base Category) | | |
| Medium | 0.88 | 0.99 | 0.88 | 1.12 |
| High | 0.63 | 0.97 | 0.84 | 1.11 |
| Income – quartile | | | | |
| Lowest income quartile | | (Base Category) | | |
| Second income quartile | 0.71 | 0.98 | 0.86 | 1.11 |
| Third income quartile | 0.63 | 0.97 | 0.84 | 1.11 |
| Highest income quartile | 0.41 | 0.94 | 0.81 | 1.09 |
| Area of living | | | | |
| Rural | | (Base Category) | | |
| Suburban | 0.27 | 0.94 | 0.85 | 1.05 |
| Urban | 0.02 | 0.87 | 0.78 | 0.98 |
| Employment | | | | |
| Unemployed | | (Base Category) | | |
| Employed | <0.01 | 0.82 | 0.72 | 0.93 |
| Living alone | | | | |
| Living alone | | (Base Category) | | |
| Living with someone | <0.01 | 1.21 | 1.10 | 1.32 |
| Worst joint | | | | |

| | | | | | |
|---|-----------------|------|------|------|--|
| Hip | (Base Category) | | | | |
| Knee | 0.35 | 0.95 | 0.86 | 1.05 | |
| Pain intensity (NRS 0-100) * | 0.49 | 1.01 | 0.98 | 1.04 | |
| Pain frequency | | | | | |
| Infrequent | (Base Category) | | | | |
| Frequent | 0.80 | 0.98 | 0.87 | 1.11 | |
| Number of painful joints | 0.01 | 1.06 | 1.01 | 1.10 | |
| Charnley score | | | | | |
| A | (Base Category) | | | | |
| B | 0.99 | 0.99 | 0.97 | 1.15 | |
| C | 0.13 | 0.91 | 0.81 | 1.03 | |
| Walking difficulties | | | | | |
| No | (Base Category) | | | | |
| Yes | 0.93 | 0.99 | 0.88 | 1.13 | |
| Fear of movement | | | | | |
| No | (Base Category) | | | | |
| Yes | 0.49 | 1.04 | 0.92 | 1.18 | |
| Desire for surgery | | | | | |
| No | (Base Category) | | | | |
| Yes | 0.26 | 0.94 | 0.83 | 1.05 | |
| Arthritis-specific self-efficacy | | | | | |
| (ASES Pain and Symptoms, 0-100)* | 0.29 | 1.02 | 0.99 | 1.05 | |

High Levels of Adherence

| | | | | | |
|---------------------------------------|-----------------|------|------|------|--|
| Assigned sex (at birth) | | | | | |
| Male | (Base Category) | | | | |
| Female | <0.01 | 0.82 | 0.75 | 0.89 | |
| Age | <0.01 | 1.01 | 1.01 | 1.02 | |
| BMI | 0.01 | 0.99 | 0.98 | 0.99 | |
| HRQoL (EQ5DVAS, 0-100)* | 0.18 | 0.98 | 0.96 | 1.01 | |
| Hours physical activity / week | 0.79 | 0.99 | 0.98 | 1.01 | |
| Educational attainment | | | | | |
| Low | (Base Category) | | | | |
| Medium | 0.02 | 0.89 | 0.81 | 0.98 | |
| High | <0.01 | 0.84 | 0.76 | 0.94 | |
| Income – quartile | | | | | |
| Lowest income quartile | (Base Category) | | | | |
| Second income quartile | 0.79 | 1.01 | 0.91 | 1.13 | |

| | | | | |
|---|-----------------|------|------|------|
| Third income quartile | 0.61 | 1.03 | 0.92 | 1.15 |
| Highest income quartile | 0.95 | 1.00 | 0.89 | 1.14 |
| Area of living | | | | |
| Rural | (Base Category) | | | |
| Suburban | <0.01 | 0.79 | 0.73 | 0.86 |
| Urban | <0.01 | 0.78 | 0.71 | 0.86 |
| Employment | | | | |
| Unemployed | (Base Category) | | | |
| Employed | <0.01 | 0.71 | 0.64 | 0.78 |
| Living alone | | | | |
| Living alone | (Base Category) | | | |
| Living with someone | 0.29 | 1.04 | 0.97 | 1.12 |
| Worst joint | | | | |
| Hip | (Base Category) | | | |
| Knee | 0.03 | 0.92 | 0.85 | 0.99 |
| Pain intensity (NRS 0-10) | 0.12 | 1.02 | 0.99 | 1.04 |
| Pain frequency | | | | |
| Infrequent | (Base Category) | | | |
| Frequent | 0.02 | 1.13 | 1.02 | 1.25 |
| Number of painful joints | 0.50 | 1.01 | 0.98 | 1.05 |
| Charnley score | | | | |
| A | (Base Category) | | | |
| B | 0.74 | 1.02 | 0.91 | 1.14 |
| C | 0.11 | 0.93 | 0.84 | 1.02 |
| Walking difficulties | | | | |
| No | (Base Category) | | | |
| Yes | 0.03 | 1.12 | 1.01 | 1.24 |
| Fear of movement | | | | |
| No | (Base Category) | | | |
| Yes | 0.93 | 1.00 | 0.91 | 1.11 |
| Desire for surgery | | | | |
| No | (Base Category) | | | |
| Yes | 0.44 | 0.96 | 0.88 | 1.06 |
| Arthritis-specific self-efficacy | | | | |
| (ASES Pain and Symptoms, 0-100)* | <0.01 | 1.04 | 1.02 | 1.07 |

Legend: n, number; EQ5DVAS, EuroQol-5D Health Visual Analogue Scale; IQ, income quartile; *, RRR are reported as an increase of 10 points in the scale; OA, osteoarthritis; NRS, numeric rating scale; ASES, Arthritis Self-Efficacy Scale.

We found that ‘female’ sex (RRR=1.13, 95% CI [1.02-1.27]), ‘living with someone’ (1.21, [1.10-1.32]) and an increase of one ‘number of joints with OA’ (1.06, 95% CI [1.01-1.10]) were positively associated with achieving medium levels of adherence. Conversely, an increase in an hour of ‘weekly physical activity’ (0.98, 85% CI [0.96-0.99]), living in an ‘urban’ area (0.87, 95% CI [0.78-0.98]) and being ‘employed’ (0.82, 95% CI [0.72-0.93]) were negatively associated with achieving medium levels of adherence.

An increase of one year in ‘age’ (1.01, 95% CI [1.01-1.02]), having ‘frequent’ pain (1.13, 95% CI [1.02-1.25]), having ‘walking difficulties’ (1.12, 95% CI [1.01-1.24]) and with a 10-point increase on the ‘arthritis-specific self-efficacy’ scale (1.04, 95% CI [1.02-1.07]) were positively associated with high levels of adherence. Instead, ‘female’ sex (0.82, 95% CI [0.75-0.89]), an increase of one point in ‘BMI’ (0.99, 95% CI [0.98-0.99]), living in a ‘suburban’ (0.79, 95% CI [0.73-0.86]) or ‘urban’ area (0.78 95% CI [0.71-0.86]), being ‘employed’ (OR=0.71, 95% CI [0.64-0.78]), having a medium (0.89, 95% CI [0.81; 0.98]) or a high ‘educational attainment’ (0.84, 95% CI [0.76-0.94]) and having ‘knee’ as ‘worst joint’ (0.92, 95% CI [0.85-0.99]) were negatively associated with high levels of adherence.

Factors Associated with Exercise Adherence – Variability Explanation

The McFadden R² (stata function ‘fitstat’) was employed to assess the capacity of the models to explain the variability of exercise adherence. This metric indicates the extent to which the variance of the dependent variable (adherence) can be explained by the independent variables in the model using a 0-100% scale. The McFadden R² was calculated for the full model, which included all variables. Subsequently, a set of variables (demographic and lifestyle, socioeconomic and disease-related) were excluded from the model, and the difference in McFadden R² was calculated to determine the contribution of each set of variables to the explanatory power of the full model.

The McFadden R² of the full model indicated that the participant's demographic and lifestyle characteristics, socioeconomic characteristics, and disease-related characteristics collectively explained approximately 1.2% of the variation in adherence. When each variable set was removed from the model, there was a difference in McFadden R² with respect to the full model of 0.3%, 0.4%, and 0.2%, respectively. Of these, disease-related characteristics contributed the most to the explanatory power of the full model, although the overall explanatory ability was low.

Study 4: Income-Related Health Inequality and Osteoarthritis Intervention Outcomes: a Cohort Study

Inequalities in OA Care

SEPs are central determinants of health [154, 155]. They are defined as the position of an individual within a society stemming from different factors such as educational attainment and income [154, 156, 157]. The uneven distribution of health or health resources between individuals or groups due to genetic or other factors such as SEPs is defined as ‘health inequality’ [158]. It is worth noticing that this term does not consider whether this difference is just or unjust. Therefore, it notices that there is a difference among individuals or groups. Conversely, unfair, avoidable differences arising from poor governance, corruption or cultural exclusion are defined as ‘health inequity’ [159].

In OA, preliminary findings indicated that people in lower SEP experienced health inequalities in pain intensity, self-efficacy and desire for surgery [27, 28, 130, 131]. Moreover, healthcare interventions can widen pre-existing inequalities gaps between people in different SEPs [132]. This phenomenon is known as ‘intervention-generated inequalities’, and it is more likely to happen with healthcare interventions targeting behavioural changes, such as exercise and education [25, 133]

However, as per the adherence to exercise, these studies on inequalities focused on measures of average association [27, 28, 130, 131]. In contrast, one of the definitions of inequalities reported by the WHO defined inequalities as differences in the distribution of health determinants among different socioeconomic groups (rather than the average association thereof) [160]. Besides, these studies did not investigate inequalities regarding the use of non-steroidal anti-inflammatory (NSAIDs) drugs, though broadly used in OA treatment, despite their adverse-associated events [161]. Finally, these studies did not explore intervention-generated inequalities and adopted SEPs (e.g., education) other than income.

Income

Income is a straightforward measure of social class that plays a critical role in maintaining people's health [162]. This role is twofold: a direct effect on the material resources needed for survival, and an indirect impact on social opportunities and ability to cope with life events [162]. In healthcare, income suggested a ‘dose-response’ association with health [163, 164]. The mechanisms through which income could affect health are several. It allows for 1) buying access to higher-quality material resources (e.g., food); 2) increasing access to service (e.g., health services); 3) providing materials that are ‘relevant’ for society, fostering individuals’ self-esteem and social standing

[164]. However, income is difficult to gather as people might refrain from providing this information [164]. Moreover, disposable income might be more interesting to collect than gross income as the latter represents how much a(n) individual / family can spend [164]. Nevertheless, gross income is mainly collected.

This study investigated income-related inequality in the outcomes (i.e., pain intensity, arthritis-specific self-efficacy, the desire for surgery and the use of NSAIDs) of the SOAR, using concentration index analysis and income information from LISA [163]. LISA allowed for having accurate individuals' disposable income. Specifically, we analysed how the concentration index changed throughout the different time points in the SOAR to see whether 'intervention-generated inequalities' arose.

Study Design

A cohort study on individual-level register data. The SOAR (for data on the OA first-line intervention) was merged with LISA (for income information) using personal identity numbers (PINs) unique to all citizens in Sweden. The research was conducted in respect of the Declaration of Helsinki and reported following the Strengthening the Reporting of Observational studies in Epidemiology (STROBE). Ethical approval was granted by the Swedish Ethics Committee (Dnr: 2019-02570).

Variables

Descriptive Characteristics

Participants reported their demographic characteristics at the baseline. Specifically, they reported their 'Assigned sex (at birth)' (categorical variable – Male / Female), 'Age' (continuous variable), and height and weight, which were merged into the 'BMI' (continuous variable). The physiotherapists established participants' 'Index Joint' (categorical variable - hip or knee) based on medical history, symptoms, and clinical assessment. In the case of multiple joints with OA, the most symptomatic joint was considered the index joint for the treatment. At the inspections of q-q plots (Stata command 'qnorm'), continuous variables followed a normal distribution and were reported as mean \pm standard deviation (SD). Categorical variables are reported as absolute and percentage frequencies.

OA Outcomes

The OA outcomes were retrieved from the SOAR and reported by the participants and the physiotherapists. We retrieved mean 'Pain intensity' during the last week (continuous variable) in patients' 'Index joint' (continuous variable 0-10, Numeric Rating Scale (NRS) [145]) and their 'Desire for surgery' (binary variable - yes / no) assessed by asking them: "Are your knee / hip symptoms so severe that you wish to undergo surgery?" [139], and the 'Self-efficacy' with the 'Arthritis self-efficacy scale

(continuous variable 10-100 – pain and symptom arthritis self-efficacy scale, ASES). The ASES scale is a reliable instrument that assesses patients’ arthritis-specific self-efficacy, namely, their beliefs about their ability to perform a specific task and cope with OA [148]. The full version is composed of three subscales: 1) ‘self-efficacy pain scale’ (5 items); 2) ‘function scale’ (9 items); 3) ‘other symptoms scale’ (6 items). Participants indicate to what extent they feel confident they can do the tasks reported in the items from 10 (‘very uncertain’) to 100 (‘very certain’). In the SOAR, only 1) and 3) were adopted and combined as suggested in the scale instruction [148]. Cronbach’s alpha for internal consistency ranged between 0.82-0.91 and test-retest correlations between 0.81-0.91, similar to the original version [148], showing that the Swedish ASES met satisfactorily psychometric standards. Similar results are reported in the two subscales adopted with a Cronbach’s alpha of 0.92 for the ‘self-efficacy pain scale’ and 0.82 for the ‘other symptoms’ scale [147]. This scale was used from 2012 to 2015. Participants report all these variables. The physiotherapists collect the ‘Use of NSAIDs’. They reported whether patients had taken any medications for their joint pain in the last three months and whether or not they were NSAIDs (binary variable - yes / no) [139]. All these variables but ‘Use of NSAIDs’ were measured at baseline, 3- and 12-month follow-ups, that was collected at the baseline and 3-month follow-up.

Income

‘Individuals’ disposable income’ (continuous variable) in the year before the enrolment to the SOAR was retrieved from LISA and considered for the analysis. The disposable income is the part of an individual’s income used for saving or consumption. Specifically, disposable income considers income from employment, social welfare, pension (both public and private), sickness benefits, income from business activities and capital minus taxes and deductions and several other benefits. The household’s disposable income per consumption unit is calculated by dividing the sum of all family members’ disposable incomes by the total consumption weight of the family. Specifically, one adult weighs 1.0, two adults as 1.51 and children as 0.56–0.76 (depending on age) in the household [165]. The income is reported in SEK (10 SEK ≈ 1 €).

Participants and Recruitment

Our cohort gathers all the participants in the SOAR with a first registration (baseline) between 2008 and 2018. Specifically, we selected participants with knee or hip OA as the primary cause of their pain, who were recorded in the SOAR only once and did not perform the programme digitally. Hence, we identified 126,308 participants, of whom 7,639 reported joints other than the hip and knee as their first cause of pain and were excluded. Moreover, we excluded 2,663 who attended exercises digitally and 603 that participated in the programme more than once. Finally, 115,403 (age: 66.2 years (SD: 9.7); sex: female 67.8%) people with knee and hip OA were identified (Figure 7). Table 8 presents the characteristics of the whole cohort. Table 9 reports the characteristics of those with and without missing values in the OA outcomes of interest.




Figure 7 Selection of the study population

Table 8 Descriptive statistics of the included population

| Variables | Baseline | 3-months | 12-months |
|----------------------------------|---------------------|---------------|---------------|
| Descriptive Variables | | | |
| Assigned sex (at birth) | n=115,403 | | |
| Female, n (%) | 78,233 (67.8) | | |
| Male, n (%) | 37,170 (32.2) | | |
| Age | n=115,403 | | |
| Mean (SD) | 66.2 (9.7) | | |
| BMI | n=107,354 | | |
| Mean (SD) | 27.7 (5.0) | | |
| Worst joint (n) | n=115,402 | | |
| Hip, n (%) | 37,212 (32.4) | | |
| Knee, n (%) | 78,191 (67.8) | | |
| Income | n=115,356 | | |
| Mean (SD) | 241,414.2 (271,223) | | |
| OA Outcomes | | | |
| Pain intensity (NRS 0-10) | n=114,908 | n=80,588 | n=47,648 |
| Mean (SD) | 5.5 (2.1) | 4.4 (2.3) | 4.4 (2.3) |
| Arthritis self-efficacy | | | |
| (ASES Pain and Symptoms, 10-100) | n=45,286 | n=33,465 | n=26,074 |
| Mean (SD) | 64.8 (17.0) | 69.1 (17.7) | 65.0 (18.8) |
| Desire for surgery (n) | n=113,713 | n=79,425 | n=47,246 |
| No, n (%) | 82,327 (72.4) | 62,027 (78.1) | 36,907 (78.1) |
| Yes, n (%) | 31,386 (27.6) | 17,398 (21.9) | 10,339 (21.9) |
| Use of NSAIDs (n) | n=113,560 | n=80,428 | |
| No, n (%) | 64,460 (56.8) | 54,890 (68.3) | |
| Yes, n (%) | 49,100 (43.2) | 25,538 (31.2) | |

Abbreviations: n, number; SD, standard deviation, ASES, Arthritis-Specific Self-Efficacy; NSAIDs, non-steroidal anti-inflammatory drugs.

Table 9 Descriptive statistics per OA outcomes confronting complete VS missing cases

| Variables | Baseline VS 3-months | | Baseline VS 12-months | |
|--------------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| | Complete cases | Missing cases | Complete Cases | Missing cases |
| Pain Intensity | | | | |
| | n=80,329 | n=35,074 | n=47,499 | N=67,904 |
| Assigned sex (at birth) | | | | |
| Female, n(%) | 54,972 (68.43) | 23,261 (66.32) | 33,270 (70.04) | 44,963 (66.22) |
| Male, n(%) | 25,357 (31.57) | 11,813 (33.68) | 14,229 (29.96) | 22,941 (33.78) |
| Age | | | | |
| Mean (SD) | 67.01 (9.35) | 65.96 (10.32) | 66.76 (9.15) | 66.64 (10.02) |
| BMI | | | | |
| Mean (SD) | 27.58 (4.84) | 27.90 (5.22) | 27.44 (4.80) | 27.85 (5.07) |
| Worst joint | | | | |
| Hip, n(%) | 25,584 (31.85) | 11,628 (33.15) | 13,802 (29.06) | 24,410 (34.48) |
| Knee, n(%) | 54,745 (68.15) | 23,446 (66.85) | 33,697 (70.94) | 44,494 (65.52) |
| Income | | | | |
| Mean (SD) | 238,435.9 (241,621.3) | 248,234.3 (329,055.5) | 235,810.5 (238,387.4) | 245,333.4 (291,948.1) |
| Self-Efficacy* | | | | |
| | n=32,614 | N=14,283 | n=25,381 | n=21,516 |
| Assigned sex (at birth) | | | | |
| Female, n(%) | 22,720 (69.66) | 9,492 (66.46) | 17,904 (70.54) | 14,308 (66.50) |
| Male, n(%) | 9,894 (30.34) | 4,791 (33.54) | 7,477 (29.46) | 7,208 (33.50) |
| Age | | | | |
| Mean (SD) | 66.59 (9.29) | 66.18 (10.17) | 66.55 (9.12) | 66.37 (10.09) |
| BMI | | | | |
| Mean (SD) | 27.53 (4.85) | 27.78 (5.34) | 27.44 (4.90) | 27.80 (5.12) |
| Worst joint | | | | |
| Hip, n(%) | 10,135 (31.08) | 4,529 (31.71) | 7,250 (28.56) | 7,414 (34.46) |
| Knee, n(%) | 22,479 (68.92) | 9,754 (68.29) | 18,131 (71.44) | 14,102 (65.54) |
| Income | | | | |
| Mean (SD) | 229,400.9 (224,671.7) | 227,699.0 (244,566.4) | 231,166.4 (234,631.4) | 226,187.8 (226,384.9) |
| Desire for Surgery | | | | |
| | n=78,461 | n=36,942 | n=46,751 | n=68,652 |
| Assigned sex (at birth) | | | | |
| Female, n(%) | 53,742 (68.50) | 24,491 (66.30) | 32,784 (70.12) | 45,449 (66.20) |
| Male, n(%) | 24,719 (31.50) | 12,451 (33.70) | 13,968 (29.88) | 23,203 (33.80) |

| Age | | | | |
|--------------------------------|--------------------------|--------------------------|--------------------------|------------------------|
| Mean (SD) | 67.00 (9.36) | 66.03 (10.27) | 66.74 (9.13) | 66.66 (10.02) |
| BMI | | | | |
| Mean(SD) | 27.57 (4.84) | 27.89 (5.21) | 27.43 (4.85) | 27.86 (5.03) |
| Worst joint | | | | |
| Hip, n(%) | 25,041 (31.92) | 12,171 (32.95) | 13,589 (29.07) | 23,623 (32.41) |
| Knee, n(%) | 53,420 (68.08) | 24,771 (67.05) | 33,162 (70.93) | 45,029 (65.59) |
| Income | | | | |
| Mean (SD) | 238,300.4 (240,809.5) | 248,026.5 (326,472.1) | 235,724.3 (239,065.9) | 245,880 (291,036.8) |
| Use of NSAIDs | | | | |
| | N=80,243 | N=35,160 | | |
| Assigned sex (at birth) | | | | |
| Female, n(%) | 54,885 (68.40) | 23,348 (66.41) | | |
| Male, n(%) | 25,358 (31.60) | 11,812 (33.59) | | |
| Age | | | | |
| Mean (SD) | 66.98 (9.38) | 66.03 (10.28) | | |
| BMI | | | | |
| Mean (SD) | 27.59 (4.85) | 27.88 (5.21) | | |
| Worst joint | | | | |
| Hip, n(%) | 25,573 (31.87) | 11,639 (33.10) | | |
| Knee, n(%) | 54,670 (68.13) | 23,521 (66.90) | | |
| Income | | | | |
| (Mean (SD)) | 238,048.3 (242,607.1) | 249,094.9 (327,178.8) | | |

Income-Related Inequalities

Wagstaff et al. (1991) reviewed and assessed different inequality measures suggesting that these measures need: (1) to reflect the socioeconomic dimension of inequalities in health; (2) to mirror the experiences of the whole population; and (3) to be sensitive to modification in the distribution of the population across different socioeconomic classes [166]. They concluded that only the concentration index meets these requirements [166].

The concentration index represents a measure of inequality that highlights how much a health outcome is distributed across a population ranked through a socioeconomic measure (e.g., income) [167]. The standard concentration index is derived from the Gini coefficient of income inequalities. This index requires the investigated health variables

to be on the same scale as the socioeconomic variable (i.e., income – ratio-scaled measure) without an upper bound, and it is calculated as follows [168].

$$C(h|y) = \frac{2cov(h_i, R_i)}{\bar{h}} = \frac{1}{n} \sum_{i=1}^n \left[\frac{h_i}{\bar{h}} (2R_i - 1) \right]$$

Legend: C, concentration index; h_i , health variables where the inequality is measured; R_i , fractional (income) rank.

However, in health economics, health measures are often bounded and either ordinal or cardinal, classified as either attainments (when the observed level of a health variable meets or surpasses the target level) or shortfalls (when the observed level falls below the target level) [169]. Moreover, the concentration index measures relative inequality, invariant to equiproportionate changes in the health variable and reflecting the proportional difference in health variables between subgroups (poor and rich people). Addressing only relative changes provides a limited vision of the extent of inequality. On the other hand, absolute health inequality reflects the absolute magnitude of difference in health variables. Therefore, both relative and absolute health-inequality indices are necessary. To account for these issues, Erreygers proposed his modified concentration index for bounded variables and an absolute and relative indicator of inequality [170].

$$E(h|y) = \frac{1}{n} \sum_{i=1}^n \left[\frac{4h_i}{(h^{max} - h^{min})} (2R_i - 1) \right]$$

Legend: E, Erreygers' concentration index; h_i , the health variable, h^{max} , the upper limit for the health variable, h^{min} , the lower limit of the health variable; R_i , fractional (income) rank.

Erreygers' concentration index (E) values range from -1 to +1. A positive (negative) sign indicates an unequal concentration of the variables of interest among those in higher (lower) SEP. A zero value means no socioeconomic inequality. Larger the magnitude of the concentration index, the greater the extent of income-related inequality [170].

The analysis was performed with Stata 17. We calculated this index using the Stata command 'conindex' [167]. We calculated Erreygers' concentration index for the variables 'Pain intensity', 'Self-efficacy' and 'Use of NSAIDs' when these variables were reported at baseline and 3- and 12-month follow-ups. Specifically, we focussed on on Erreygers' concentration index changes over time to see if any 'intervention generated inequalities' might be present in this Swedish OA first-line intervention. Therefore, we calculated Erreygers' concentration indices and their differences at baseline and 3-month follow-up. We excluded participants with missing values in the SOAR outcomes at either of these time points (listwise deletion). We repeated this process for the baseline and 12-month follow-up, excluding participants with missing values in the SOAR outcomes at either of these time points. To calculate the confidence intervals (CIs) of the indices' differences, we used 1000 bootstrap replications as the

Stata command ‘conindex’ does not provide this parameter. As for ‘Use of NSAIDs’, we only calculated baseline vs 3 months as data after one year were not reported in the registry. However, before analysing the inequalities through Erreygers’ concentration index, we compared those included and excluded in this analysis, and no clinical differences were found among them (Table 10).

Table 10 OA outcomes in those included and excluded in the CI analysis

| | Baseline | 3 months | Baseline | 12 months |
|---|---------------|---------------|---------------|---------------|
| Pain Intensity (NRS 0-10) | | | | |
| Included | | | | |
| | n=80,329 | n=80,329 | n=47,499 | n=47,499 |
| (Mean (SD)) | 5.4 (2.1) | 4.4 (2.3) | 5.2 (2.0) | 4.4 (3.3) |
| Excluded | | | | |
| | n=34,579 | n=259 | n=67,409 | n=149 |
| (Mean (SD)) | 5.7 (2.1) | 4.7 (2.4) | 5.7 (2.1) | 4.5 (2.4) |
| Self-efficacy (ASES Pain and Symptoms, 10-100) | | | | |
| Included | | | | |
| | n=32,614 | n=32,614 | n=25,381 | n=25,381 |
| (Mean (SD)) | 65.5 (16.6) | 69.2 (17.7) | 66.8 (16.2) | 65.1 (18.7) |
| Excluded | | | | |
| | n=12,672 | n=851 | n=19,905 | n=693 |
| (Mean (SD)) | 62.9 (17.8) | 67.0 (18.6) | 62.2 (17.7) | 62.8 (20.2) |
| Desire for Surgery | | | | |
| Included | | | | |
| | n=78,461 | n=78,461 | n=46,751 | n=46,751 |
| No (n (%)) | 58,242 (74.2) | 61,343 (78.2) | 37,504 (80.2) | 36,565 (78.2) |
| Yes (n (%)) | 20,219 (25.8) | 17,118 (21.8) | 9,247 (19.8) | 10,186 (21.8) |
| Excluded | | | | |
| | n=35,252 | n=964 | n=66,962 | n=495 |
| No (n (%)) | 24,085 (68.3) | 684 (71.0) | 44,823 (66.9) | 342 (69.1) |
| Yes (n (%)) | 11,167 (31.7) | 280 (29.1) | 22,139 (33.1) | 153 (30.9) |
| Use of NSAIDs | | | | |

| | Included | |
|-------------|-----------------|---------------|
| | n=80,243 | n=80,243 |
| No (n (%)) | 45,458 (56.7) | 54,749 (68.2) |
| Yes (n (%)) | 34,785 (43.4) | 25,494 (31.8) |
| | Excluded | |
| | n=33,317 | n=185 |
| No (n (%)) | 19,002 (57.0) | 141 (76.2) |
| Yes (n (%)) | 14,315 (43.0) | 44 (23.8) |

Abbreviations: CI, Concentration index; n, number; ASES, arthritis self-efficacy scale; NSAIDs, non-steroidal anti-inflammatory drugs.

Then, Erreygers' concentration indices were calculated for the four investigated OA outcomes in the included cohort (Table 11 and Figure 8). Specifically, the variable 'pain' was more concentrated among people with lower income at the baseline ($E=-0.027$) and became even more concentrated among them after attending the intervention at 3-month follow-up (difference with baseline: $E=-0.011$ [95% CI: -0.015; -0.007]) and 12-month follow-up (difference with baseline: $E=-0.016$ [95% CI: -0.021; -0.012]). Similarly, 'Desire for surgery' was more concentrated among people with lower income at the baseline ($E=-0.008$) and became even more concentrated in this group at 3-month follow-up (difference with baseline: $E=-0.012$ [95% CI: -0.018; -0.005]) and 12-month follow-up (difference with baseline: $E=-0.012$ [95% CI: -0.022; -0.002]). Conversely, the variable 'Self-efficacy' was more concentrated among people with higher income at baseline ($E=0.057$) and became even more concentrated among them after attending the intervention at 3-month follow-up (difference with baseline: $E=0.008$ [95% CI: 0.004; 0.0012]) and 12-month follow-up (difference with baseline: $E=0.017$ [95% CI: 0.012; 0.021]). Finally, the variable 'Use of NSAIDs' was more concentrated among people with lower income at the baseline ($E=0.068$), but this concentration narrowed after attending the intervention at 3-month follow-up (difference with baseline: $E=-0.029$ [95% CI: -0.038; -0.021]).

Table 11 Concentration indices in the OA outcomes

| Variables | N | | Differences | | N | | Differences | |
|---------------------------|--------------------------------|-------------------|-------------------|----------------------------|--------------------------------|--------------------|--------------------|----------------------------|
| | Baseline | 3-month follow-up | 3-month follow-up | Baseline | Baseline | 12-month follow-up | 12-month follow-up | Baseline |
| | Erreycers' Concentration Index | | | | Erreycers' Concentration Index | | | |
| Pain | n=80,329 | -0.027 | -0.039 | -0.011 [-0.015; -0.007] | n=47,499 | -0.027 | -0.043 | -0.016 [-0.021; -0.012] |
| | Erreycers' Concentration Index | | | | Erreycers' Concentration Index | | | |
| Self-Efficacy | n=32,614 | 0.057 | 0.065 | 0.008 [0.004; 0.012] | n=25,370 | 0.058 | 0.075 | 0.017 [0.012; 0.021] |
| | Erreycers' Concentration Index | | | | Erreycers' Concentration Index | | | |
| Desire for Surgery | N=78,461 | -0.008 | -0.020 | -0.012 [-0.018; -0.005] | n=46,751 | -0.012 | -0.024 | -0.012 [-0.022; -0.002] |
| | Erreycers' Concentration Index | | | | Erreycers' Concentration Index | | | |
| Use of NSAIDs | n=80,243 | 0.068 | 0.038 | -0.029 [-0.038; -0.021] | | | | |

Abbreviations: N, number; NSAIDs, non-steroidal anti-inflammatory drugs.




Figure 8 Concentration index values and changes throughout timepoints

Abbreviations: n, number; f, follow-up value; b, baseline value; d: the difference between baseline and reported follow-up; NSAIDs, non-steroidal anti-inflammatory drugs.

Ethical Considerations

Albeit the studies of this thesis could not harm patients, several strict procedures were put in place to conduct this thesis ethically. Regarding data protection, the online survey (Study 1) was delivered through RedCap, a secure web application to build and manage online surveys and databases, respecting the European General Data Protection Regulations. Only the participants who agreed to partake in the study could gain access to it. Moreover, we did not record their IP. Regarding the qualitative study (Study 2), patients were thoroughly instructed and informed about the aim of the research and about how we would treat their data. They had to sign an informed consent and were assigned an identifier (ID). All their data was related to this ID and stored anonymously in a University of Genova OneDrive folder, which was only accessible to those who analysed the interviews. Moreover, all the other documents (i.e., video and audio recordings) were deleted once the interviews were transcribed and anonymised. The register-based studies in Sweden (Studies 3 and 4) generally do not require informed consent. Still, it is assumed that the study participants do not object to registry-based research, provided that such research is deemed ethical by an Ethics Committee. Therefore, we asked for ethics committee approval in Sweden for studies 3 and 4 (Dnr: 2019-02570) and in Italy for studies 1 and 2 (CERA2020.07).

General Discussions

This thesis highlighted some critical areas in the OA care process that have been largely neglected but require further attention. As per the cultural factors, physiotherapists acknowledged the importance of first-line interventions but failed to implement them. Meanwhile, patients have often viewed these interventions merely as a way to pass the time until surgery, with little understanding of their fundamental role. Then, our investigation into the sociodemographic and economic factors affecting adherence to OA treatment did not yield any definitive answers. Therefore, there is a need for further research to identify the underlying factors at play. Finally, although OA self-management programmes have been developed to empower patients, they may inadvertently deepen pre-existing inequalities. Therefore, it is crucial to address these inequalities head-on and ensure that self-management interventions are implemented in a way that does not exacerbate inequalities. Focusing on a more comprehensive and inclusive approach to OA care can improve patients' HRQoL and help them manage their condition effectively.

CPGs are essential to bridge the gap between best evidence and clinical practice. Therefore, clinicians need to be aware of their recommendations to provide high-quality care. In Study 1, over 90% of our sample seemed aware of the importance of exercise and education, consistent with findings from other studies worldwide [87–89, 171]. However, this knowledge was not consistently reflected in their clinical practice. Physiotherapists often excluded weight loss advice, and rest was often considered, although the patient in the clinical vignette had a high BMI and presented moderate symptoms. In fact, some physiotherapists may consider recommending weight loss to be beyond their clinical scope [17]. Also in previous studies, physiotherapists suggested muscle-strengthening exercises, but they refrained from recommending aerobic exercise and weight loss [17, 88]. Providing physiotherapists with specific recommendations for managing people with high BMI can be a strategy to improve patients' outcomes and increase the overall level of adherence to CPGs.

On the other hand, drawing conclusions on the relatively high inclusion of rest and load reduction in the treatment is challenging. A possible explanation is that OA CPGs do not provide specific guidance to tailor exercises to patients' conditions. As a result, physiotherapists may feel unsure and unprepared to deal with exercise in OA [172]. On top of that, different barriers exist to implementing OA CPGs in the complex primary care setting [115]. Just to mention a few, patients' preferences, resource availability, discrepancies between guidelines, lack of English knowledge and access to information

[84, 173]. Furthermore, we found an insufficient level of knowledge of the criteria for the OA clinical diagnosis: only 10% of our sample considered NICE criteria (i.e., being older than 45, having pain and joint stiffness for less than 30 minutes in the morning) sufficient to diagnose OA. The lack of agreement on clinical diagnostic criteria may explain the relatively high percentage (>40%) of physiotherapists considering radiographic findings essential to diagnose OA. This finding about the importance of imaging in OA is coherent with the result of a similar survey by Ayanniyi *et al.* [89]. In OA, radiographic findings should be considered if other diseases are the suspected cause of the symptoms (e.g. infection, cancer, rheumatoid arthritis) or once planning surgical intervention [11]. Likewise, establishing clinical decisions on imaging reinforces the perception of OA as a wear-and-tear disease and may promote fear-avoidance behaviours [116, 174].

Finally, more than 70% of our sample would provide manual therapy in OA. From a cultural perspective, manual therapy is a core physiotherapists' competence that historically set the basis of this healthcare profession. Thus, patients often expect manual therapy [175]. Meeting patients' expectations can improve clinician-patient relationships and enhance the care process [59, 176]. In fact, the quality of the care process depends not only on the appropriateness of treatments delivered by health professionals (technical aspects) but also on different relational and functional aspects investigated in Study 2 [177].

The functional aspects are the basic expectations about how care should be delivered, and they underscore the delivery of effective treatments by trustworthy professionals [74, 177]. Our interviewees did not perceive the treatment indications they received from healthcare professionals to be based on evidence, leading to a sense of uncertainty. Uncertainty is a poorly addressed issue in healthcare that can result in patients' inefficient coping strategies, dysfunctional adaptation to illness and conflicting relationships with healthcare professionals [178–180]. Moreover, our interviewees resorted to looking for information from different sources (e.g., different professionals, the Internet and word-of-mouth) as a result of this perceived uncertainty. Patients' need to interrogate various sources can stem from dissatisfaction with the information received from a healthcare professional [181].

The relational aspect includes all the interactions between the patient and the health professional(s) [177]. The interviewees complained about a lack of empathy in the health professionals they met. Treatments devoid of empathy jeopardise the relationship between patients and health providers, leading to dissatisfied patients that might not adhere to the recommended interventions [182]. In particular, this lack of empathy was specifically perceived during the diagnosis and prognosis of OA. The people who were interviewed sensed a feeling of despair after encountering some healthcare providers. This feeling stemmed from viewing OA as an inevitable consequence of ageing,

treatable only through surgery. In addition, this feeling was reinforced by the structural degeneration shown by their X-rays, which doctors deemed necessary for making a diagnosis and commented on using phrases such as “if I did not know that these x-rays belonged to you, I would think that they belong to someone who is at least 30 years older than you”. However, as reported above, radiographic findings are considered by CPGs as complementary to the assessment of OA [11].

Regarding the technical aspects, patients reported a treatment that was mainly (if not entirely) based on passive therapies while waiting for surgery. Not only was movement considered as dangerous, it was also considered a possible risk factor for anticipated surgery and valuable only after joint replacement. However, people with OA seemed open to undergoing exercise and physiotherapy interventions once educated on their benefits [181, 183]. In our interviews, patients shared a similar experience as they understood the importance of being active and saw exercise as a real support to their care.

According to the findings of this paper, several aspects may function as facilitators or barriers to patients’ exercise adherence, such as the lack of time and unwillingness to change life habits. The former is a well-known cause of lack of adherence to exercise [184–186]. The latter is a shared barrier to implementing diet strategies into patients’ care. Individuals are reluctant to engage in health-promoting behaviours, even though they can reduce mortality and improve their well-being [187].

Study 3 examined exercise adherence in OA. This study was unique as it sought to investigate to what extent demographic, lifestyle, socioeconomic, and disease-related factors were related to adherence to a face-to-face supervised exercise programme for OA in a large group of people with the condition. In our sample, almost 30% had low, 20% had medium, and 50% had high adherence levels. This distribution of adherence levels aligns with that of participants in a similar Danish intervention[188]. Still, it is different from the distribution observed in the online version of the same intervention, which had a higher proportion of people with high levels of adherence [189]. Although multiple factors were linked to adherence, the complete model only accounted for 1% of the variation, indicating that these factors are unlikely to impact exercise adherence substantially.

As far as the demographic and lifestyle factors were concerned, the female sex was negatively associated with a high level of adherence. Women may face societal expectations of caregiving responsibilities, making it harder to find time to exercise[190–194]. However, in the digital version of the investigated Swedish interventions, the female sex was positively associated with high levels of exercise

adherence [189], suggesting that the delivery modality may play a role in the association between sex and adherence. Nevertheless, addressing the root causes of the gender disparities is essential rather than focusing solely on the delivery mode. It is worth noticing that our study only collected information on participants' assigned sex (at birth), limiting generalisability to non-cisgender individuals. Further research is needed to explore gender identity and exercise adherence in people with OA. Then, according to the study's findings, there was a positive correlation between participants' age and high levels of adherence. This result is consistent with previous research that suggested older adults to be more likely to adhere to self-paced exercises, such as those offered in this study, as opposed to moderate-intensity exercise [195]. Finally, BMI was negatively associated with high levels of adherence, as people with high BMI seemed less keen on engaging in physical exercise [196]. In the digital version of the Swedish first-line intervention, BMI was negatively associated with high levels of adherence [189]. Therefore, there is a need to find new strategies to motivate people with high levels of BMI to partake to exercise.

Among the socioeconomic factors, people with lower SEP seemed to adhere more to this intervention. These results contrast the previous literature, where higher SEP seemed to be linked with higher exercise engagement [137, 197]. However, these findings arise primarily from secondary analyses of RCTs not designed to investigate adherence [198]. RCTs per se tend to enhance adherence to treatment which bias the observed associations [199]. Furthermore, in RCTs, people are volunteers selected on specific inclusion and exclusion criteria that may not accurately reflect the socioeconomic diversity of the population from which the sample was drawn [200]. Moreover, individuals with lower SEP may experience a more severe burden of OA than their higher counterparts [27]. Severe symptoms can motivate and drive exercise adherence [197, 201]. Also our study revealed this phenomenon once examining disease-related factors, as frequent pain and difficulty walking were associated with high levels of adherence.

Moreover, we found self-efficacy to be associated with exercise adherence, as in previous evidence [202], but with a modest RRR. Bandura defines self-efficacy as “an individual’s belief in his or her own ability to organise and implement action to produce the desired achievements and results”, such as a behavioural change [203]. Self-efficacy seems characterised by a u-shaped relationship with task accomplishment [204]. People with low self-efficacy tend to doubt their ability to complete a task successfully [204]. In contrast, those with high self-efficacy may exhibit complacency, inadequate preparation, and a focus solely on achieving task-related goals [204]. As a result, low and high levels of self-efficacy may result in a similar outcome, such as low adherence to a task (e.g., exercise). Considering the large number of people in our study, the effect of someone’s self-efficacy might be diluted due to the significant heterogeneity within our population.

However, our model could explain only 1% of the variability, as pointed out by the McFadden R^2 . Thus, it is unlikely to improve adherence significantly by designing strategies based on demographic and lifestyle, socioeconomic and disease-related factors. Therefore, other factors should be taken into account and are further discussed in the 'Future perspective' section of this thesis.

Finally, the socioeconomic factors were further investigated in Study 4 with a concentration index approach. Specifically, we analysed income inequalities in different OA outcomes (i.e., 'Pain intensity', 'Self-efficacy', 'Desire for Surgery' and 'Use of NSAIDs') before and after attending an OA first-line intervention. Our results suggested that pain and desire for surgery were more concentrated among people with lower income, while self-efficacy and use of NSAIDs were more concentrated among people with higher income. Attending the intervention widened income-related inequalities in pain, self-efficacy and desire for surgery, while a narrowing income-related inequality was seen for the use of NSAIDs. In line with that, our data suggest the possible presence of intervention-generated inequalities in the SOAR.

Regarding pain, previous evidence indicated that people in Sweden with lower SEP experienced a higher disease burden, regardless of the socioeconomic index [27, 28]. In Unevik et al., people with lower educational attainment and foreign-born experienced higher pain levels than their higher educated and non-foreign counterparts [27]. In Kiadaliri et al., they found educational attainment and occupation inequalities in the prevalence of knee pain [28]. Hence, our study indicated a similar trend using income as a measure of SEP. Similar scenarios of the association between SEP and higher OA pain were also found worldwide [205–207].

The greater perceived pain intensity can be one potential explanation for the higher concentration of the desire for surgery among individuals with lower income [208, 209]. There is a correlation between total joint replacement (TJR) and health-seeking behaviour, and those with lower SEP require TJR the most due to their more severe symptoms [131]. Our results support those reported in Sweden and other European and non-European countries, showing an association between lower SEP and a stronger desire for surgery [131, 210, 211]. Another reason for the higher willingness for surgery in those with lower income can be the lower reported self-efficacy. In first-line interventions for OA, individuals with this condition play an active role in their care process by collaborating with healthcare providers, sharing expertise, setting goals, and taking responsibility for their treatment by selecting and adopting self-management strategies that impact their lifestyle. Therefore, individuals with higher income and higher levels of self-efficacy may be more self-assured in their ability to manage the

burden of their disease by adopting active treatments (such as those proposed in OA first-line intervention programmes) rather than opting for surgery.

The greater reported benefits experienced by individuals with higher income may also explain why the pro-rich inequality in the use of NSAIDs was reduced after attending the intervention. Prior to the intervention, individuals with higher income had a higher prevalence of NSAID use. This tendency could be explained by the fact that people with higher income may have greater access to healthcare services, resulting in more opportunities to receive prescriptions for NSAIDs. However, following the intervention led to a reduction in these pro-rich inequalities, underscoring the fact that people with higher income derived the greatest benefit from the intervention. Existing research on the link between income and NSAID use has yielded inconsistent findings. Bonnesen et al. found that the lowest-income people had a lower probability of getting additional NSAID prescriptions [212]. Nielsen et al. found that having a low income was slightly associated with the prescription of NSAID but not with their over-the-counter use [213]. Finally, Fosbøl et al. found no differences in NSAID use among those in the highest or lowest income categories [214]. Previous studies on income and health outcomes have often categorised or dichotomised income instead of treating it as a continuous variable. However, such an approach may result in statistical analysis limitations, including reduced statistical power due to the loss of information [215]. In addition, the other studies used average association measures such as OR, which may not effectively capture differences in the distribution of NSAID use across the entire population, as they rely only on average associations between different income groups [216].

Studies' Limitations

In Study 1, our sample was mainly composed of physiotherapists with a post-graduate degree (47%). Therefore, we might have overestimated the real level of knowledge of and adherence to osteoarthritis CPGs. In Study 2, we gathered a small sample size that limits the generalisability of the results [217]. However, qualitative studies do not aim at data generalisation [110]. Besides, all the interviewees lived in a similar geographical area (i.e. northern Italy) and were mostly women and Caucasian. This is particularly important since meanings attached to an experience might be influenced by gender, ethnicity and geographical location. Finally, the interviewees were at different stages of their care process. However, all patients agreed with what emerged during the member checking phase that was held at the end of the study by sending our results to all the interviewees. Additionally, some variables had missing values, but the missingness of our data is likely to be completely at random and not introduce significant bias in our results. Furthermore, our study lacks data on other predictors of adherence, and certain reported variables (e.g., weekly physical activity) may have limited explanatory power due to the way they were measured. Our results may not generalise to other forms of exercise (e.g., unsupervised home exercise) due to the specific focus of our study. Finally, while certain variables were found to have medium but not high levels of association with adherence (such as 'physical activity hours', 'number of painful joints', and 'living alone'), this finding may be influenced by chance or the adherence categorisation used in the SOAR study. In Study 4, those registered in the SOAR have higher SEP than the general Swedish population [150]. Therefore, our results might underestimate the impact of this OA first-line intervention in generating inequalities. Then, this intervention does not have a comparable control group. Thus, we do not know whether the observed results can be attributed to the intervention itself. Moreover, a few variables were missing as per Study 3. To conclude, the observational nature of our studies, did not allow us to identify any causal relationship. Overall, this thesis included studies from two countries that are very different to one another. Nevertheless, we compared our results to similar studies conducted worldwide, rather than focussing on Italy and Sweden solely.

Future Perspectives

This thesis further proves the need to improve OA care at various levels. Healthcare professionals could be targeted to bridge the gap between evidence and clinical practice. Therefore, future studies should investigate new strategies for accomplishing this and are discussed hereafter. CPGs should include essential practical elements, such as algorithms for clinical decision-making in complicated cases and inclusion-exclusion criteria for patients, once compiling CPGs. Providing physiotherapists with CPGs in their native language and promoting their use in university programmes may be another possible solution to improve compliance with recommendations. There is also a need to reconceptualise the professional image of physiotherapists within society. Specifically, we need to shift towards a new vision of physiotherapists no longer associated solely with physical and manual therapy but focused on improving patients' functioning through active treatment strategies such as exercise and education, tailored to specific contexts and grounded in scientific evidence.

As per the patients, this thesis sheds some light on common themes in the experience of people with OA. People with hip and knee OA often encounter an uncertain care process, characterised by the lack of clear explanations, insufficient empathy, and negative attitudes towards first-line non-surgical treatments. These factors underscore the importance of implementing effective communication and providing patients with adequate information about treatment options. As a result, patients' beliefs can be positively impacted, increasing their awareness and compliance with first-line interventions.

Moreover, we need to understand which factors are associated with exercise adherence to create tailored interventions to improve it. Modern and well-designed RCTs are exposing the weaknesses in the evidence supporting the effectiveness of exercise in curbing OA pain [218–220]. However, exercise leads to benefit that goes beyond joint health as exercise positively affects body weight, lipid metabolism, hyperglycaemia, mood and systemic inflammation, preventing different OA-related comorbidities (e.g., diabetes and depression) [33, 34]. In this thesis, we indicated that strategies based on demographic, lifestyle, socioeconomic and disease-related factors are unlikely to improve adherence significantly. On the other hand, preliminary evidence suggested that improving mindsets about exercise can increase adherence [221]. Mindsets are “core assumptions about a domain or category that orient individuals to a particular set of attributions, expectations, and goals” [221, 222]. Mindsets are responsible for encoding contextual factors (e.g., the facility's structure, the clinician's communication style, and

the ability to motivate patients), affecting people's outcomes via a placebo (or nocebo) response if positively (placebo) or negatively (nocebo) judged by one's brain [109]. Booster sessions, reminders, and behavioural change techniques can also improve exercise adherence by increasing motivation to engage in exercise [223, 224]. These strategies seem to ground their efficacy on contextual factors too (e.g., communication with the clinicians, feeling to be taken care of by them etc.). Future studies should explore the relationship between contextual factors, mindsets, and exercise adherence to understand better how to improve adherence in patients with OA.

Finally, we need to reduce inequalities in OA care. Bearing in mind that there is no society without inequalities, it is essential to find new strategies to reduce them in OA care. Our results suggested the possible presence of intervention-generated inequalities in this OA first-line intervention. Moreover, past evidence showed that people in socially disadvantaged backgrounds might not even begin this intervention, though delivered in the public healthcare system [150]. Therefore, we need to evaluate the systemic, institutional, and power-related harms of interventions by adopting a health-equity approach to make these interventions more accessible to underserved populations. Past evidence showed that upstream interventions are necessary to tackle the root causes of health disparities [133, 225, 226]. These interventions involve political, social, and economic policies (e.g., fiscal food policies, urban planning policies, transport policies, increase wage etc.) [133, 227]. Conversely, downstream interventions like healthcare and individual behaviour change play a limited (to no) role in reducing inequalities and should not be the main focus of decision-making related to health inequalities [133].

Conclusions

Several areas in the OA care process require attention and are currently being neglected. To improve the quality of OA care, we need to take further actions. Firstly, we need to close the evidence-to-practice gap among healthcare professionals working with people with OA. Secondly, we need to raise awareness among people with OA of their active role in their care and the importance of first-line interventions. Thirdly, we need to acknowledge that adherence to exercise is a complex behaviour that cannot be explained solely by people's lifestyle, demographic, socioeconomic and disease-related status. Therefore, we need to invest more resources in understanding which factors are related to exercise adherence and develop tailored interventions to target and improve it. Finally, while self-management programmes have been designed to empower patients, they may inadvertently exacerbate existing inequalities. Therefore, we need to find new ways to reduce inequalities in OA care, prioritising upstream interventions.

References

1. Martel-Pelletier J, Barr AJ, Cicuttini FM, Conaghan PG, Cooper C, Goldring MB, et al. Osteoarthritis. *Nat Rev Dis Primers*. 2016;2.
2. Brandt KD, Radin EL, Dieppe PA, van de Putte L. Yet more evidence that osteoarthritis is not a cartilage disease. *Ann Rheum Dis*. 2006;65:1261–4.
3. Hunter DJ, Bierma-Zeinstra S. Osteoarthritis. *The Lancet*. 2019;393:1745–59.
4. Long H, Liu Q, Yin H, Wang K, Diao N, Zhang Y, et al. Prevalence Trends of Site-Specific Osteoarthritis From 1990 to 2019: Findings From the Global Burden of Disease Study 2019. *Arthritis and Rheumatology*. 2022;74:1172–83.
5. Long H, Liu Q, Yin H, Wang K, Diao N, Zhang Y, et al. Prevalence Trends of Site-Specific Osteoarthritis From 1990 to 2019: Findings From the Global Burden of Disease Study 2019. *Arthritis and Rheumatology*. 2022;74:1172–83.
6. Kolasinski SL, Neogi T, Hochberg MC, Oatis C, Guyatt G, Block J, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Management of Osteoarthritis of the Hand, Hip, and Knee. *Arthritis and Rheumatology*. 2020;72:220–33.
7. Vos T, Abajobir AA, Abbafati C, Abbas KM, Abate KH, Abd-Allah F, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: A systematic analysis for the Global Burden of Disease Study 2016. *The Lancet*. 2017;390:1211–59.
8. Hunter DJ, Schofield D, Callander E. The individual and socioeconomic impact of osteoarthritis. *Nat Rev Rheumatol*. 2014;10:437–41.
9. Weng Q, Goh S-L, Wu J, Persson MSM, Wei J, Sarmanova A, et al. Comparative efficacy of exercise therapy and oral non-steroidal anti-inflammatory drugs and paracetamol for knee or hip osteoarthritis: a network meta-analysis of randomised controlled trials. *Br J Sports Med*. 2023;0:bjssports-2022-105898.
10. Bannuru RR, Osani MC, Vaysbrot EE, Arden NK, Bennell K, Bierma-Zeinstra SMA, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis Cartilage*. 2019;27:1578–89.
11. National Clinical Guidelines Centre. Osteoarthritis: Care and Management in Adults. NICE Clinical Guidelines. 2014;:1–498.

12. Basedow M, Esterman A. Assessing appropriateness of osteoarthritis care using quality indicators: a systematic review. *J Eval Clin Pract.* 2015;21:782–9.
13. Theis KA, Brady TJ, Sacks JJ. Where Have All the Patients Gone? Profile of US Adults Who Report Doctor-Diagnosed Arthritis But Are Not Being Treated. *JCR: Journal of Clinical Rheumatology.* 2018;25:341–7.
14. Runciman WB, Hunt TD, Hannaford NA, Hibbert PD, Westbrook JI, Coiera EW, et al. CareTrack: Assessing the appropriateness of health care delivery in Australia. *Medical Journal of Australia.* 2012;197:100–5.
15. Battista S, Salvioli S, Millotti S, Testa M, Dell’Isola A. Italian physiotherapists’ knowledge of and adherence to osteoarthritis clinical practice guidelines: a cross-sectional study. *BMC Musculoskelet Disord.* 2021;22:380.
16. Holden MA, Nicholls EE, Young J, Hay EM, Foster NE. UK-based physical therapists’ attitudes and beliefs regarding exercise and knee osteoarthritis: Findings from a mixed-methods study. 2009;61:1511–21.
17. Holden MA, Waterfield J, Whittle R, Bennell K, Quicke JG, Chesterton L, et al. How do UK physiotherapists address weight loss among individuals with hip osteoarthritis? A mixed-methods study. *Musculoskeletal Care.* 2019;17:133–44.
18. Battista S, Dell’Isola A, Manoni M, Englund M, Palese A, Testa M. Experience of the COVID-19 pandemic as lived by patients with hip and knee osteoarthritis: an Italian qualitative study. *BMJ Open.* 2021;11:e053194.
19. Battista S, Manoni M, Dell’Isola A, Englund M, Palese A, Testa M. Giving an account of patients’ experience: A qualitative study on the care process of hip and knee osteoarthritis. *Health Expectations.* 2022. <https://doi.org/10.1111/hex.13468>.
20. Marks R. Knee osteoarthritis and exercise adherence: a review. *Curr Aging Sci.* 2012;5:72–83.
21. Taulaniemi A, Kankaanpää M, Rinne M, Tokola K, Parkkari J, Suni JH. Fear-avoidance beliefs are associated with exercise adherence: secondary analysis of a randomised controlled trial (RCT) among female healthcare workers with recurrent low back pain. *BMC Sports Sci Med Rehabil.* 2020;12.
22. Bennell KL, Dobson F, Hinman RS. Exercise in osteoarthritis: moving from prescription to adherence. *Best Pract Res Clin Rheumatol.* 2014;28:93–117.
23. Loew L, Brosseau L, Kenny GP, Durand-Bush N, Poitras S, de Angelis G, et al. Factors influencing adherence among older people with osteoarthritis. *Clin Rheumatol.* 2015;35:2283–91.
24. Duong V, Hunter DJ, Nicolson PJA. Predictors and Measures of Adherence to Core Treatments for Osteoarthritis. *Clin Geriatr Med.* 2022;38:345–60.

25. Lorenc T, Petticrew M, Welch V, Tugwell P. What types of interventions generate inequalities? Evidence from systematic reviews. *J Epidemiol Community Health*. 2013;67:190–3.
26. Peat G, Jordan KP, Wilkie R, Corp N, van der Windt DA, Yu D, et al. Do recommended interventions widen or narrow inequalities in musculoskeletal health? An equity-focussed systematic review of differential effectiveness. *J Public Health (Bangkok)*. 2022;44:e376–87.
27. Unevik E, Abbott A, Fors S, Rolfson O. Sociodemographic inequalities in outcomes of a swedish nationwide self-management program for osteoarthritis: Results from 22,741 patients between years 2008–2017. *J Clin Med*. 2020;9:1–14.
28. Kiadaliri AA, Gerhardsson de Verdier M, Turkiewicz A, Lohmander LS, Englund M. Socioeconomic inequalities in knee pain, knee osteoarthritis, and health-related quality of life: a population-based cohort study in southern Sweden. *Scand J Rheumatol*. 2017;46:143–51.
29. Fernandes L, Hagen KB, Bijlsma JWJ, Andreassen O, Christensen P, Conaghan PG, et al. EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. *Annals of the Rheumatic Diseases*. 2013;72:1125–35.
30. National Clinical Guidelines Centre. Osteoarthritis in over 16s: diagnosis and management. 2022.
31. Zeng CY, Zhang ZR, Tang ZM, Hua FZ. Benefits and Mechanisms of Exercise Training for Knee Osteoarthritis. *Front Physiol*. 2021;12:2267.
32. Raposo F, Ramos M, Lúcia Cruz A. Effects of exercise on knee osteoarthritis: A systematic review. *Musculoskeletal Care*. 2021;19:399–435.
33. Wesseling J, Welsing PMJ, Bierma-zeinstra SMA, Dekker J, Gorter KJ, Kloppenburg M, et al. Impact of self-reported comorbidity on physical and mental health status in early symptomatic osteoarthritis: the CHECK (Cohort Hip and Cohort Knee) study. *Rheumatology (Oxford)*. 2013;52:180–8.
34. Swain S, Sarmanova A, Coupland C, Doherty M, Zhang W. Comorbidities in Osteoarthritis: A Systematic Review and Meta-Analysis of Observational Studies. *Arthritis Care Res (Hoboken)*. 2020;72:991–1000.
35. Messier SP, Mihalko SL, Legault C, Miller GD, Nicklas BJ, DeVita P, et al. Effects of intensive diet and exercise on knee joint loads, inflammation, and clinical outcomes among overweight and obese adults with knee osteoarthritis: The IDEA randomized clinical trial. *JAMA - Journal of the American Medical Association*. 2013;310:1263–73.
36. Messier SP, Newman JJ, Scarlett MJ, Mihalko SL, Miller GD, Nicklas BJ, et al. Changes in Body Weight and Knee Pain in Adults With Knee Osteoarthritis Three-and-a-Half Years After Completing Diet and Exercise Interventions: Follow-Up Study for a

- Single-Blind, Single-Center, Randomized Controlled Trial. *Arthritis Care Res (Hoboken)*. 2022;74:607–16.
37. Dunlop DD, Song J, Semanik PA, Chang RW, Sharma L, Bathon JM, et al. Objective physical activity measurement in the Osteoarthritis Initiative: Are guidelines being met? *Arthritis Rheum*. 2011;63:3372.
38. de Groot IB, Bussmann JB, Stam HJ, Verhaar JAN. Actual everyday physical activity in patients with end-stage hip or knee osteoarthritis compared with healthy controls. *Osteoarthritis Cartilage*. 2008;16:436–42.
39. Dunlop DD, Song J, Semanik PA, Sharma L, Chang RW. Physical activity levels and functional performance in the Osteoarthritis Initiative: a graded relationship. *Arthritis Rheum*. 2011;63:127.
40. Hauner H. Secretory factors from human adipose tissue and their functional role. *Proc Nutr Soc*. 2005;64:163–9.
41. Thomas S, Browne H, Mobasher A, Rayman MP. What is the evidence for a role for diet and nutrition in osteoarthritis? *Rheumatology (Oxford)*. 2018;57 suppl_4:iv61–74.
42. Messier SP, Mihalko SL, Legault C, Miller GD, Nicklas BJ, DeVita P, et al. Effects of intensive diet and exercise on knee joint loads, inflammation, and clinical outcomes among overweight and obese adults with knee osteoarthritis: the IDEA randomized clinical trial. *JAMA*. 2013;310:1263–73.
43. Wluka AE, Lombard CB, Cicuttini FM. Tackling obesity in knee osteoarthritis. *Nat Rev Rheumatol*. 2013;9:225–35.
44. Fibel KH. State-of-the-Art management of knee osteoarthritis. *World J Clin Cases*. 2015;3:89.
45. Hunter DJ, McDougall JJ, Keefe FJ. The Symptoms of Osteoarthritis and the Genesis of Pain. *Medical Clinics of North America*. 2009;93:83–100.
46. Fu K, Robbins SR, McDougall JJ. Osteoarthritis: the genesis of pain. *Rheumatology*. 2018;57 suppl_4:iv43–50.
47. Hawker GA, Stewart L, French MR, Cibere J, Jordan JM, March L, et al. Understanding the pain experience in hip and knee osteoarthritis – an OARSI/OMERACT initiative. *Osteoarthritis Cartilage*. 2008;16:415–22.
48. Neogi T. The epidemiology and impact of pain in osteoarthritis. *Osteoarthritis Cartilage*. 2013;21:1145–53.
49. Neogi T, Nevitt MC, Yang M, Curtis JR, Torner J, Felson DT. Consistency of knee pain: correlates and association with function. *Osteoarthritis Cartilage*. 2010;18:1250–5.

50. Kittelson AJ, Schmiege SJ, Maluf K, George SZ, Stevens-Lapsley JE. Determination of Pain Phenotypes in Knee Osteoarthritis Using Latent Profile Analysis. *Pain Med.* 2021;22:653–62.
51. Felson DT, Chaisson CE, Hill CL, Totterman SMS, Gale ME, Skinner KM, et al. The association of bone marrow lesions with pain in knee osteoarthritis. *Ann Intern Med.* 2001;134:541–9.
52. Baker K, Grainger A, Niu J, Clancy M, Guermazi A, Crema M, et al. Relation of synovitis to knee pain using contrast-enhanced MRIs. *Ann Rheum Dis.* 2010;69:1779–83.
53. Sofat N, Ejindu V, Kiely P. What makes osteoarthritis painful? The evidence for local and central pain processing. *Rheumatology (Oxford).* 2011;50:2157–65.
54. Schaible HG, Richter F, Ebersberger A, Boettger MK, Vanegas H, Natura G, et al. Joint pain. *Exp Brain Res.* 2009;196:153–62.
55. Schaible HG, Ebersberger A, Segond Von Banchet G. Mechanisms of pain in arthritis. *Ann N Y Acad Sci.* 2002;966:343–54.
56. Lluch E, Torres R, Nijs J, van Oosterwijck J. Evidence for central sensitization in patients with osteoarthritis pain: a systematic literature review. *Eur J Pain.* 2014;18:1367–75.
57. Richter F, Natura G, Löser S, Schmidt K, Viisanen H, Schaible HG. Tumor necrosis factor causes persistent sensitization of joint nociceptors to mechanical stimuli in rats. *Arthritis Rheum.* 2010;62:3806–14.
58. Lentz TA, George SZ, Manickas-Hill O, Malay MR, O'Donnell J, Jayakumar P, et al. What General and Pain-associated Psychological Distress Phenotypes Exist Among Patients with Hip and Knee Osteoarthritis? *Clin Orthop Relat Res.* 2020;478:2768–83.
59. Rossetini G, Carlino E, Testa M. Clinical relevance of contextual factors as triggers of placebo and nocebo effects in musculoskeletal pain. *BMC Musculoskelet Disord.* 2018;19:27.
60. Hunter DJ, Guermazi A, Roemer F, Zhang Y, Neogi T. Structural correlates of pain in joints with osteoarthritis. *Osteoarthritis Cartilage.* 2013;21:1170–8.
61. Swain S, Sarmanova A, Coupland C, Doherty M, Zhang W. Comorbidities in Osteoarthritis: A Systematic Review and Meta-Analysis of Observational Studies. *Arthritis Care Res (Hoboken).* 2020;72:991–1000.
62. Kadam UT, Jordan K, Croft PR. Clinical comorbidity in patients with osteoarthritis: a case-control study of general practice consultants in England and Wales. *Ann Rheum Dis.* 2004;63:408–14.
63. Wolfe F, Hawley DJ, Peloso PM, Wilson K, Anderson J. Back pain in osteoarthritis of the knee. *Arthritis Rheum.* 1996;9:376–83.

64. Sale JEM, Gignac M, Hawker G. The relationship between disease symptoms, life events, coping and treatment, and depression among older adults with osteoarthritis. *Journal of Rheumatology*. 2008;35:335–42.
65. Wang ST, Ni GX. Depression in Osteoarthritis: Current Understanding. *Neuropsychiatr Dis Treat*. 2022;18:375–89.
66. Gignac MAM, Cott C, Badley EM. Adaptation to Chronic Illness and Disability and Its Relationship to Perceptions of Independence and Dependence. *J Gerontol B Psychol Sci Soc Sci*. 2000;55:362–72.
67. Hawker GA. Osteoarthritis is a serious disease. *Clin Exp Rheumatol*. 2019;37 Suppl 1:3–6.
68. Hall AJ, Stubbs B, Mamas MA, Myint PK, Smith TO. Association between osteoarthritis and cardiovascular disease: Systematic review and meta-analysis. *Eur J Prev Cardiol*. 2016;23:938–46.
69. Puenpatom RA, Victor TW. Increased prevalence of metabolic syndrome in individuals with osteoarthritis: An analysis of NHANES III data. *Postgrad Med*. 2009;121:9–20.
70. Eyles JP, Sharma S, Telles RW, Namane M, Hunter DJ, Bowden JL. Implementation of Best-Evidence Osteoarthritis Care: Perspectives on Challenges for, and Opportunities From, Low and Middle-Income Countries. *Frontiers in Rehabilitation Sciences*. 2022;2:126.
71. Zadro J, O’Keeffe M, Maher C. Do physical therapists follow evidence-based guidelines when managing musculoskeletal conditions? Systematic review. *BMJ Open*. 2019;9.
72. Chou L, Ellis L, Papandony M, Maheeka Seneviwickrama KLD, Cicuttini FM, Sullivan K, et al. Patients’ perceived needs of osteoarthritis health information: A systematic scoping review. *PLoS One*. 2018;13.
73. Horne R. Patients’ beliefs about treatment: the hidden determinant of treatment outcome? *J Psychosom Res*. 1999;47:491–5.
74. Chi-Lun-Chiao A, Chehata M, Broeker K, Gates B, Ledbetter L, Cook C, et al. Patients’ perceptions with musculoskeletal disorders regarding their experience with healthcare providers and health services: an overview of reviews. *Arch Physiother*. 2020;10.
75. Laferton JAC, Kube T, Salzmann S, Auer CJ, Shedden-Mora MC. Patients’ expectations regarding medical treatment: A critical review of concepts and their assessment. *Frontiers in Psychology*. 2017;8 FEB:233.
76. Institute of Medicine (IOM). *Clinical Practice Guidelines We Can Trust*. Washington DC: National Academies Press; 2011.

77. Juhl C, Christensen R, Roos EM, Zhang W, Lund H. Impact of exercise type and dose on pain and disability in knee osteoarthritis: A systematic review and meta-regression analysis of randomized controlled trials. *Arthritis and Rheumatology*. 2014;66:622–36.
78. Thorstensson CA, Garellick G, Rystedt H, Dahlberg LE. Better Management of Patients with Osteoarthritis: Development and Nationwide Implementation of an Evidence-Based Supported Osteoarthritis Self-Management Programme. *Musculoskeletal Care*. 2015;13:67–75.
79. McGlynn EA, Asch SM, Adams J, Keesey J, Hicks J, DeCristofaro A, et al. The quality of health care delivered to adults in the United States. *New England Journal of Medicine*. 2003;348:2635–45.
80. Egerton T, Diamond LE, Buchbinder R, Bennell KL, Slade SC. A systematic review and evidence synthesis of qualitative studies to identify primary care clinicians' barriers and enablers to the management of osteoarthritis. *Osteoarthritis and Cartilage*. 2017;25:625–38.
81. Dziedzic KS, French S, Davis AM, Geelhoed E, Porcheret M. Implementation of musculoskeletal Models of Care in primary care settings: Theory, practice, evaluation and outcomes for musculoskeletal health in high-income economies. *Best Practice and Research: Clinical Rheumatology*. 2016;30:375–97.
82. Fischer F, Lange K, Klose K, Greiner W, Kraemer A. Barriers and Strategies in Guideline Implementation—A Scoping Review. *Healthcare*. 2016;4:36.
83. Gupta S, Rai N, Bhattacharyya O, Cheng AYY, Connelly KA, Boulet LP, et al. Optimizing the language and format of guidelines to improve guideline uptake. *CMAJ*. 2016;188:E362–8.
84. Cutolo R, Battista S, Testa M. Actual and perceived level of scientific english across italian physiotherapy courses: A cross-sectional study. *Healthcare (Switzerland)*. 2021;9.
85. Holden MA, Nicholls EE, Hay EM, Foster NE. Physical Therapists' Use of Therapeutic Exercise for Patients With Clinical Knee Osteoarthritis in the United Kingdom: In Line With Current Recommendations? *Phys Ther*. 2008;88:1109–21.
86. Holden MA, Nicholls EE, Young J, Hay EM, Foster NE. UK-based physical therapists' attitudes and beliefs regarding exercise and knee osteoarthritis: Findings from a mixed-methods study. *Arthritis Rheum*. 2009;61:1511–21.
87. Tittlemier BJ, Wittmeier KD, Robinson DB, Webber SC. Knee Osteoarthritis: An Investigation into the Clinical Practice of Physiotherapists in Canada. *Physiotherapy Canada*. 2020;:e20190068.

88. Tang CY, Pile R, Croft A, Watson NJ. Exploring Physical Therapist Adherence to Clinical Guidelines When Treating Patients With Knee Osteoarthritis in Australia: A Mixed Methods Study. *Phys Ther.* 2020;100:1084–93.
89. Ayanniyi O, Egwu RF, Adeniyi AF. Physiotherapy management of knee osteoarthritis in Nigeria—A survey of self-reported treatment preferences. *Hong Kong Physiotherapy Journal.* 2017;36:1–9.
90. Holden MA, Nicholls EE, Hay EM, Foster NE. Physical Therapists' Use of Therapeutic Exercise for Patients With Clinical Knee Osteoarthritis in the United Kingdom: In Line With Current Recommendations? *Phys Ther.* 2008;88:1109–21.
91. Kristensen N, Nymann C, Konradsen H. Implementing research results in clinical practice- the experiences of healthcare professionals. *BMC Health Serv Res.* 2015;16:48.
92. Mazzola P, Rimoldi SML, Rossi P, Noale M, Rea F, Facchini C, et al. Aging in Italy: The Need for New Welfare Strategies in an Old Country. *Gerontologist.* 2016;56:383–90.
93. Romanini E, Decarolis F, Luzi I, Zanoli G, Venosa M, Laricchiuta P, et al. Total knee arthroplasty in Italy: reflections from the last fifteen years and projections for the next thirty. *Int Orthop.* 2019;43:133–8.
94. Vliet Vlieland TPM, van den Ende CHM, Alliot-Launois F, Beauvais C, Gobbo M, Iagnocco A, et al. Educational needs of health professionals working in rheumatology in Europe. *RMD Open.* 2016;2:e000337.
95. Eysenbach G, Wyatt J. Using the Internet for surveys and health research. *J Med Internet Res.* 2002;4:76–94.
96. de Leeuw D, Hox J, Dillman D. *International Handbook of Survey Methodology.* 1st edition. New York: Taylor and Francis group; 2008.
97. Eysenbach G. Improving the Quality of Web Surveys: The Checklist for Reporting Results of Internet E-Surveys (CHERRIES) | Eysenbach | *Journal of Medical Internet Research.* *J Med Internet Res.* 2004;6:e34+.
98. von Elm E, Altman D, Egger M, Pocock S, Gøtzsche P, Vandenbroucke J, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Ann Intern Med.* 2007;147:573–8.
99. The European Parliament and the Council of the European Union. Regulation (EU) 2016/679 of the European Parliament and of the Council of 27 April 2016 on the protection of natural persons with regard to the processing of personal data and on the free movement of such data, and repealing Directive 95/46/EC. 2016.

100. Suárez-Alvarez J, Pedrosa I, Lozano LM, García-Cueto E, Cuesta M, Muñiz J. Using reversed items in likert scales: A questionable practice. *Psicothema*. 2018;30:149–58.
101. Caffini G, Battista S, Raschi A, Testa M. Physiotherapists' knowledge of and adherence to evidence-based practice guidelines and recommendations for ankle sprains management: a cross-sectional study. *BMC Musculoskelet Disord*. 2022;23:1–12.
102. Teo PL, Hinman RS, Egerton T, Dziedzic KS, Bennell KL. Identifying and Prioritizing Clinical Guideline Recommendations Most Relevant to Physical Therapy Practice for Hip and/or Knee Osteoarthritis. *Journal of Orthopaedic & Sports Physical Therapy*. 2019;49:501–12.
103. Rutten GMJ, Harting J, Rutten STJ, Bekkering GE, Kremers SPJ. Measuring physiotherapists' guideline adherence by means of clinical vignettes: A validation study. *J Eval Clin Pract*. 2006;12:491–500.
104. Katz JN, Arant KR, Loeser RF. Diagnosis and Treatment of Hip and Knee Osteoarthritis: A Review. *JAMA*. 2021;325:568–78.
105. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: An emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924–6.
106. Sebrant L, Jong M. What's the meaning of the concept of caring?: a meta-synthesis. *Scandinavian Journal of Caring Sciences*. 2020;35:353–65.
107. Mortari L, Saiani L. Gestures and thoughts of caring: A Theory Of Caring From The Voices Of Nurses. 1st edition. 2014.
108. Benedetti F. Placebo-Induced Improvements: How Therapeutic Rituals Affect the Patient's Brain. *JAMS Journal of Acupuncture and Meridian Studies*. 2012;5:97–103.
109. Testa M, Rossettini G. Enhance placebo, avoid nocebo: How contextual factors affect physiotherapy outcomes. *Man Ther*. 2016;24:65–74.
110. Moser A, Korstjens I. Series: Practical guidance to qualitative research. part 1: Introduction. *European Journal of General Practice*. 2017;23:271–3.
111. Moustakas C. Phenomenological research methods. CA: SAGE Publications, Inc.; 1994.
112. Spiegelberg H. Doing Phenomenology: Essays on and in Phenomenology. Springer; 1975.
113. Matua GA, van der Wal DM. Differentiating between descriptive and interpretive phenomenological research approaches. *Nurse Res*. 2015;22:22–7.
114. Korstjens I, Moser A. Series: Practical guidance to qualitative research. part 2: Context, research questions and designs. *European Journal of General Practice*. 2017;23:274–9.

115. Swaithe L, Paskins Z, Dziedzic K, Finney A. Factors influencing the implementation of evidence-based guidelines for osteoarthritis in primary care: A systematic review and thematic synthesis. *Musculoskeletal Care*. 2020;Ahead of P:1–10.
116. Darlow B, Brown M, Thompson B, Hudson B, Grainger R, McKinlay E, et al. Living with osteoarthritis is a balancing act: An exploration of patients' beliefs about knee pain. *BMC Rheumatol*. 2018;2:15.
117. Hurley M, Dickson K, Hallett R, Grant R, Hauari H, Walsh N, et al. Exercise interventions and patient beliefs for people with hip, knee or hip and knee osteoarthritis: A mixed methods review. *Cochrane Database of Systematic Reviews*. 2018;2018.
118. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International Journal for Quality in Health Care*. 2007;19:349–57.
119. Korstjens I, Moser A. Series: Practical guidance to qualitative research. Part 4: Trustworthiness and publishing. *European Journal of General Practice*. 2018;24:120–4.
120. Birt L, Scott S, Cavers D, Campbell C, Walter F. Member Checking: A Tool to Enhance Trustworthiness or Merely a Nod to Validation? *Qual Health Res*. 2016;26:1802–11.
121. Moser A, Korstjens I. Series: Practical guidance to qualitative research. Part 3: Sampling, data collection and analysis. *European Journal of General Practice*. 2018;24:9–18.
122. Saunders B, Sim J, Kingstone T, Baker S, Waterfield J, Bartlam B, et al. Saturation in qualitative research: exploring its conceptualization and operationalization. *Qual Quant*. 2018;52:1893–907.
123. Jefferis BJ, Sartini C, Lee IM, Choi M, Amuzu A, Gutierrez C, et al. Adherence to physical activity guidelines in older adults, using objectively measured physical activity in a population-based study. *BMC Public Health*. 2014;14:1–9.
124. Hemmingsson E, Ekelund U. Is the association between physical activity and body mass index obesity dependent? *Int J Obes (Lond)*. 2007;31:663–8.
125. Marmot M. Social determinants of health inequalities. *The Lancet*. 2005;365:1099–104.
126. Austin S, Qu H, Shewchuk RM. Association between adherence to physical activity guidelines and health-related quality of life among individuals with physician-diagnosed arthritis. *Quality of Life Research*. 2012;21:1347–57.
127. Pisters MF, Veenhof C, Schellevis FG, Twisk JWR, Dekker J, de Bakker DH. Exercise adherence improving long-term patient outcome in patients with osteoarthritis of the hip and/or knee. *Arthritis Care Res (Hoboken)*. 2010;62:1087–94.

128. Safiri S, Kolahi AA, Smith E, Hill C, Bettampadi D, Mansournia MA, et al. Global, regional and national burden of osteoarthritis 1990-2017: a systematic analysis of the Global Burden of Disease Study 2017. *Ann Rheum Dis*. 2020;79.
129. Battista S, Kiadaliri A, Jönsson T, Gustafsson K, Englund M, Testa M, et al. Factors Associated with Adherence To a Supervised Exercise Intervention for Osteoarthritis: Data From the Swedish Osteoarthritis Registry. *Arthritis Care Res (Hoboken)*. 2023. <https://doi.org/10.1002/ACR.25135>.
130. Olsson CB, Ekelund J, Degerstedt Å, Thorstensson CA. Change in self-efficacy after participation in a supported self-management program for osteoarthritis - an observational study of 11 906 patients. *Disabil Rehabil*. 2020;42:2133–40.
131. Brennan SL, Lane SE, Lorimer M, Buchbinder R, Wluka AE, Page RS, et al. Associations between socioeconomic status and primary total knee joint replacements performed for osteoarthritis across Australia 2003–10: data from the Australian Orthopaedic Association National Joint Replacement Registry. *BMC Musculoskelet Disord*. 2014;15.
132. Peat G, Yu D, Grønne DT, Marshall M, Skou ST, Roos EM. Do patients with intersectional disadvantage have poorer outcomes from osteoarthritis management programmes? A tapered-balancing study of patient outcomes from the Good Life with osteoArthritis in Denmark (GLA:D®) programme. *Arthritis Care Res (Hoboken)*. 2022. <https://doi.org/10.1002/ACR.24987>.
133. Williams O, Fullagar S. Lifestyle drift and the phenomenon of ‘citizen shift’ in contemporary UK health policy. *Sociol Health Illn*. 2019;41:20–35.
134. World Health Organization. *World Health Report, Adherence to Long Term Therapies, Evidence for Action*. Geneva; 2003.
135. World Health Organization. *WHO Fact sheet on physical activity*. 2018.
136. Parker R, Bergman E, Mntambo A, Stubbs S, Wills M. Levels of physical activity in people with chronic pain. *S Afr J Physiother*. 2017;73.
137. Areerak K, Waongenngarm P, Janwantanakul P. Factors associated with exercise adherence to prevent or treat neck and low back pain: A systematic review. *Musculoskelet Sci Pract*. 2021;52:102333.
138. Merlo J, Mulinari S, Wemrell M, Subramanian S v., Hedblad B. The tyranny of the averages and the indiscriminate use of risk factors in public health: The case of coronary heart disease. *SSM Popul Health*. 2017;3:684–98.
139. Dell’Isola A, Jönsson T, Nero H, Eek F, Dahlberg L. Factors Associated With the Outcome of a First-Line Intervention for Patients With Hip or Knee Osteoarthritis or Both: Data From the BOA Register. *Phys Ther*. 2020;100:1771–81.

140. Ludvigsson JF, Svedberg P, Olén O, Bruze G, Neovius M. The longitudinal integrated database for health insurance and labour market studies (LISA) and its use in medical research. *Eur J Epidemiol.* 2019;34.
141. Statistics Sweden. Longitudinal integration database for health insurance and labour market studies (LISA).
142. Jönsson T, Eek F, Dell’Isola A, Dahlberg LE, Ekvall Hansson E. The Better Management of Patients with Osteoarthritis Program: Outcomes after evidence-based education and exercise delivered nationwide in Sweden. *PLoS One.* 2019;14:e0222657.
143. Dell’Isola A, Jönsson T, Ranstam J, Dahlberg LE, Ekvall Hansson E. Education, Home Exercise, and Supervised Exercise for People With Hip and Knee Osteoarthritis As Part of a Nationwide Implementation Program: Data From the Better Management of Patients With Osteoarthritis Registry. *Arthritis Care Res (Hoboken).* 2020;72:201–7.
144. Socialstyrelsen. Nationella riktlinjer för rörelseorganens sjukdomar. Reumatoid artrit, axial spondylartrit, psoriasisartrit, artros och osteoporos. 2021.
145. Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). *Arthritis Care Res (Hoboken).* 2011;63:S240–52.
146. Bjorgul K, Novicoff WM, Saleh KJ. Evaluating comorbidities in total hip and knee arthroplasty: available instruments. *J Orthop Traumatol.* 2010;11:203–9.
147. Lomi C. Evaluation of a Swedish Version of the Arthritis Self-efficacy Scale. *Scand J Caring Sci.* 1992;6:131–8.
148. Lorig K, Chastain RL, Ung E, Shoor S, Holman HR. Development and evaluation of a scale to measure perceived self-efficacy in people with arthritis. *Arthritis Rheum.* 1989;32:37–44.
149. Balestroni G, Bertolotti G. EuroQol-5D (EQ-5D): an instrument for measuring quality of life. *Monaldi Arch Chest Dis.* 2012;78:155–9.
150. Gustafsson K, Kvist J, Eriksson M, Dahlberg LE, Rolfson O. Socioeconomic status of patients in a Swedish national self-management program for osteoarthritis compared with the general population—a descriptive observational study. *BMC Musculoskeletal Disord.* 2020;21:10.
151. Valveny N, Gilliver S. How to interpret and report the results from multivariable analyses. *Medical Writing.* 2016;25:37.
152. Neely JG, Paniello RC, Cho Lieu JE, Voelker CCJ, Grindler DJ, Sequeira SM, et al. Practical guide to understanding multivariable analyses: Part A. *Otolaryngol Head Neck Surg.* 2013;148:185–90.

153. Komorowski M, Marshall DC, Saliccioli JD, Crutain Y. Chapter 15 - Exploratory Data Analysis. In: Secondary Analysis of Electronic Health Records. Springer; 2016. p. 185–203.
154. Braveman P, Gottlieb L. The Social Determinants of Health: It's Time to Consider the Causes of the Causes. *Public Health Reports*. 2014;129 Suppl 2:19.
155. Islam MM. Social Determinants of Health and Related Inequalities: Confusion and Implications. *Front Public Health*. 2019;7 FEB:11.
156. Shafiei S, Yazdani S, Jadidfard MP, Zafarmand AH. Measurement components of socioeconomic status in health-related studies in Iran. *BMC Res Notes*. 2019;12.
157. Shavers VL. Measurement of socioeconomic status in health disparities research. *J Natl Med Assoc*. 2007;99:1013.
158. Kawachi I, Subramanian S V., Almeida-Filho N. A glossary for health inequalities. *J Epidemiol Community Health* (1978). 2002;56:647–52.
159. Whitehead M. The concepts and principles of equity and health. *Int J Health Serv*. 1992;22:429–45.
160. World Health Organization. WHO. Health Impact Assessment, glossary of terms. <https://www.who.int/publications/m/item/glossary-of-terms-used-for-health-impact-assessment-hia>. Accessed 15 Nov 2022.
161. Cooper C, Chapurlat R, Al-Daghri N, Herrero-Beaumont G, Bruyère O, Rannou F, et al. Safety of Oral Non-Selective Non-Steroidal Anti-Inflammatory Drugs in Osteoarthritis: What Does the Literature Say? *Drugs Aging*. 2019;36 Suppl 1:15.
162. Marmot M. The influence of income on health: views of an epidemiologist. *Health Aff (Millwood)*. 2002;21:31–46.
163. Zhang S, Xiang W. Income gradient in health-related quality of life - The role of social networking time. *Int J Equity Health*. 2019;18:1–10.
164. Galobardes B, Shaw M, Lawlor DA, Lynch JW, Davey Smith G. Indicators of socioeconomic position (part 1). *J Epidemiol Community Health*. 2006;60:7–12.
165. Ludvigsson JF, Svedberg P, Olén O, Bruze G, Neovius M. The longitudinal integrated database for health insurance and labour market studies (LISA) and its use in medical research. *Eur J Epidemiol*. 2019;34:423.
166. Wagstaff A, Paci P, van Doorslaer E. On the measurement of inequalities in health. *Soc Sci Med*. 1991;33:545–57.
167. O'Donnell O, O'Neill S, Van Ourti T, Walsh B. conindex: Estimation of concentration indices. *Stata J*. 2016;16:112.
168. Kjellsson G, Gerdtham UG. On correcting the concentration index for binary variables. *J Health Econ*. 2013;32:659–70.

169. Madden D. Ordinal and cardinal measures of health inequality: An empirical comparison. *Health Econ.* 2010;19:243–50.
170. Erreygers G. Correcting the concentration index. *J Health Econ.* 2009;28:504–15.
171. Holden MA, Nicholls EE, Young J, Hay EM, Foster NE. UK-based physical therapists' attitudes and beliefs regarding exercise and knee osteoarthritis: Findings from a mixed-methods study. *Arthritis Care Res (Hoboken).* 2009;61:1511–21.
172. Morin Chabane S, Coutinho F, Laliberte M, Feldman D. Outpatient physiotherapists' attitudes and beliefs toward patients with chronic pain: A qualitative study. *Physiother Theory Pract.* 2020;36:85–94.
173. Fischer F, Lange K, Klose K, Greiner W, Kraemer A. Barriers and Strategies in Guideline Implementation—A Scoping Review. *Healthcare.* 2016;4:36.
174. Gunn AH, Schwartz TA, Arbeeva LS, Callahan LF, Golightly Y, Goode A, et al. Fear of Movement and Associated Factors Among Adults With Symptomatic Knee Osteoarthritis. *Arthritis Care Res (Hoboken).* 2017;69:1826–33.
175. Briggs AM, Houlding E, Hinman RS, Desmond LA, Bennell KL, Darlow B, et al. Health professionals and students encounter multi-level barriers to implementing high-value osteoarthritis care: a multi-national study. *Osteoarthritis Cartilage.* 2019;27:788–804.
176. Geri T, Viceconti A, Minacci M, Testa M, Rossetini G. Manual therapy: Exploiting the role of human touch. *Musculoskelet Sci Pract.* 2019;44:102044.
177. Doyle C, Lennox L, Bell D. A systematic review of evidence on the links between patient experience and clinical safety and effectiveness. *BMJ Open.* 2013;3:e001570.
178. Lillrank A. Back pain and the resolution of diagnostic uncertainty in illness narratives. *Soc Sci Med.* 2003;57:1045–54.
179. Lütze U, Archenholtz B. The impact of arthritis on daily life with the patient perspective in focus. *Scand J Caring Sci.* 2007;21:64–70.
180. Etkind SN, Bristowe K, Bailey K, Selman LE, Murtagh FE. How does uncertainty shape patient experience in advanced illness? A secondary analysis of qualitative data. *Palliat Med.* 2017;31:171.
181. Chou L, Ellis L, Papandony M, Seneviwickrama KLMD, Cicuttini FM, Sullivan K, et al. Patients' perceived needs of osteoarthritis health information: A systematic scoping review. *PLoS One.* 2018;13:e0195489.
182. Riess H. The Science of Empathy. *J Patient Exp.* 2017;4:77.
183. Vitaloni M, Botto-van Bemden A, Sciortino R, Carné X, Quintero M, Santos-Moreno P, et al. A patients' view of OA: the Global Osteoarthritis Patient Perception Survey (GOAPPS), a pilot study. *BMC Musculoskelet Disord.* 2020;21:727.

184. Escoto KH, Laska MN, Larson N, Neumark-Sztainer D, Hannan PJ. Work Hours and Perceived Time Barriers to Healthful Eating Among Young Adults. *Am J Health Behav.* 2012;36:786.
185. Withall J, Jago R, Fox KR. Why some do but most don't. Barriers and enablers to engaging low-income groups in physical activity programmes: a mixed methods study. *BMC Public Health.* 2011;11:507.
186. Herazo-Beltrán Y, Pinillos Y, Vidarte J, Crissien E, Suarez D, García R. Predictors of perceived barriers to physical activity in the general adult population: a cross-sectional study. *Braz J Phys Ther.* 2017;21:44.
187. Ford E, Zhao G, JTsai J, Li C. Low-risk lifestyle behaviors and all-cause mortality: findings from the National Health and Nutrition Examination Survey III Mortality Study. *Am J Public Health.* 2011;101:1922–9.
188. Skou ST, Roos EM. Good Life with osteoArthritis in Denmark (GLA:D™): evidence-based education and supervised neuromuscular exercise delivered by certified physiotherapists nationwide. *BMC Musculoskelet Disord.* 2017;18:1–13.
189. Kiadaliri A, Dell'isola A, Lohmander LS, Hunter DJ, Dahlberg LE, Se AKL. Assessing the importance of predictors of adherence to a digital self-management intervention for osteoarthritis. *Journal of Orthopaedic Surgery and Research* 2023 18:1. 2023;18:1–10.
190. Gommans LNM, Scheltinga MRM, van Sambeek MRHM, Maas AHM, Bendermacher BLW, Tejjink JAW. Gender differences following supervised exercise therapy in patients with intermittent claudication. *J Vasc Surg.* 2015;62:681–8.
191. Oosenbrug E, Marinho RP, Zhang J, Marzolini S, Colella TJF, Pakosh M, et al. Sex Differences in Cardiac Rehabilitation Adherence: A Meta-analysis. *Canadian Journal of Cardiology.* 2016;32:1316–24.
192. McArthur D, Dumas A, Woodend K, Beach S, Stacey D. Factors influencing adherence to regular exercise in middle-aged women: a qualitative study to inform clinical practice. *BMC Womens Health.* 2014;14:49.
193. Verhoef M, Love EJ, Rose SA. Women's social roles and their exercise participation. *Women Health.* 1992;19:15–29.
194. Cheung C, Wyman JF, Peden-McAlpine C. Long-Term Yoga and Aerobic/Strength Exercise Adherence in Older Women with Knee Osteoarthritis: A Mixed Methods Approach. *Int J Yoga Therap.* 2022;32.
195. Lee HH, Dunsiger S, Connell Bohlen L, Boyle HK, Emerson JA, Williams DM. Age Moderates the Effect of Self-Paced Exercise on Exercise Adherence among Overweight Adults. *J Aging Health.* 2020;32:154–61.

196. Baillot A, Chenail S, Polita NB, Simoneau M, Libourel M, Nazon E, et al. Physical activity motives, barriers, and preferences in people with obesity: A systematic review. *PLoS One*. 2021;16:e0253114.
197. Tuakli-Wosornu YA, Selzer F, Losina E, Katz JN. Predictors of Exercise Adherence in Patients With Meniscal Tear and Osteoarthritis. *Arch Phys Med Rehabil*. 2016;97:1945–52.
198. Kahan BC, Rehal S, Cro S. Risk of selection bias in randomised trials. *Trials*. 2015;16.
199. van Onzenoort HAW, Menger FE, Neef C, Verberk WJ, Kroon AA, de Leeuw PW, et al. Participation in a clinical trial enhances adherence and persistence to treatment: a retrospective cohort study. *Hypertension*. 2011;58:573–8.
200. Collado-Mateo D, Lavín-Pérez AM, Peñacoba C, del Coso J, Leyton-Román M, Luque-Casado A, et al. Key Factors Associated with Adherence to Physical Exercise in Patients with Chronic Diseases and Older Adults: An Umbrella Review. *Int J Environ Res Public Health*. 2021;18:1–24.
201. Essery R, Geraghty AWA, Kirby S, Yardley L. Predictors of adherence to home-based physical therapies: a systematic review. *Disabil Rehabil*. 2017;39:519–34.
202. Daniali SS, Darani FM, Eslami AA, Mazaheri M. Relationship between Self-efficacy and Physical Activity, Medication Adherence in Chronic Disease Patients. *Adv Biomed Res*. 2017;6:63.
203. Bandura A. *Self-efficacy: The exercise of control*. New York, NY: Freeman; 1997.
204. Bachrach DG, Rapp TL, Rapp AA, Ogilvie J. “Too Much” Self-Efficacy? Understanding the Curvilinear Consequences of Between-Person Self-Efficacy through a Moderated-Mediation Model of Perceived Proximity and Employee Effort. *Group Organ Manag*. 2022;2022:1–38.
205. Peters TJ, Sanders C, Dieppe P, Donovan J. Factors associated with change in pain and disability over time: a community-based prospective observational study of hip and knee osteoarthritis. *The British Journal of General Practice*. 2005;55:205.
206. Cleveland RJ, Luong MLN, Knight JB, Schoster B, Renner JB, Jordan JM, et al. Independent associations of socioeconomic factors with disability and pain in adults with knee osteoarthritis. *BMC Musculoskelet Disord*. 2013;14.
207. Jhun HJ, Sung NJ, Kim SY. Knee pain and its severity in elderly Koreans: prevalence, risk factors and impact on quality of life. *J Korean Med Sci*. 2013;28:1807–13.
208. Dell’Isola A, Turkiewicz A, Jönsson T, Rolfson O, Dahlberg L, Englund M. The role of pain and walking difficulties in shaping willingness to undergo joint surgery for osteoarthritis: Data from the Swedish BOA register. *Osteoarthr Cartil Open*. 2021;3:100157.

209. Dell'Isola A, Jönsson T, Rolfson O, Cronström A, Englund M, Dahlberg L. Willingness to Undergo Joint Surgery Following a First-Line Intervention for Osteoarthritis: Data From the Better Management of People With Osteoarthritis Register. *Arthritis Care Res (Hoboken)*. 2021;73:818–27.
210. Judge A, Welton NJ, Sandhu J, Ben-Shlomo Y. Geographical variation in the provision of elective primary hip and knee replacement: the role of socio-demographic, hospital and distance variables. *J Public Health (Oxf)*. 2009;31:413–22.
211. Dieppe P, Judge A, Williams S, Ikwueke I, Guenther KP, Floeren M, et al. Variations in the pre-operative status of patients coming to primary hip replacement for osteoarthritis in European orthopaedic centres. *BMC Musculoskelet Disord*. 2009;10:19.
212. Bonnesen K, Ehrenstein V, Grønkjær MS, Pedersen L, Lash TL, Schmidt M. Impact of lifestyle and socioeconomic position on use of non-steroidal anti-inflammatory drugs: A population-based cohort study. *Pharmacoepidemiol Drug Saf*. 2022. <https://doi.org/10.1002/PDS.5571>.
213. Nielsen MW, Hansen EH, Rasmussen NK. Prescription and non-prescription medicine use in Denmark: Association with socio-economic position. *Eur J Clin Pharmacol*. 2003;59:677–84.
214. Fosbøl EL, Gislason GH, Jacobsen S, Abildstrom SZ, Hansen ML, Schramm TK, et al. The pattern of use of non-steroidal anti-inflammatory drugs (NSAIDs) from 1997 to 2005: A nationwide study on 4.6 million people. *Pharmacoepidemiol Drug Saf*. 2008;17:822–33.
215. Altman DG, Royston P. The cost of dichotomising continuous variables. *BMJ*. 2006;332:1080.
216. Oakes J, Kaufam J, editors. *Methods in Social Epidemiology*, Chapter 6. 2nd edition. Wiley; 2017.
217. Sim J, Saunders B, Waterfield J, Kingstone T. Can sample size in qualitative research be determined a priori? <https://doi.org/10.1080/1364557920181454643>. 2018;21:619–34.
218. Messier SP, Beavers DP, Queen K, Mihalko SL, Miller GD, Losina E, et al. Effect of Diet and Exercise on Knee Pain in Patients With Osteoarthritis and Overweight or Obesity: A Randomized Clinical Trial. *JAMA*. 2022;328:2242–51.
219. Messier SP, Mihalko SL, Beavers DP, Nicklas BJ, Devita P, Carr JJ, et al. Effect of High-Intensity Strength Training on Knee Pain and Knee Joint Compressive Forces Among Adults With Knee Osteoarthritis: The START Randomized Clinical Trial. *JAMA*. 2021;325:646–57.

220. Bandak E, Christensen R, Overgaard A, Kristensen LE, Ellegaard K, Guldborg-Møller J, et al. Exercise and education versus saline injections for knee osteoarthritis: a randomised controlled equivalence trial. *Ann Rheum Dis.* 2022;81:537–43.
221. Boswell MA, Evans KM, Zion SR, Boles DZ, Hicks JL, Delp SL, et al. Mindset is associated with future physical activity and management strategies in individuals with knee osteoarthritis. *Ann Phys Rehabil Med.* 2022;65.
222. Crum AJ, Salovey P, Achor S. Rethinking stress: the role of mindsets in determining the stress response. *J Pers Soc Psychol.* 2013;104:716–33.
223. Cinthuja P, Krishnamoorthy N, Shivapatham G. Effective interventions to improve long-term physiotherapy exercise adherence among patients with lower limb osteoarthritis. A systematic review. *BMC Musculoskelet Disord.* 2022;23:1–11.
224. Willett M, Duda J, Gautrey C, Fenton S, Greig C, Rushton A. Effectiveness of behavioural change techniques in physiotherapy interventions to promote physical activity adherence in patients with hip and knee osteoarthritis: a systematic review protocol. *BMJ Open.* 2017;7:e015833.
225. Smith KE, Eltanani MK. What kinds of policies to reduce health inequalities in the UK do researchers support? *J Public Health (Oxf).* 2015;37:6–17.
226. Bouraoui A, Rutter M, Williamson L, Fisher C, Sofat R, Sen D. The urgent need to move upstream in caring for people with rheumatic and musculoskeletal diseases. *Rheumatol Adv Pract.* 2022;6.
227. McMahon NE. Framing action to reduce health inequalities: what is argued for through use of the ‘upstream–downstream’ metaphor? *J Public Health (Bangkok).* 2022;44:671–8.

Acknowledgements (English)

This chapter is of great importance to me as gratitude towards all the people who have supported, guided and advised me has been the constant of my journey.

I would like to start by thanking my supervisors, each for different reasons, often overlapping. I thank **Dr Marco Testa** because defining him as a guide is an understatement. I thank him for always listening to me and guiding me and for making me understand that it's not the final goal that matters but how you feel while trying to achieve it. I thank **Dr Andrea Dell'Isola** for providing the tools to become a good researcher and constantly pushing me to ask questions rather than provide answers. I thank **Prof Martin Englund** for overseeing every phase of my PhD, providing the right advice at the right time.

I would like to thank my mother, **Elena Nascimbene**. Among the many traits she passed on to me that make me similar to her, passion, curiosity, and the pursuit of answers have been pillars in my formation. I thank my father, **Pierpaolo Battista** because he is the certainty in this world of question marks. I thank my brother, **Davide Battista**, for always bring there. I thank my grandmother, **Giuseppina Pusterla**, because she has taught me more than any book or research I will ever read. I thank my grandmother, **Enrica Dell'Acqua** because her passion for physiotherapy and medicine has flowed through generations, reaching me. And I thank my grandfathers, **Costantino Battista** and **Luigi Nascimbene** because they are always by my side.

I thank my travel companions from Italy. **Mirko Job**, **Lucia Grazia Sansone**, **Antonello Viceconti**, **Mattia Manoni**, **Gaia Leuzzi**, **Luca Francini**, **Gianluca Bertoni**, and **Evelin Marchesini** because without them, the **REHELab** (and myself) would not be what we are today. I thank the entire **Clinical Epidemiology Unit** of Lund University because they have always made me feel part of a group, even when kilometres apart. Three special thanks go to **Dr Ali Kiadaliri** and **Dr Aleksandra Turkiewicz** because I would have never believed I would fall in love with numbers and statistics before knowing them, and to **Liselotte Højgård Hansen** for her ability to take care of others.

I can only thank the people who have been closest to me (also) outside of academic life. I start with **Benedetto Giardulli**, roommate and chosen brother, because I have never seen a stronger form of gentleness. I continue with my other roommate and chosen sister, **Ilaria Ghiso** because certain people are destined to enter in your life. I thank **Ludovica Paravelli** and **Fanny Mariette** for being the constant in my life. I thank **Carola Cosentino** and **Ilaria Coppola** because it is not about how much time we've spent together but how we've spent it. I thank **Prof Yeliz Prior** because she taught me to respond to every call that excites my spirit. I thank **Dr Marialuisa Gandolfi** because all this would not have been possible without her. I thank all the physiotherapists and doctors at the **San Paolo Hospital** in Milan because they give me the reason to start this journey years ago. I thank **Raffaele Cutolo** because our paths may have diverged, but the portion of the road we travelled together will not be forgotten. I thank **Irsida Zaimaj** for making me embrace what I thought it was my greatest enemy. I thank **Sunny Gurumayum** and **Luca Buzzatti** for being family during my time abroad. I thank **Ezia Zulfarino** for believing in me before I even started to do so. I thank my lifelong friends, **Edoardo Boidi, Federica Puppo, Valerio Bianchi, Marco Giuliani, Allegra Marletti, Sara Milia, Melissa Pellizzoni, Sara Federico, Ioana Gaga, Federica Fullone, Zambotti Jessica, Simone Cecchetto, Jennifer Barteloni** and **Davide Marazzi** because they witnessed the departure, not just the arrival.

I thank each one of you because, thanks to you, the person writing these acknowledgements is not the same as four years ago. I also thank the people who will become part of my life in the forthcoming years for what they will teach me. These years have given me so much, but I am only eager to discover what lies ahead.

Thank you.

Acknowledgements (Italian)

Questo capitolo è per me di grande importanza in quanto la gratitudine verso tutte le persone che mi hanno seguito, indirizzato, consigliato in questi quattro anni è stata la costante che mi ha accompagnato in questo viaggio.

Vorrei iniziare ringraziando i miei supervisor, ognuno per dei motivi diversi, spesso sovrapponibili. Ringrazio il **Dr Marco Testa** perché definirlo una guida è riduttivo. Lo ringrazio per avermi sempre ascoltato e indirizzato, e avermi fatto comprendere che non è l'obiettivo finale ciò che importa, ma come ti senti mentre cerchi di raggiungerlo. Ringrazio il **Dr Andrea Dell'Isola** per avermi fornito gli strumenti per diventare un bravo ricercatore ed avermi sempre spinto a farmi delle domande piuttosto che a dare risposte. Ringrazio il **Prof Martin Englund** per aver sorvegliato ogni fase, dando i giusti consigli nei momenti opportuni.

Vorrei ringraziare mia madre, **Elena Nascimbene**, perché delle tante caratteristiche che mi ha passato e che mi rendono simile a lei, la passione, la curiosità e la ricerca di risposte sono stati dei pilastri nella mia formazione. Ringrazio mio padre, **Pierpaolo Battista**, perché è certezza in questo mondo di punti interrogativi. Ringrazio mio fratello, **Davide Battista**, per esserci sempre stato. Ringrazio mia nonna **Giuseppina Pusterla**, perché mi ha insegnato più di ogni libro/ricerca che potrò mai leggere. Ringrazio mia nonna **Enrica Dell'Acqua**, perché la sua passione per la fisioterapia e la medicina l'ha fatta scorrere di generazione in generazione, arrivando a me. E ringrazio i miei nonni, **Costantino Battista** e **Luigi Nascimbene**, perché, in qualche modo, sono sempre al mio fianco.

Ringrazio i miei compagni di viaggio dall'Italia. **Mirko Job**, **Lucia Grazia Sansone**, **Antonello Viceconti**, **Mattia Manoni**, **Gaia Leuzzi**, **Luca Francini**, **Gianluca Bertoni** ed **Evelin Marchesini** perché senza di loro il REHELab (e io stesso) non saremmo ciò che siamo oggi. Ringrazio tutta la **Clinical Epidemiology Unit** dell'Università di Lund perché mi ha sempre fatto sentire parte di un gruppo anche a chilometri di distanza. Tre ringraziamenti speciali vanno al **Dr Ali Kiadaliri** e alla **Dr.ssa Aleksandra Turkiewicz** perché mai avrei creduto che mi sarei innamorato dei numeri e della statistica prima di conoscerli e a **Liselotte Højgård Hansen** per la sua capacità di prendersi cura dell'altro.

Non posso che ringraziare le persone che mi sono state vicine anche all'esterno della vita accademica. Inizio con **Benedetto Giardulli**, coinquilino e fratello scelto, perché non ho mai visto una forma di delicatezza più forte. Proseguo con l'altra mia sorella scelta e coinquilina, **Ilaria Ghiso**, perché certe persone sono destinate a entrare nella tua vita. Ringrazio **Ludovica Paravelli** e **Fanny Mariette** per essere le costanti della mia vita. Ringrazio **Carola Cosentino** e **Ilaria Coppola** perché non è quanto tempo si è passato insieme, ma come lo si è passato. Ringrazio la **Prof.ssa Yeliz Prior** perché mi ha insegnato a rispondere a qualunque chiamata che faccia risuonare la mia anima. Ringrazio la **Dr.ssa Marialuisa Gandolfi** perché, senza di lei, tutto questo non sarebbe iniziato. Ringrazio tutte/i le/i fisioterapiste/i e medici **dell'U.O. dell'Ospedale San Paolo di Milano** per aver inserito il primo tassello. Ringrazio **Raffaele Cutolo** perché le strade si saranno divise, ma il pezzo di strada percorso assieme non sarà dimenticato. Ringrazio **Irsida Zaimaj** per avermi fatto abbracciare colui che pensavo essere il mio nemico più grande. Ringrazio **Sunny Gurumayum** e **Luca Buzzatti** per essere stati famiglia durante i miei periodi all'estero. Ringrazio **Ezia Zulfarino** per aver creduto in me prima ancora che io iniziassi a farlo. Ringrazio i compagni e le compagne di una vita, **Edoardo Boidi, Federica Puppo, Valerio Bianchi, Marco Giuliani, Allegra Marletti, Sara Milia, Melissa Pellizzoni, Sara Federico, Ioana Gaga, Federica Fullone, Zambotti Jessica, Simone Cecchetto, Jennifer Barteloni** e **Davide Marazzi** perché hanno visto la partenza e non solo l'arrivo.

Ringrazio anche le persone che entreranno a far parte della mia vita nei prossimi anni per ciò che mi insegneranno. Questi anni mi avranno anche dato tantissimo, ma non posso che essere impaziente di scoprire cosa ci sarà dopo. Vi ringrazio tutti e tutte perché per merito vostro, colui che sta scrivendo questi ringraziamenti, non è la stessa persona di quattro anni fa.

Grazie.

Manuscripts

Study 1

RESEARCH ARTICLE

Open Access

Italian physiotherapists' knowledge of and adherence to osteoarthritis clinical practice guidelines: a cross-sectional study



Simone Battista^{1,2}, Stefano Salvioli¹, Serena Millotti¹, Marco Testa¹ and Andrea Dell'Isola^{2*}

Abstract

Introduction: Implementation of clinical practice guidelines (CPGs) to manage musculoskeletal conditions among physiotherapists appears suboptimal. Osteoarthritis is one of the most disabling conditions worldwide and several studies showed a lack of knowledge of and adherence to osteoarthritis CPGs in physiotherapists' clinical practice. However, those studies are not conclusive, as they examine the knowledge of and adherence to CPGs only in isolation, or only by focussing on a single treatment. Thus, analysis of the knowledge of and adherence to CPGs in the same sample would allow for a better understanding of the evidence-to-practice gap, which, if unaddressed, can lead to suboptimal care for these patients. This study aims at assessing Italian physiotherapists' evidence-to-practice gap in osteoarthritis CPGs.

Methods: An online survey divided into two sections investigating knowledge of and adherence to CPGs was developed based on three high-quality, recent and relevant CPGs. In the first section, participants had to express their agreement with 24 CPG statements through a 1 (completely disagree) to 5 (completely agree) scale. We defined a $\geq 70\%$ agreement with a statement as consensus. In the second section, participants were shown a clinical case, with different interventions to choose from. Participants were classified as 'Delivering' (all recommended interventions selected), 'Partially Delivering' (some recommended interventions missing) and 'Non-Delivering' (at least one non-recommended interventions selected) the recommended intervention, depending on chosen interventions.

Results: 822 physiotherapists (mean age (SD): 35.8 (13.3); female 47%) completed the survey between June and July 2020. In the first section, consensus was achieved for 13/24 statements. In the second section, 25% of the participants were classified as 'Delivering', 22% as 'Partially Delivering' and 53% as 'Non-Delivering'.

(Continued on next page)

* Correspondence: andrea.dellisola@med.lu.se

²Clinical Epidemiology Unit, Orthopedics, Department of Clinical Sciences
Lund, Lund University, Lund, Sweden

Full list of author information is available at the end of the article



© The Author(s). 2021, corrected publication 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

Conclusions: Our findings revealed an adequate level of knowledge of osteoarthritis CPGs regarding the importance of exercise and education. However, an adequate level of adherence has yet to be reached, since many physiotherapists did not advise weight reduction, but rest from physical activity, and often included secondary treatments (e.g. manual therapy) supported by low-level evidence. These results identify an evidence-to-practice gap, which may lead to non-evidence based practice behaviours for the management of patients with osteoarthritis.

Keywords: Osteoarthritis, Osteoarthritis, knee, Osteoarthritis, hip, Practice guidelines as topic, Clinical governance, Physical therapy specialty, Physical therapists, Education, public health professional

Introduction

Osteoarthritis is the most prevalent joint disease, and one of the most common causes of disability worldwide [1], for whose management several international Clinical Practice Guidelines (CPGs) have been released [2–4]. CPGs are collections of graded recommendations supported by a systematic review of evidence, intended to help clinicians to optimise patient care [5]. In osteoarthritis, the European League Against Rheumatism (EULAR), Osteoarthritis Research Society International (OARSI) and the National Institute for Health and Care Excellence (NICE) CPGs [2–4] recommend exercise and education [6] as first-line interventions for their ability to reduce pain and disability, regardless of the severity of the disease [7, 8].

Despite the differences in grading the strength of recommendations between them, all abovementioned CPGs categorised the level of evidence, of the different treatments, into six categories from Ia to IV [2–4]. Treatments categorised as Ia are derived from systematic reviews of randomised controlled trials (RCTs) [2–4, 9]; whereas, treatments categorised from IIa to IV are based on lower quality RCTs, cohort studies or the opinion of experts in the field [2–4, 9]. The level of evidence in all CPGs is one of the factors that contribute to defining the strength of the recommendations together with the balance between benefits and harms, considerations of values and preferences, and resources implications [10]. By taking into account all these factors, treatments such as therapeutic exercise and patients education (Ia, level of evidence) are categorised as first-line interventions, whereas treatments such as hyaluronic acid injection and manual therapy (IV, level of evidence) are categorised as conditional recommendations [2–4].

In spite of the availability of several high-quality osteoarthritis CPGs, Egerton et al. highlighted that clinicians who work with patients with osteoarthritis perceived themselves as under-prepared and unfamiliar with CPGs [11]. Besides, implementation of CPGs for the management of musculoskeletal conditions among physiotherapists appears suboptimal [12] and osteoarthritis often remains under-diagnosed and under-treated, with less

than 40% of people with osteoarthritis receiving the recommended first-line intervention [13, 14].

Several studies have explored physiotherapists' knowledge of and adherence to osteoarthritis CPGs, showing major gaps in the implementation of weight reduction strategies, therapeutic exercise and patients education, as opposed to what has been noticed for other passive modalities (e.g. manual therapy) [15–19]. The reasons behind this gap are several. First of all, the care of people with knee and hip osteoarthritis is a complex process in which the clinician is required to balance knowledge of the best evidence with the patients' preferences and beliefs [20]. Secondly, clinicians may face several barriers to the implementation of CPGs, such as clinical applicability, language and lack of time, which can widen this gap even further [21–23].

However, the aforementioned studies on the application of osteoarthritis CPGs are not conclusive, as they either examined the knowledge of or adherence to CPGs, [16–19] or focus solely on a particular treatment (e.g. therapeutic exercise) [15]. In fact, knowledge of the CPGs does not automatically translate into clinical practice [24, 25], thus an analysis of the knowledge of and adherence to CPGs in the same sample would allow for a better understanding of the so-called *evidence-to-practice gap* [21].

In line with this, this study aimed at exploring the knowledge of and adherence to osteoarthritis CPGs in a cohort of Italian physiotherapists in order to identify the possible evidence-to-practice gap. By analysing this gap in Italy, this study gathered information that might be more easily transferred to other Mediterranean countries which seem to have higher educational needs compared to the Northern-European ones [26].

Methods

Study design

A quantitative web-based cross-sectional survey investigating physiotherapists' knowledge of and adherence to osteoarthritis CPGs was developed according to the *International Handbook of Survey Methodology* and to the *Checklist for Reporting Results of Internet E-Survey*

through the use of distinct and iterative steps [27, 28]. The study was conducted following the Declaration of Helsinki. Ethical approval was obtained from the Ethics Committee for University Research (CERA: Comitato Etico per la Ricerca di Ateneo), University of Genova (approval date: 15/06/2020; CERA2020.07) and follows the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) recommendations for reporting observational studies [29].

Survey development

The questionnaire was developed based on the EULAR, OARSI and NICE CPGs [2–4]. Before the online dissemination, the survey was tested on a sample of six physiotherapists specialised in musculoskeletal rehabilitation. The online version of the questionnaire was delivered through *Microsoft 365 Forms*, a secure web application to build and manage online surveys and databases, respecting the European General Data Protection Regulations [30]. The questionnaire included a brief cover letter, and the informed consent outlining the aim of the study. The cover letter emphasised that participation in the survey was voluntary and that anonymity and confidentiality were guaranteed.

The questionnaire was delivered in Italian and it was divided into two sections investigating (1) the knowledge of and (2) the adherence to osteoarthritis CPGs. Before the first section, a paragraph investigating the socio-demographic variables (e.g. sex, age, years of experience) and if the physiotherapists read, at least, one osteoarthritis CPG, was included. The first section comprised 24 statements on knee and hip osteoarthritis management, adapted from the aforementioned CPGs. Each statement was acquired from the synoptic review of the three CPGs (Table 1).

If disagreement was found between the CPGs, the most recent recommendation was considered when phrasing the statement. To measure agreement with the statement, we used a 5-point Likert scale ranging from completely disagree (score 1) to completely agree (score 5) [31]. Participants who partially or completely agreed (scores 4–5) were considered to agree with the statements. Furthermore, to limit acquiescence bias, i.e. the tendency to agree with all the survey statements, eleven reversed statements were put into the questionnaire so that disagreement with those statements (scores 1–2) would indicate an agreement with the CPGs [32].

The second section presented a clinical vignette illustrating a case of knee osteoarthritis (Table 2). Clinical vignettes are considered a valid tool to assess healthcare professionals' clinical reasoning and behaviour, including physiotherapists, as they are easy to administer, and all the variables within them can be easily manipulated [33].

The participants were, therefore, invited to express how they would manage that specific patient by selecting from a list of options. The options were grouped into three sub-sections representing three different clinical moments (management, assessment and treatment). In the management section, the participants were asked whether they would opt to treat the presented patient or refer her to a specialist for pharmacological or surgical treatment. Participants who decided to refer the patient to a specialist without performing any assessment or treatment were directed to the end of the questionnaire, considered non-delivering any possible clinical options.

Participants

An online version of the questionnaire, attainable through a hyperlink, was delivered through the Italian Association of Physiotherapists (AIFI) and the University of Genova newsletters. We included physiotherapists who had treated at least one person with osteoarthritis in the previous six months within the targeted population. To do so, after the cover letter, the questionnaire included a preliminary question asking the respondents if they had treated any patients, with hip or knee osteoarthritis in the last six months. Participants who answered "No" were shown a Thank-You page and were not allowed to continue the questionnaire.

Variables

The primary outcome of the present study was the level of knowledge of and adherence to CPGs of a sample of Italian physiotherapists.

Analysis

Descriptive analysis was carried out to understand the sample's characteristics. Moreover, the frequencies of physiotherapists who declared to have read at least one osteoarthritis CPG were reported.

Section 1: level of knowledge of clinical practice guidelines – statement consensus

In section one, participants who partially or completely agreed with a statement (scores 4–5) or partially or completely disagreed (scores 1–2) with a reversed statement were considered to agree with the CPGs recommendation. The overall consensus with each statement was investigated. In the absence of a standard threshold, we defined a $\geq 70\%$ agreement with a statement as consensus [34].

Section 2: level of adherence to clinical practice guidelines – clinical vignette

In section two, participants were classified as 'Delivering', 'Partially Delivering' and 'Non-Delivering' the core treatments depending on the interventions chosen.

Table 1 Section 1: Statements and synoptic review of Clinical Practice Guidelines

| Statements | Clinical Practice Guidelines |
|--|--|
| 1) Exercise can be effective on all patients, regardless of the pain severity. | NICE (1.2.5–1.4.1) [4]; EULAR (3–6–7) [2]; OARSI (Tables 2–3) [3] |
| 2) In an advanced stage of the disease, exercise can damage the joint (reversed statement). | NICE (1.2.5–1.4.1) [4]; EULAR (2–3–6–7) [2]; OARSI (Tables 2–3) [3] |
| 3) The rehabilitation programme must always include a part of education on the pathophysiology of osteoarthritis and self-management strategies. | NICE (1.3.1–1.3.2–1.3.3) [4]; EULAR (3–5) [2]; OARSI (Tables 2–3) [3] |
| 4) The rehabilitation programme should always include a part of manual treatment (reversed statement) | NICE (1.4.2) [4]; EULAR (–); OARSI (–) |
| 5) Exercise should only be undertaken after prescribing drug treatment to control pain (reversed statement). | NICE (1.2.5–1.4.1) [4]; EULAR (3–6–7) [2]; OARSI (Tables 2–3) [3] |
| 6) The use of topical anti-inflammatory drugs is effective for pain relief for knee osteoarthritis. | NICE (1.5.3) [4]; EULAR (–); OARSI (Table 2) [3] |
| 7) Radiographic findings are needed to express a functional diagnosis of osteoarthritis (reversed statement). | NICE (1.1.1) [4]; EULAR (1) [2]; OARSI (–) |
| 8) Radiographic findings are needed to plan the physiotherapy treatment (reversed statement). | NICE (1.1.1) [4]; EULAR (1) [2]; OARSI (–) |
| 9) Physical activity should be avoided because it can damage the joint (reversed statement). | NICE (1.2.5–1.4.1) [4]; EULAR (3–6–7) [2]; OARSI (Tables 2–3) [3] |
| 10) The use of topical anti-inflammatory drugs is effective for pain relief for hip osteoarthritis. | NICE (–); EULAR (–); OARSI (–) |
| 11) In case of severe joint degeneration, it is necessary to recommend rest from physical activity (reversed statement). | NICE (1.2.5–1.4.1) [4]; EULAR (2–3–6–7) [2]; OARSI (Tables 2–3) [3] |
| 12) In cases of severe pain (VAS \geq 6/10), arthroplasty surgery should be preferred to rehabilitation (reversed statement). | NICE (1.6) [4]; EULAR (–); OARSI (–) |
| 13) The use of TENS should be considered. | NICE (1.4.4) [4]; EULAR (–); OARSI (–) |
| 14) The use of physical therapies such as lasers, TECAR and ultrasound therapy should be considered (reversed statement). | NICE (1.4.4) [4]; EULAR (–); OARSI (–) |
| 15) In addition to the rehabilitation treatment, it is useful to recommend physical activity (for example, yoga, swimming, Nordic walking). | NICE (1.2.5–1.3.2–1.4.1) [4]; EULAR (–); OARSI (Tables 2–3) [3] |
| 16) It is important to recommend weight loss to overweight or obese patients. | NICE (1.2.5–1.4.3) [4]; EULAR (3–8) [2]; OARSI (Tables 2–3) [3] |
| 17) Age > 45, pain and absence of joint stiffness (or < 30 min) in the morning are sufficient to diagnose osteoarthritis. | NICE (1.1.1) [4]; EULAR (–); OARSI (–) |
| 18) The use of comfortable footwear, braces or aids should be considered. | NICE (1.3.2–1.4.7–1.4.8–1.4.9) [4]; EULAR (3–9–10) [2]; OARSI (Tables 2–3) [3] |
| 19) It is advisable to refer the patient for arthroscopy surgery to reduce symptoms and start/continue treatment (reversed statement). | NICE (1.4.10) [4]; EULAR (–); OARSI (–) |
| 20) It is necessary to assess the impact of osteoarthritis on function, quality of life and disability. | NICE (1.2.1) [4]; EULAR (1) [2]; OARSI (–) |
| 21) At least 10–12 sessions are needed to ensure proper treatment for osteoarthritis. | NICE (1.4.1) [4]; EULAR (6) [2]; OARSI (–) |
| 22) In the treatment for osteoarthritis, the patient's adherence to the treatment must be motivated. | NICE (1.3.2–1.4.1–1.7.1) [4]; EULAR (–); OARSI (–) |
| 23) Joint hyaluronic acid and/or corticosteroid infiltrations should be considered. | NICE (1.5.12–1.5.13) [4]; EULAR (–); OARSI (Table 2) [3] |
| 24) The supplements of chondroitin and glucosamine should be considered (reversed statement). | NICE (1.4.5) [4]; EULAR (–); OARSI (–) |

Legend: (n), CPGs paragraph into which the statements were originally reported; (–), the CPGs did not adopt a position on that statement; [n], CPGs reference

Briefly, they were considered as delivering the recommended intervention if they chose all the treatments recommended by the CPGs for the patient described in the vignette, without selecting non-recommended treatments. They were considered partially delivering the recommended intervention if they chose only some of the recommended treatments but none of the non-recommended ones. Lastly, they were considered as non-delivering the recommended intervention if they chose at least one of the non-recommended treatments or if they decided either not to treat the patient or to treat her for fewer than five sessions, and therefore the percentage of physiotherapists' 'Delivering', 'Partially Delivering' and 'Non-Delivering' the recommended intervention was calculated.

Results

Participants

Through the AIFI and the University of Genova newsletter, we were able to reach a total of 1582 physiotherapists, of which 1062 (response rate: 67%) completed the online survey between 16 June 2020 and 6 July 2020. Among them, 40 (4%) had not treated any patient with osteoarthritis in the previous six months, and 200 (19%) did not complete the survey in all its sections. Thus, 822 (77%; (mean age (SD):35.8 (13.3); female 47%; male 53%) physiotherapists compiled the questionnaire in all its sections (Fig. 1) and were included in the analysis (Table 3). Of these, 465 physiotherapists (57%) declared to have read at least one osteoarthritis CPG, whereas 357 (43%) did not.

Table 2 Section 2: Clinical vignette proposed treatments

| Clinical Scenario: | | |
|---|---|--|
| <p>Maria, a 72-year-old housewife, lives with her husband, who is in good health. She cultivates the hobby of gardening. For the past ten years, she has been suffering from knee pain which, in certain periods, forces her to take NSAIDs and to limit daily activities for a few days. Over the past two years, the pain has become increasingly frequent (VAS 5/10), so that she has decided to find some help with the housework and she is struggling to take care of the garden. She also suffers from diabetes and is overweight (BMI 28). She decides to consult her physician, who recommends her to do a visit to the physiotherapist.</p> | | |
| Management: | | |
| Core Treatment | Partially Core Treatment | Non-Core Treatment |
| <ul style="list-style-type: none"> Evaluation and planning of the rehabilitation treatment; Weight loss advice. | <ul style="list-style-type: none"> Referral to the physician for drug therapy. | <ul style="list-style-type: none"> Referral to the physician for arthroscopic surgery (joint debridement); Referral to the physician for prosthetic intervention. |
| Assessment: | | |
| Core Treatment | | |
| <ul style="list-style-type: none"> Assessment of the quantity and quality of pain; Assessment of the function; Assessment of disability and participation. | | |
| Treatment: | | |
| Core Treatment | Partially Core Treatment | Non-Core Treatment |
| <ul style="list-style-type: none"> Specific exercise on the joint (muscle strengthening); Generic exercise (aerobic exercise or generic physical activity); Education on the pathophysiology of osteoarthritis. | <ul style="list-style-type: none"> Manual therapy (mobilisation and/or massage); TENS; Load reduction devices (braces, insoles or walking aids); Hyaluronic acid and corticosteroid injections. | <ul style="list-style-type: none"> Activity rest (reduce the load on the joint); Other physical therapies (Laser, Ultrasound etc.); Supplement integration: glucosamine and chondroitin |
| <p>For how many sessions would you treat this patient?</p> <ul style="list-style-type: none"> For less than 5 sessions; Between 5 and 10 sessions; For more than 10 sessions. | | |

Section 1: level of knowledge of clinical practice guidelines – statement consensus

Overall, consensus was achieved for 13 (54%) statements (1–3, 5, 8, 9, 12, 14–16, 19, 20, 22) out of 24 (Fig. 2).

These statements addressed the role of clinical assessment, exercise, education, weight loss, and the effectiveness of physical therapies in the management of people with knee or hip osteoarthritis. Conversely, the consensus was not reached for those statements dealing with the role of supplements, radiographic findings, manual therapy, topical non-steroidal drugs, TENS, the number of sessions and the criteria for clinical diagnosis.

Section 2: level of adherence to clinical practice guidelines – clinical vignette

Demographic characteristics, percentages of the selection of each item and the classification of the


participants in ‘Delivering’, ‘Partially Delivering’ and ‘Non-Delivering’ the recommended intervention are reported in Table 4 and Table 5.

The ‘Delivering’ group ($N = 202$; 25%) provided the patient with all the CPGs recommended treatments. In the ‘Partially Delivering’ group ($N = 181$; 22%), all the participants performed the physiotherapy treatment, but only half delivered weight loss advice. The majority of the sample assessed functionality, disability, participation, pain and, delivered muscle-strengthening exercises and education. General exercise (e.g. aerobic exercise and general physical activity) was prescribed by about half of the group. Finally, in the ‘Non-Delivering Group’ ($N = 439$; 53%), the majority of participants performed the physiotherapy treatment. As far as non-recommended treatments are concerned, load reduction (rest), and other physical therapies (e.g. ultrasound and laser) were the most often delivered. Among the recommended treatments, the ones that were delivered by the majority of the members of the group were muscle strengthening and pathophysiology education. About half of this group prescribed generic exercises (aerobic exercise or generic physical activity).

Discussions

Clinical practice guidelines are an important tool that aims at bridging the gap between best evidence and clinical practice. Although our study highlighted an overall good level of knowledge of the core first-line intervention, adherence to CPGs was low. Many physiotherapists did not advise losing weight, but advised rest, while often including secondary treatments (e.g. manual therapy) supported by low-level evidence.

Most physiotherapists (> 90%) participating in the survey were aware of the importance of therapeutic exercise, education and enhancing patients’ adherence to the treatment, in the caretaking process of people with hip or knee osteoarthritis. These results are in line with the ones reached by other physiotherapists worldwide [16–19]. Despite this, 56% of the physiotherapists participating in the survey considered it essential to include manual therapy (e.g. manual mobilisation, massage) in the treatment. Current evidence shows that when manual therapy is compared with exercise therapy alone, it provides only short term benefits in reducing pain, improving function, and physical performance [35]. However, this conclusion on manual therapy is gathered from low-quality evidence, and therefore CPGs rated it as lower quality when compared to the evidence supporting exercise, which makes manual therapy only a conditional treatment [3, 4]. In line with this, prioritising manual therapy in patients’ management may reduce the time allocated to exercise. However, we did not ask the participants whether they considered manual therapy more



effective than exercise, or which treatment they would prioritise, thus leaving uncertainties regarding the clinical impact of this finding. Future studies, with a mixed-method design are needed to better understand how different treatments are weighed by clinicians in the management of people with osteoarthritis.

Furthermore, we found an insufficient level of knowledge in three distinct areas: a) the criteria for the clinical diagnosis of osteoarthritis, b) the role of other non-surgical interventions that could enhance therapeutic exercise benefits (e.g. topical anti-inflammatory drugs and TENS), c) the number of sessions needed to ensure an optimal outcome. As far as the clinical diagnosis and


drug prescription are concerned, this lack of knowledge might be due to the fact that, in Italy, physiotherapists are not allowed to perform clinical diagnosis and prescribe drugs. However, they are often the first health care professionals that people with osteoarthritis refer to. Thus, they should be aware of the recommended pharmacological management to facilitate the integration of drug therapy with physiotherapy and the proper clinical diagnostic criteria in order to refer patients to relevant healthcare professionals, when necessary.

Our study showed that only 10% of the respondents considered being 45 years old or older, having pain, and joint stiffness for less than 30 min in the morning,

Table 3 Participants' demographic characteristics

| Demographic Characteristics | |
|--|--------------------|
| Age (years)(mean,(SD)) | 35.77 (13.3) |
| Sex (female); (male) (N (%)): | 387 (47); 435 (53) |
| Years of Practice (N (%)): | |
| Less than 1 year | 87 (11) |
| From 1 to 5 years | 319 (39) |
| From 6 to 10 years | 149 (18) |
| More than 10 years | 267 (32) |
| Highest Academic Level Reached (N (%)): | |
| Bachelor of Science (BSc)/Equivalent title | 282 (34) |
| Post-Graduate I Level Degree* | 382 (47) |
| Master of Science (MSc)/Post-Graduate II Level Degree† | 122 (15) |
| Doctor of Philosophy (PhD) | 36 (4) |
| Read at least one osteoarthritis CPGs (N (%)): | |
| Yes | 465 (57) |
| No | 357 (43) |

Legend: N, number; %, percentage; *Academic degree that can be gained after BSc (Italian education system); † Academic degree that can be gained after MSc (Italian education system)



sufficient criteria for the diagnosis of osteoarthritis, as recommended by the NICE CPGs. These seemed to be the most appropriate criteria to ensure that even younger patients with osteoarthritis would receive appropriate care in line with CPGs [36]. The lack of agreement regarding clinical diagnostic criteria may partially explain the relatively high percentage (> 40%) of physiotherapists who considered radiographic findings necessary to

express a clinical diagnosis of osteoarthritis. This data is consistent with the findings of a similar survey conducted by Ayanniyi et al. [19]. Radiographic findings should be taken into account only when other diseases are the suspected cause of the symptoms (e.g. infection, cancer, rheumatoid arthritis) or when surgical intervention is planned [4]. These recommendations are based on evidence that shows a weak association between the

Table 4 Participants’ profile by level of adherence for the Clinical Vignette

| | ‘Delivering’ (N = 202) | ‘Partially Delivering’ (N = 181) | ‘Non-Delivering’ (N = 439) |
|---|------------------------|----------------------------------|----------------------------|
| Age (years)(mean,(SD)) | 31.2 (10.9) | 37.4 (13.7) | 37.2 (13.7) |
| Sex (N (%)): | | | |
| Female | 81 (40) | 86 (48) | 220 (50) |
| Male | 121 (60) | 95 (52) | 219 (50) |
| Years of Practice (N (%)): | | | |
| Less than 1 year | 30 (15) | 18 (10) | 39 (9) |
| From 1 to 5 years | 100 (50) | 71 (39) | 148 (34) |
| From 6 to 10 years | 40 (20) | 28 (16) | 81 (18) |
| More than 10 years | 32 (15) | 64 (35) | 171 (39) |
| Highest Academic Level Reached (N (%)): | | | |
| Bachelor of Science (BSc)/Equivalent level | 57 (28) | 64 (35) | 161 (37) |
| I Level Master Degree* | 115 (57) | 79 (44) | 188 (43) |
| Master of Science (MSc)/II Level Mastert | 23 (11) | 32 (18) | 67 (15) |
| Doctor of Philosophy (PhD) | 7 (4) | 6 (3) | 23 (5) |
| Read at least one osteoarthritis CPGs (N (%)): | | | |
| Yes | 124 (61) | 114 (63) | 240 (55) |
| No | 78 (39) | 67 (37) | 199 (45) |

Legend: N, number; %, percentage; *Academic degree that can be gained after BSc (Italian education system); † Academic degree that can be gained after MSc (Italian education system)

Table 5 Frequencies of answers to Clinical Vignette by level of adherence with Clinical Practice Guidelines

| Question | All (N = 822; 100%) | | 'Delivering' (N = 202; 25%) | | 'Partial Delivering' (N = 181; 22%) | | 'Non-Delivering' (N = 439; 53%) | |
|--|---------------------|----------|-----------------------------|-----------|-------------------------------------|-----------|---------------------------------|----------|
| | YES | NO | YES | NO | YES | NO | YES | NO |
| Section 1: Management (N (%)) | | | | | | | | |
| 1) Physiotherapy treatment | 794 (97) | 28 (3) | 202 (100) | 0 (0) | 181 (100) | 0 (0) | 411 (94) | 28 (6) |
| 2) Referral pharmacological | 159 (19) | 663 (81) | 40 (20) | 162 (80) | 15 (8) | 166 (92) | 104 (24) | 335 (76) |
| 3) Referral for debridement | 15 (2) | 807 (98) | 0 (0) | 202 (100) | 0 (0) | 181 (100) | 15 (3) | 424 (97) |
| 4) Referral for surgery | 26 (3) | 796 (97) | 0 (0) | 202 (100) | 0 (0) | 181 (100) | 26 (6) | 413 (94) |
| 5) Weight Loss | 645 (79) | 177 (21) | 202 (100) | 0 (0) | 90 (50) | 91 (50) | 353 (80) | 86 (20) |
| Section 2: Assessment (N (%)) | | | | | | | | |
| 1) Pain | 727 (88) | 95 (12) | 202 (100) | 0 (0) | 138 (76) | 43 (24) | 387 (88) | 52 (12) |
| 2) Functionality | 756 (92) | 66 (8) | 202 (100) | 0 (0) | 164 (91) | 17 (9) | 390 (89) | 49 (11) |
| 3) Disability and Participation | 722 (88) | 100 (12) | 202 (100) | 0 (0) | 149 (82) | 32 (18) | 371 (85) | 68 (15) |
| Section 3: Treatment (N (%)) | | | | | | | | |
| 1) Load reduction (rest) | 244 (30) | 578 (70) | 0 (0) | 202 (100) | 0 (0) | 181 (100) | 244 (56) | 195 (44) |
| 2) Manual therapy | 612 (75) | 210 (25) | 145 (72) | 57 (28) | 135 (75) | 46 (25) | 332 (76) | 107 (24) |
| 3) Muscles strengthening | 702 (85) | 120 (15) | 202 (100) | 0 (0) | 143 (79) | 38 (21) | 357 (81) | 82 (19) |
| 4) Generic exercise (e.g. aerobic exercise or generic physical activity) | 525 (64) | 297 (36) | 202 (100) | 0 (0) | 92 (51) | 89 (49) | 231 (53) | 208 (47) |
| 5) Education on the pathophysiology of osteoarthritis | 698 (85) | 124 (15) | 202 (100) | 0 (0) | 142 (78) | 39 (21) | 354 (81) | 85 (19) |
| 6) TENS | 111 (14) | 711 (86) | 25 (12) | 177 (88) | 27 (15) | 154 (85) | 59 (13) | 380 (87) |
| 7) Other Physical Therapies (e.g. ultrasound and laser) | 137 (17) | 685 (83) | 0 (0) | 202 (100) | 0 (0) | 181 (100) | 137 (31) | 302 (69) |
| 8) Load reduction devices (e.g. braces, insoles or walking aids). | 185 (23) | 637 (77) | 30 (15) | 172 (85) | 34 (19) | 147 (82) | 121 (28) | 318 (72) |
| 9) Hyaluronic acid and corticosteroids | 111 (14) | 711 (86) | 7 (4) | 195 (96) | 19 (11) | 162 (89) | 85 (19) | 354 (81) |
| 10) Treatment: Supplements | 279 (34) | 543 (66) | 0 (0) | 202 (100) | 0 (0) | 181 (100) | 77 (18) | 362 (82) |
| Section 4: Number of sessions (N (%))* | | | | | | | | |
| Less than 5 sessions | 71 (9) | 0 (0) | 0 (0) | 71 (17) | Less than 5 sessions | 71 (9) | 0 (0) | 0 (0) |
| Between 5 and 10 sessions | 465 (58) | 138 (68) | 121 (67) | 206 (50) | Between 5 and 10 sessions | 465 (58) | 138 (68) | 121 (67) |
| More than 10 sessions | 258 (33) | 64 (32) | 60 (33) | 134 (33) | More than 10 sessions | 258 (33) | 64 (32) | 60 (33) |

Legend: N, number; %, percentage; *Percentage calculated on N = 794; N = 28 could not access to this section as they didn't check the "Management section: 1) Physiotherapy treatment" option

severity of radiographic findings and pain and disability levels [37, 38]. Furthermore, basing clinical decisions on imaging fosters the perception of osteoarthritis as a wear-and-tear disease which may, in turn, induce fear-avoidance behaviours [38, 39]. However, since physiotherapists in Italy are unable to prescribe radiographic investigations, the impact of this finding on the clinical management of the patients is uncertain.

The second section of the survey showed mismatching between the knowledge of CPGs and their application in clinical practice. Although most physiotherapists showed adequate knowledge of first-line interventions, only a minority (25%) wholly adhered to CPGs, whereas more than 50% included at least one non-recommended strategy or treatment.

One of the challenges in the implementation of CPGs seems to be the fact that health care professionals view osteoarthritis as a “non-serious” disease. This may depend on inadequate preparation at undergraduate level [11, 20, 40]. These erroneous beliefs might be carried on in post-graduate degrees, since the percentage of “Non-Delivering” did not change throughout the different levels of academic degrees achieved (> 50%). Moreover, in Italy, there are no MSc degrees available, specifically on musculoskeletal and rheumatic conditions, and PhD curricula are very specific, therefore, if not focussed on osteoarthritis, they cannot bridge the pre-existing evidence-to-practice gap. In addition, implementing osteoarthritis CPGs in the complex setting of clinical care can be challenging [20] since clinicians have to face several barriers among which patients’ preferences, resource availability, discrepancies between CPGs, lack of English knowledge and limited access to information [22].

Although the patient in the clinical vignette was overweight and presented with moderate symptoms, the interviewed physiotherapists often excluded advising weight loss, whilst rest was often considered. In fact, recommending weight loss may be considered by some physiotherapists as beyond their clinical scope [41]. This data is in line with other studies showing that both Australian and British physiotherapists recommended muscle strengthening exercises, but seemed less confident in prescribing aerobic exercise and recommending weight loss [18, 41]. Providing physiotherapists with additional specific training aimed at dealing with overweight patients may enhance the patients’ outcomes and increase the overall level of adherence to CPGs.

The interpretation of the relatively high inclusion of rest as well as the load reduction in the treatment is difficult to explain, especially in light of the good level of knowledge shown in the first part of the questionnaire. However, the CPGs available do not specify how to adapt the therapeutic exercise in those cases with severe

osteoarthritis symptoms, where pain can be easily triggered by joint movement or weight-bearing activities. This is also highlighted by the fact that about 50% of the physiotherapists in the “Non-Delivering” group declared to have read at least one osteoarthritis CPG. Thus, in light of these results, it can be hypothesised that physiotherapists may feel unsure and unprepared when having to deal with this pain condition [42]. Discrepancies between CPGs knowledge and application may also depend on factors that are external to the physiotherapist, and they may vary by country. Regarding osteoarthritis CPGs, this is the first study that pointed towards this discrepancy, starting from several CPGs and by considering a plethora of treatments.

Beneath the differences between the three groups, a transversal trait was found regarding the application of manual therapy which was delivered by more than 70% of the sample. From a cultural perspective, manual therapy is a core competence of physiotherapy, which set the basis of this professional figure in the past, and patients often expect this type of treatments from physiotherapists [40]. Meeting patients’ expectations is thought to foster a positive clinician-patient relationship while enhancing the treatment outcome by inducing analgesia, regulating patients’ emotions, and reorganising the body’s mental representations [43, 44]. Therefore, this data may reflect the contrast between treatments recommendations and patients’ expectations, which can be itself the results of a specific cultural belief that needs to be investigated.

Some limitations of this study need to be discussed. Firstly, our sample was mainly based on physiotherapists who completed a post-graduate degree, therefore our results might overestimate the real level of knowledge of and adherence to osteoarthritis CPGs. Secondly, we did not investigate the participants’ clinical practice setting (e.g. private practice, public care etc.) which might have had an impact on the participants’ level of adherence to CPGs.

Our findings revealed that Italian physiotherapists are aware of the core treatments for patients with osteoarthritis. However, they showed a low level of knowledge of the clinical diagnostic criteria and of the usefulness of other non-surgical treatments that can support first-line intervention (e.g. TENS and non-steroidal anti-inflammatory medications). Moreover, an adequate level of adherence is yet to be reached. These results identify an evidence-to-practice gap which may lead to non evidence-based practice behaviours for the management of the patients with hip and knee osteoarthritis.

Finding new strategies to bridge the gap between evidence and clinical practice appears to be necessary, therefore providing physiotherapists with CPGs in their native language and fostering their use through

university programmes could be one of the possible solutions proposed. Moreover, the use of recognised manuals aimed at developing CPGs is advocated. These should ascertain that all search stages are documented for transparency and reproducibility and that the most important elements for a real practical implementation, such as algorithms for clinical decision-making for complicated cases, and patients' inclusion-exclusion criteria, are included [45].

Finally, the professional image of physiotherapists within society should be reconceptualised. In particular, we should continue to foster a new vision of physiotherapists, as no longer anchored to treatments that are mainly based on physical and manual therapy, but as figures whose treatment paradigm focusses on improving the patient's individual functioning by specific treatment strategies, such as exercise and education, that take into account scientific evidence conveyed into specific contexts.

Acknowledgements

The researchers would like to thank all the participants that joined the study, the Italian Physiotherapist Association (AIFI) and the people that helped with the participants recruitment. This work was developed within the DINOEMI Department of Excellence framework of MIUR 2018-2022 (Legge 232 del 2016).

Authors' contributions

SB, SS, SM, MT and AD made substantial contributions to the conception and design, data acquisition, or analysis and interpretation of data. SB, MT and AD participated in drafting the article or revising it critically for important intellectual content. SS, SM, MT and AD gave final approval of the version to be published. All authors agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The author(s) read and approved the final manuscript.

Funding

This research did not receive external funding. Open Access funding provided by Lund University.

Availability of data and materials

The datasets used and analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Ethical approval was obtained from the Ethics Committee for University Research (CERA: Comitato Etico per la Ricerca di Ateneo), University of Genova (approval date: 15/06/2020; CERA2020.07). The participants signed digital informed consent before participation.

Consent for publication

The participants signed digital consent for publication by inclusion in the study.

Competing interests

The authors report no conflicts of interest.

Author details

¹Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genova, Campus of Savona, Genova, Italy. ²Clinical Epidemiology Unit, Orthopedics, Department of Clinical Sciences Lund, Lund University, Lund, Sweden.

Received: 24 January 2021 Accepted: 8 April 2021

Published online: 23 April 2021

References

- James SL, Abate D, Abate KH, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the global burden of disease study 2017. *Lancet*. 2018;392:1789-858.
- Fernandes L, Hagen KB, Bijlsma JWJ, et al. EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. *Ann Rheum Dis*. 2013;72:1125-35.
- Bannuru RR, Osani MC, Vaysbrot EE, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthr Cartil*. 2019;27:1578-89.
- National Clinical Guidelines Centre. Osteoarthritis: Care and Management in Adults. NICE Clin Guidel. 2014. p. 1-498.
- Institute of Medicine (IOM). Clinical practice guidelines we can trust. *Clin Pract Guidel We Can Trust*. 2011; <https://doi.org/10.17226/13058>.
- Swärdh E, Opava CH, Nygård L, Lindquist I. Understanding exercise promotion in rheumatic diseases: a qualitative study among physical therapists. *Physiother Theory Pract*. 2019;30:1-10.
- Juhl C, Christensen R, Roos EM, Zhang W, Lund H. Impact of exercise type and dose on pain and disability in knee osteoarthritis: a systematic review and meta-regression analysis of randomized controlled trials. *Arthritis Rheumatol*. 2014;66:622-36.
- Thorstensson CA, Garellick G, Rystedt H, Dahlberg LE. Better Management of Patients with osteoarthritis: development and Nationwide implementation of an evidence-based supported osteoarthritis self-management Programme. *Musculoskeletal Care*. 2015;13:67-75.
- Dougados M, Betteridge N, Burmester GR, et al. EULAR standardised operating procedures for the elaboration, evaluation, dissemination, and implementation of recommendations endorsed by the EULAR standing committees. *Ann Rheum Dis*. 2004;63:1172-6.
- Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924-6.
- Egerton T, Diamond LE, Buchbinder R, Bennell KL, Slade SC. A systematic review and evidence synthesis of qualitative studies to identify primary care clinicians' barriers and enablers to the management of osteoarthritis. *Osteoarthr Cartil*. 2017;25:625-38.
- Zadro J, O'Keefe M, Maher C. Do physical therapists follow evidence-based guidelines when managing musculoskeletal conditions? Systematic review. *BMJ Open*. 2019;9:e032329.
- Basedow M, Esterman A. Assessing appropriateness of osteoarthritis care using quality indicators: a systematic review. *J Eval Clin Pract*. 2015;21:782-9.
- Theis KA, Brady TJ, Sacks JJ. Where have all the patients gone? Profile of US adults who report doctor-diagnosed arthritis but are not being treated. *JCR J Clin Rheumatol*. 2018;25:341-7.
- Holden MA, Nicholls EE, Hay EM, Foster NE. Physical therapists' use of therapeutic exercise for patients with clinical knee osteoarthritis in the United Kingdom: in line with current recommendations? *Phys Ther*. 2008;88:1109-21.
- Holden MA, Nicholls EE, Young J, Hay EM, Foster NE. UK-based physical therapists' attitudes and beliefs regarding exercise and knee osteoarthritis: Findings from a mixed-methods study. *Arthritis Rheum*. 2009;61:1511-21.
- Tittlemier BJ, Wittmeier KD, Robinson DB, Webber SC. Knee Osteoarthritis: An Investigation into the Clinical Practice of Physiotherapists in Canada. *Physiother Canada*. 2020;73:e20190068.
- Tang CY, Pile R, Croft A, Watson NJ. Exploring physical therapist adherence to clinical guidelines when treating patients with knee osteoarthritis in Australia: a mixed methods study. *Phys Ther*. 2020;100:1084-93.
- Ayanniyi O, Egwu RF, Adeniyi AF. Physiotherapy management of knee osteoarthritis in Nigeria—a survey of self-reported treatment preferences. *Hong Kong Physiother J*. 2017;36:1-9.
- Swaithes L, Paskins Z, Dziedzic K, Finney A. Factors influencing the implementation of evidence-based guidelines for osteoarthritis in primary care: a systematic review and thematic synthesis. *Musculoskeletal Care*. 2020; 18:101-10.
- Dziedzic KS, French S, Davis AM, Geelhoed E, Porcheret M. Implementation of musculoskeletal models of care in primary care settings: theory, practice,

- evaluation and outcomes for musculoskeletal health in high-income economies. *Best Pract Res Clin Rheumatol*. 2016;30:375–97.
22. Fischer F, Lange K, Klose K, Greiner W, Kraemer A. Barriers and strategies in guideline implementation—a scoping review. *Healthcare*. 2016;4:36.
 23. Gupta S, Rai N, Bhattacharyya O, Cheng AYY, Connelly KA, Boulet LP, et al. Optimizing the language and format of guidelines to improve guideline uptake. *CMAJ*. 2016;188:E362–8.
 24. Lilienfeld SO, Ritschel LA, Lynn SJ, Brown AP, Cautin RL, Latzman RD. The research-practice gap: bridging the schism between eating disorder researchers and practitioners. *Int J Eat Disord*. 2013;46:386–94.
 25. Kristensen N, Nymann C, Konradsen H. Implementing research results in clinical practice—the experiences of healthcare professionals. *BMC Health Serv Res*. 2015;16:48.
 26. Vliet Vlieland TPM, Van Den Ende CHM, Alliot-Launois F, et al. Educational needs of health professionals working in rheumatology in Europe. *RMD Open*. 2016;2:e000337.
 27. de Leeuw D, Hox JDD. *International handbook of survey methodology* (European Association of Methodology Series). 1st ed. New York: Taylor and Francis group; 2008.
 28. Eysenbach G. Improving the quality of web surveys: the checklist for reporting results of internet E-surveys (CHERRIES) | Eysenbach | journal of medical internet research. *J Med Internet Res*. 2004;6:e34+.
 29. Von Elm E, Altman D, Egger M, Pocock S, Gøtzsche P, Vandenbroucke J, et al. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Ann Intern Med*. 2007;147:573–8.
 30. The European Parliament and the Council of the European Union (2016). Regulation (EU) 2016/679 of the European Parliament and of the Council of 27 April 2016 on the protection of natural persons with regard to the processing of personal data and on the free movement of such data, and repealing Directive 95/46/EC. 1–88.
 31. Joshi A, Kale S, Chandel S, Pal DK. Likert scale: explored and explained. *Br J Appl Sci Technol*. 2015;7:396–403.
 32. Suárez-Alvarez J, Pedrosa I, Lozano LM, García-Cueto E, Cuesta M, Muñoz J. Using reversed items in likert scales: a questionable practice. *Psicothema*. 2018;30:149–58.
 33. Rossetini G, Palese A, Geri T, Fiorio M, Colloca L, Testa M. Physical therapists' perspectives on using contextual factors in clinical practice: findings from an Italian national survey. *PLoS One*. 2018;13:e0208159.
 34. Teo PL, Hinman RS, Egerton T, Dziedzic KS, Bennell KL. Identifying and prioritizing clinical guideline recommendations Most relevant to physical therapy practice for hip and/or knee osteoarthritis. *J Orthop Sport Phys Ther*. 2019;49:501–12.
 35. Anwer S, Alghadir A, Zafar H, Brismée JM. Effects of orthopaedic manual therapy in knee osteoarthritis: a systematic review and meta-analysis. *Physiother (United Kingdom)*. 2018;104:264–76.
 36. Skou ST, Koes BW, Grønne DT, Young J, Roos EM. Comparison of three sets of clinical classification criteria for knee osteoarthritis: a cross-sectional study of 13,459 patients treated in primary care. *Osteoarthr Cartil*. 2020;28:167–72.
 37. Helminen E-E, Sinikallio SH, Valjakka AL, Väisänen-Rouvali RH, Arokoski JP. Determinants of pain and functioning in knee osteoarthritis: a one-year prospective study. *Clin Rehabil*. 2016;30:890–900.
 38. Darlow B, Brown M, Thompson B, Hudson B, Grainger R, McKinlay E, et al. Living with osteoarthritis is a balancing act: an exploration of patients' beliefs about knee pain. *BMC Rheumatol*. 2018;2:15.
 39. Gunn AH, Schwartz TA, Arbeeva LS, et al. Fear of movement and associated factors among adults with symptomatic knee osteoarthritis. *Arthritis Care Res*. 2017;69:1826–33.
 40. Briggs AM, Houlding E, Hinman RS, et al. Health professionals and students encounter multi-level barriers to implementing high-value osteoarthritis care: a multi-national study. *Osteoarthr Cartil*. 2019;27:788–804.
 41. Holden MA, Waterfield J, Whittle R, Bennell K, Quicke JG, Chesterton L, et al. How do UK physiotherapists address weight loss among individuals with hip osteoarthritis? A mixed-methods study. *Musculoskeletal Care*. 2019;17:133–44.
 42. Morin Chabane S, Coutinho F, Laliberte M, Feldman D. Outpatient physiotherapists' attitudes and beliefs toward patients with chronic pain: a qualitative study. *Physiother Theory Pract*. 2020;36:85–94.
 43. Rossetini G, Carlino E, Testa M. Clinical relevance of contextual factors as triggers of placebo and nocebo effects in musculoskeletal pain. *BMC Musculoskelet Disord*. 2018;19:27.
 44. Geri T, Viceconti A, Minacci M, Testa M, Rossetini G. Manual therapy: exploiting the role of human touch. *Musculoskelet Sci Pract*. 2019;44:102044.
 45. Rosenfeld RM, Shiffman RN. Clinical practice guideline development manual: a quality-driven approach for translating evidence into action. *Otolaryngol - Head Neck Surg*. 2009;140:1–43.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions



Study 2

Giving an account of patients' experience: A qualitative study on the care process of hip and knee osteoarthritis

Simone Battista MSc, PhD Student^{1,2}  | Mattia Manoni MSc, PhD Student¹ |
Andrea Dell'Isola PhD, Post-Doc Research Fellow² | Martin Englund PhD, Professor² |
Alvisa Palese PhD, Professor³ | Marco Testa PhD, Professor¹

¹Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genova, Savona, Italy

²Department of Clinical Sciences, Clinical Epidemiology Unit, Orthopedics, Lund University, Lund, Sweden

³Department of Medical Sciences, School of Nursing, University of Udine, Udine, Italy

Correspondence

Andrea Dell'Isola, PhD, Post-Doc Research Fellow, Department of Clinical Sciences, Clinical Epidemiology Unit, Orthopedics, Lund University, Wigerthuset, Remissgatan 4, Skåne University Hospital, SE-221 85 Lund, Sweden.
Email: andrea.dellisola@med.lu.se

Funding information

European Alliance of Associations for Rheumatology (EULAR), Grant/Award Number: The EULAR Health Professionals research grant 2020

Abstract

Introduction: Despite the publication of clinical practice guidelines, the quality of the care process as experienced by patients with osteoarthritis (OA) appears suboptimal. Hence, this study investigates how patients with OA experience their disease and care process, highlighting potential elements that can enhance or spoil it, to optimise their quality of care.

Methods: A qualitative study based on semi-structured interviews. Patients with hip and knee OA in Italy were interviewed. The interview guide was created by a pool of health professionals and patients. The interviews were analysed through a theme-based analysis following a philosophy of descriptive phenomenological research.

Results: Our analysis revealed seven main themes: (1) *Experiencing a sense of uncertainty*, as interviewees perceived treatment choices not to be based on medical evidence; (2) *Establishing challenging relationships with the self and the other*, as they did not feel understood and felt ashamed or hopeless about their condition; (3) *Being stuck in one's own or the health professionals' beliefs about the disease management*, as a common thought was the perception of movement as something dangerous together with a frequent prescription of passive therapies; (4) *Dealing with one's own attitudes towards the disease*; (5) *the barriers to and* (6) *the facilitators of the adherence to therapeutic exercise*, which revolve around the therapy cost, the time needed and the patients' willingness to change their life habits and (7) *Developing an uneasy relationship with food* since the diet was considered as something that "you force yourself to follow" and overeating as a way "to eat your feelings".

Conclusion: The lack of clear explanations and a negative attitude towards first-line nonsurgical treatments (mainly physical exercise), which are considered as a way to fill the time while waiting for surgery, underlines the importance of providing patients with adequate information about OA treatments and to better explain the

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Health Expectations* published by John Wiley & Sons Ltd.

role of first-line intervention in the care of OA. This will enhance patient-centred and shared decision-making treatments.

Patient Contribution: Patients with hip and knee OA participated in creating the interview and contributed with their experience of their care process.

KEYWORDS

clinical governance, osteoarthritis, phenomenology, physical therapists, physical therapy speciality, policy making, practice guidelines as topic

1 | INTRODUCTION

The care process is often a complex and intimate experience as lived by patients. Caring is one of the expressions of the ontology of human beings since people are dependent on one another.^{1,2} Due to its grounded origin in anthropology, the care process can be considered as a 'ritual' performed around the patient.³ It is a complex psychosocial context with a nonspecific effect on patients' brain that can amplify or reduce the specific effect of the treatment, tapping into people's beliefs, expectations and feelings.³⁻⁵ An example of this complexity can be found in the care of individuals with hip and knee osteoarthritis (OA).

OA is the most common form of arthritis and one of the foremost causes of compromised health-related quality of life worldwide.^{6,7} Its first-line intervention includes therapeutic exercises and education programmes, often delivered by physiotherapists, and diet when needed.⁸⁻¹¹ Therefore, this intervention is grounded on highly demanding nonsurgical treatments in terms of patients' compliance and on a cultural change at the individual and societal levels that need to accept and foster the importance of these interventions.^{12,13}

To answer this need, several national and international clinical practice guidelines (CPGs) were published all over the world.⁸⁻¹¹ However, different international studies brought to the forefront that only one-third of patients are receiving the recommended first-line interventions.¹⁴⁻¹⁶ Basedow and Esterman¹⁴ analysed the appropriateness of OA care through the synthesis of quality indicators retrieved from global medical records, administrative databases and patient questionnaires and interviews. These indicators are generally expressed as pass-rates, and they concluded that the quality of OA care was suboptimal for all treatment domains, with first-line interventions reaching the lowest pass-rate.¹⁴

The reasons behind the failure of the implementation of first-line interventions are several. They include, among others, patients' preferences, beliefs and experiences about their disease and the care process behind it.⁴ In the healthcare setting of musculoskeletal conditions, such as OA, efforts on assessing and understanding patients' experience could help health policy-makers foster improvements in healthcare providers' interpersonal aspects and patients' expectations on how healthcare should be delivered.¹⁷

Therefore, it is essential to investigate the experience of people with OA about their disease, focussing on the care process they

received, exploring their experiences, preferences and beliefs, so as to highlight potential elements that affect it, both positively or negatively. For what concerns the Italian healthcare system, it provides its citizens with universal coverage essentially free of charge.¹⁸ However, Italy lacks national CPGs for OA management that are officially recognised by the Italian Higher Institute of Health. This results in the absence of a standardised care process for this disease. Therefore, health professionals can only rely on international CPGs, which we showed elsewhere to be scarcely followed by Italian physiotherapists.¹⁹ Hence, this qualitative study explored the experience of people with OA about the care process they received in Italy.

2 | METHODS

2.1 | Study design

A descriptive phenomenological study was performed from October 2020 to March 2021. The descriptive phenomenological inquiry aims at ensuring "direct explorations, analysis and descriptions of particular phenomena (as in this case—the care process), as free as possible from unexamined presuppositions, maximising intuitive presentation".²⁰⁻²⁴ The underpinning intent of the phenomenological researcher is to give voice and power to individuals who experienced, or are experiencing, at first-hand the phenomenon of interest as, in this instance, people with OA.²²

The research was conducted in respect of the Declaration of Helsinki and reported following the Consolidated Criteria for Reporting Qualitative Research (COREQ; Table S1).²⁵ Ethical approval was obtained from the Ethics Committee for University Research (CERA: Comitato Etico per la Ricerca di Ateneo), University of Genova (Approval date: 15 June 2020; CERA2020.07).

2.2 | Participants

A purposeful sampling method was adopted to ensure the maximum variations of the experiences.²⁶ Participants living in different geographic locations, both urban (core areas of cities) and suburban (residential areas that surround main cities), with hip OA, knee OA or

hip and knee OA, were considered eligible according to the need to reach a broad understanding of the studied phenomenon, which stems from different experiences.²⁶ Specifically, individuals with physician-diagnosed knee and hip OA, able to speak Italian and willing to participate were considered eligible to join this study. Those patients reporting joints other than hip or knee as the primary joints for OA symptoms were not considered eligible.

A network, including physicians (i.e., orthopaedics, rheumatologists and general practitioners) and other health professionals (i.e., physiotherapists and nurses), specialised in the rehabilitation of rheumatic and musculoskeletal diseases was created to help with the participants' recruitment. Health professionals were first approached individually by the research team and informed about the study aims and procedures of data collection. After obtaining their collaboration, the eligible participants were contacted by the health professionals in the network and informed on the aim of the study, the interview process (i.e., places and proposed dates) and the data confidentiality and anonymity. Eligible participants were left free to join the research and to withdraw from it at any time. Only those who expressed their interest in partaking in the study were contacted by S. B. to collect the informed consent and arrange the interview. Then, a snowball sampling was also adopted to access individuals in the network of participants with OA who had been contacted and agreed to participate first.²⁶ The recruitment was concluded once the data saturation was reached, as judged by the two authors (S. B. and M. M.) who analysed interviews. An inductive thematic saturation was followed to assess the data saturation: S. B. and M. M. kept interviewing and analysing the interviews simultaneously until no new themes were found.²⁴

2.3 | Data collection method

A semi-structured interview was designed based on the existing literature. An interview guide (Table 1) was specifically developed for this study by a pool of physiotherapists, psychologists, nurses and people with OA.^{8-11,13,19,27-30} The interview guide consisted of open questions exploring different topics related to the OA care process: (a) experiential and emotional dimensions; (b) expectations and (c) beliefs. Follow-up questions were frequently asked to investigate participants' experiences further. Examples of these questions were, 'Can you give me an example?' and 'Can you explain to me what you mean with this sentence?'. At the beginning of each interview, the participants filled in the informed consent and provided their demographic (i.e., age, gender, nationality, retirement, area of living) and clinical information (i.e., height and weight to calculate body mass index, joint(s) with OA and years living with the pathology) which were registered on an electronic sheet. Only the interviewer and the interviewee were present during the interview process. No follow-up interviews were performed.

The interviews were performed by S. B. and lasted approximately one hour each. S. B. is a physiotherapist and a PhD candidate trained in advanced qualitative methodologies, with proficiency in conducting

qualitative studies. S. B. recognises himself as male. Participants were not aware of his professional background, and none of them had close relationships with him. The interviews were performed online, by videoconferencing, and they were conducted only with the interviewee. An audio-visual recording of each interview was produced and transcribed *verbatim* by two authors (S. B. and M. M.).

2.4 | Data analysis

As far as the analysis was concerned, a theme-based analysis was performed.³¹ Thematic analysis is an independent qualitative descriptive approach described as 'a method for identifying, analysing and reporting patterns (themes) within data'.³² Since this study explored the experience of the care process, the thematic analysis was conducted within the framework of a descriptive phenomenology study, as reported and explained above.

Two authors (S. B. and M. M.) read the transcribed interviews several times to obtain a global impression of the content. S. B. and M. M. are PhD candidates (S. B. is a physiotherapist, M. M. is a psychologist), trained in qualitative methods, who both identify themselves as male. They both analysed first independently, and then jointly, the interview transcripts. Specifically, initial coding involved examining the data line by line to search for subthemes, themes, concepts and patterns. Meaning units (i.e., words, phrases and sentences that described the phenomenon of interest) were identified and framed into codes, representing significant and central aspects of the reported statements.³³ Throughout this process, emerging codes were compared to previous codes to understand the experiences of the OA care process as lived by the participants and to generate focussed codes. Finally, the focussed codes and coding were merged and synthesised to extract final subthemes and themes. For each theme, exemplary quotes were identified and reported anonymously. The themes were derived from the data and not determined in advance.

2.5 | Rigour and trustworthiness

To ensure the study rigour and trustworthiness, multiple strategies were promoted. Firstly, S. B. documented field notes ('Memos') after completing each interview to promote reflexivity.³⁴ These memos were shared during research meetings for reflexive thoughts. Secondly, the research team met frequently to refine the themes and subthemes until a consensus on the final themes was achieved. Thirdly, an audit trail containing meeting notes, analysis discussions and research decisions was continuously reorganised by the two authors who analysed the interviews (S. B. and M. M.) to stress the dependability and confirmability of the study.³⁴ An example of it is reported in Table 2. Lastly, a Synthesised Member Checking was exploited to improve the credibility of the analysis.³⁵ At the end of each interview, participants were asked if they wanted to participate in the member checking phase. All of them agreed to partake in it.

TABLE 1 Interview guide and domains investigated

| Questions | Domains |
|---|--|
| 1) I would like to start this interview by asking you how you realised that you have OA? | Experiential and emotional experiences |
| 2) How did you manage your disease? | Experiential and emotional experiences |
| 3) What prompted you to go to a physician/health professional? | Experiential and emotional experiences |
| 4) What did you expect from your physician/health professional the first time you saw them for OA? | Expectations |
| 5) How were you diagnosed? | Experiential and emotional experiences |
| 6) Would you like to tell me how you believed your disease would evolve in the future? | Beliefs |
| 7) Did you have any family or friends that supported you during your care process? | Experiential and emotional experiences |
| 8) Would like to tell how you felt when you received your diagnosis? | Experiential and emotional experiences |
| 9) How would you describe the impact of OA in your life/work? | Experiential and emotional experiences |
| 10) Would you like to tell me which treatments you expected to be suggested to manage your disease? | Expectations |
| 11) Eventually, which treatments were in fact suggested? | Experiential and emotional experiences |
| 12) Would you like to describe which treatments you deem useful in the management of OA? | Beliefs |
| 12) ...For example, physical activity? Manual therapy? | |
| 13) What did you expect from the treatments that you have received so far? | Expectations |
| 14) Would you like to tell me which roles physical activity and diet play in the management of OA? | Beliefs |
| 15) What does a healthy diet mean for you? | Beliefs |
| 16) Would you like to tell me the reasons why a person with OA may not be willing to change their lifestyle, integrating physical activity and a healthy diet into their daily routine? | Beliefs |
| 17) Would you like to tell me the role of the physiotherapist in the management of OA? ... And what would you expect from this professional figure? | Beliefs |
| 18) Which attitudes did you expect from the health professionals you met during your care process towards OA? And which one(s) did they adopt? | Expectations |
| 19) If you've ever been shown, would you like to tell me how you felt when you saw your radiographic findings? | Experiential and emotional experiences |
| 20) In your opinion, how important were radiographic findings in your OA care process? ... How important were they for the health professionals you met? | Beliefs/experiential and emotional experiences |
| 21) Would you like to tell me how you live with OA now? | Experiential and emotional experiences |
| 22) Is there anything else you would like to add? | Closing question |

Abbreviation: OA, osteoarthritis.

The member checking phase was yielded at the end of the interview and analysis process. The participants were provided with a one-page summary, highlighting the main themes and subthemes identified in the study, together with a brief plain explanation of the key findings. They were then asked to read it thoroughly, feeding back the researchers with any doubts or concerns they might have had about this summary. All the participants agreed with what was retrieved, and no further modifications to the results were done.

2.6 | Patient and public involvement

Two people with both knee and hip OA (one identified herself as female whilst the other one as male), representatives of two different patients associations focussed on rheumatologic and musculoskeletal diseases,

were involved in the design of the study and participated in the creation of the interview guide to ensure that the questions included were relevant for the studied population.³⁶ The patients' representatives did not attend the interviews and did not participate in the study.

3 | RESULTS

Thirteen Italian people from northern Italy agreed to participate in the interviews. Two participants were not able to do the interviews because they did not understand the questions on both telephone and videoconference since they had critical auditory impairments. Therefore, eleven participants were included. Table 3 reports the sample's demographic and clinical characteristics in detail for each participant.

TABLE 2 Data synthesis by extracting and abstracting findings in common themes and subthemes

| Abstraction: Themes | Abstraction: Subthemes | Codes defined by researchers | Example of quotes extracted from the interviews |
|-------------------------------------|---|---|--|
| Experiencing a sense of uncertainty | Need of a straightforward treatment | Need of guidelines to follow | '... Yes, I would like to have a precise guideline, also regarding nutrition... It looks as if there are some things that are left to our own intuition'. (P2, female, 68) |
| | | Precise treatment | 'Erm. I expected them to give me clear indications on how to deal with my disease'. (P8, male, 66) |
| | Doubts (treatments to follow and pathology genesis) | Lack of understanding of the disease's mechanisms | 'They tried to explain to me how OA works somehow, but I still don't have a clear idea of how it works'. (P3, female, 73) |
| | | Need to hear several health professionals to have a definitive answer | 'In my experience, I've had to consult two or three physicians, unless the first two agree'. (P5, female, 72) |
| | | Need to explain the disease set-up biomechanically | 'I thought [OA] was a consequence of bad posture, as I've been using my leg wrongly after slipping on ice once'. (P6, male, 55) |
| | Different opinions heard by various health professionals | Feeling that the physicians' decisions are not evidence-based | 'There is an almost religious way of thinking about how to deal with the pathology. It is not an exact science; when you choose the physicians, you choose the treatment'. (P1, male, 49) |
| | | Confusion caused by consulting different physicians | 'Maybe the fact that I did not have only one physician at the very beginning did not help me to understand how to deal with OA'. (P9, female, 73) |
| | Frustration and anger | Worrying for hearing different opinions | 'I was worried because we, as patients, hear different opinions coming from our friends and acquaintances that give us their personal point of view on how they take care of their disease'. (P9, female, 73) |
| | | Anger/frustration for different opinions | 'It is very frustrating for a patient [not to have a precise indication] because you expect to have a disease, and a common one too, so the care process should be clear'. (P1, male, 49) |
| | Establishing a challenging relationship with the self and the other | Not being understood and the importance of empathy | Seeing the patients as a diagnosis and not as human beings |
| Lack of empathy | | | 'The orthopaedists did not give me much attention, and they told me that I have OA and that I have to live with it'. (P2, female, 68) |
| Shame | | Feeling shameful | 'I felt it [shameful] recently, I went to the beach with my granddaughter [...] she wanted me to be involved in her games, and she said "Grandma come, sit down next to me". I had to kneel down to play in the sand with her... I felt, how can I say... erm... like a piece of wood, like someone who can no longer manage their body'. (P2, female, 68) |
| | | Limping as an unpleasant sensation | 'It is an unpleasant sensation, it feels as if you are limping, mentally though, you see? Because I do not know if it is visible or if that is only a perception [that I am limping]'. (P4, female, 47) |
| Hopelessness | | Hopelessness for the prognosis | |

(Continues)

TABLE 2 (Continued)

| Abstraction: Themes | Abstraction: Subthemes | Codes defined by researchers | Example of quotes extracted from the interviews |
|---|---|--|---|
| | | | 'When I received my diagnosis, they told me I had only few years left [before the surgery], and they told me "Chill and don't do anything". I asked the second orthopaedic who visited me "If I keep on being active will I undergo surgery in 5 years instead of 10 years?" and he answered, "You are quite optimistic in both scenarios". (P1, male, 49) |
| | | No possibilities to do other interventions | 'Erm. Yes [I can only do surgery], because I dragged it on for too long and they told me that I have no other possibilities with other [nonsurgical] interventions'. (P10, male, 65) |
| | Use of metaphors to describe the pathology | Using a relevant metaphor/simile | 'I see it [the joint affected by OA] as a mountain which is crumbling'. (P3, female, 73) |
| | | Associating OA to something realistic to understand it | 'My physiotherapist once told me that [OA] is like having a rusty gate, the orthopaedist decides to break it open, but if you try to grease it, it can last longer. I think that this gives you the idea'. (P1, male, 49) |
| <i>Being stuck in one's own or the health professionals' beliefs about the disease management</i> | Sealed faith (surgery) | Surgery as an obtrusive thought | 'It [OA surgery] is something you think about every day, something you try to resist, but that is your fate'. (P1, male, 49) |
| | | Surgery as the final and obvious stage of OA | 'However, everyone told me: try and resist for as long as you can, but sooner or later you will have to undergo surgery, and end up under the knife, full stop'. (P10, male, 65) |
| | OA as a pathology of the old adults | Misperception between radiographic findings and patients' perception | 'The doctor told me: "You know that if I did not know that these x-rays belong to you, I would think that they belong to another person who is at least 30 years older than you"... but, I guess I did not feel as bad as he was describing me'. (P11, male, 56) |
| | | OA as an ageing process | 'They told me that I was starting to get old'. (P3, female, 73) |
| | Necessity of radiographic findings (diagnose/treatment) | Diagnosis only through X-ray | 'They told me: here we have the problem, and it is evident as we can see from the x-ray'. (P11, male, 56) |
| | | OA as a wear and tear disease | 'Well, I had some medical check-ups through x-rays... and from them you could see some wear and tear joint surfaces, but they weren't uniform, nor regular, right? And so it was clear that there was something special even for a neophyte like me, right? And he [the doctor] said to me, "look, this is linked to this pathology." And by looking at it I became aware of what my problem was'. (P6, male, 55) |
| | (Ab)use of passive therapies | Recommendation of ice instead of movement in young patient | 'And he [the doctor] told me that I was too young for surgery and he recommended I do this therapy, to put some ice on my joint'. (P5, female, 72) |
| | | Recommended physical therapy for OA | 'The same rheumatologist I saw in September confirmed that the only thing I could do for my hip was Extracorporeal Shockwave Therapy'. (P4, female, 47) |

TABLE 2 (Continued)

| Abstraction: Themes | Abstraction: Subthemes | Codes defined by researchers | Example of quotes extracted from the interviews |
|--|---|--|--|
| | Movement as dangerous for the joint | Refrain from moving to postpone surgery | 'The doctor told me: "You have to try to postpone surgery for as long as you can. So please stop [any physical exercise]"'. (P1, male, 49) |
| | | Avoiding exercise with a load on the joint | 'They recommended I do pilates, go swimming etc. so as to go easy on the joint, not to go to the gym, not to run etc. Basically, avoid everything that could have a violent stress on the joint at the moment'. (P4, female, 47) |
| <i>Dealing with one's own attitudes towards the disease</i> | Fight, resignation and acceptance | Disease as something to fight | 'From a certain perspective, I found it [OA] positive, since it is something that I have to fight against'. (P1, male, 49) |
| | | Resignation towards getting older | 'Maybe I am accepting my becoming old, what can I say...'. (P3, female, 73) |
| | | Acceptance of the future | 'Besides, I am also a fatalist, things happen in life and when they do, you face them'. (P4, female, 47) |
| | Coping strategies (primarily passive) | (Over)use of drugs | 'It's not an issue for me to take some pills not to feel any pain'. (P4, female, 47) |
| | | Impromptu strategies to manage OA | 'I proceeded by myself with some clay, with some palliatives like unguents and things like that'. (P8, male, 66) |
| <i>Understanding the facilitators of the adherence to therapeutic exercise</i> | Importance of being active | Movement as an intrinsic need of the body | '...The body has to move...'. (P3, female, 73) |
| | | The awareness of the importance of movement | 'And, this kind of movement (walking), I realised, was good for me'. (P2, female, 68) |
| | Perceived exercises as concrete support to the cure | Witnessing the importance of training | 'Then the situation improved, always with training, thanks to the workouts'. (P7, female, 45) |
| | | Difference between active and passive treatments | 'It [physical exercise] is not like taking supplements with hyaluronic acid, those (supplements) you do not see what they do'. (P1, male, 49) |
| | Mean to maintain functionality | The importance of movement to maintain good functionality | 'OA is difficult to cure, even impossible, it is a natural tear, only some palliatives exist. I believe that the only way, or rather, the best way is to strengthen the muscle structure so that bones and joints suffer less from the weight load on them'. (P11, male, 56) |
| | | Movement as a way to reduce OA's impact | 'I felt well, because I kept on walking... and this allowed for reducing OA impact'. (P10, male, 65) |
| | Willingness to change life-habits | The importance of determination and willpower in active care | '...Determination and willpower [to change life-habits]'. (P7, female, 45) |
| | | Willpower as a compulsory step to change life habits | 'A great willpower is necessary [to change life-habits]'. (P9, female, 73) |
| <i>Understanding the barriers to the adherence to therapeutic exercise</i> | Cost and lack of time | Economic consequences of the care process | 'Yes... but also from an economic point of view [it is difficult to do supervised exercises]'. (P2, female, 68) |
| | | Being overwhelmed with life duties and forgetting about the self | 'We are trapped into a spiral in which work, we can say, takes up a lot of energy and a lot of time, and then that time is taken away from us...'. (P6, male, 55) |

(Continues)

TABLE 2 (Continued)

| Abstraction: Themes | Abstraction: Subthemes | Codes defined by researchers | Example of quotes extracted from the interviews |
|--|---|--|---|
| | Lack of clear indications | Do not know what to do with exercise | 'That is, there were some, just some things [decisions in the care process]... Erm... I don't know... they were left to our intuition, to our perception but just because you understand that by acting in a certain way, maybe you will limit its progress [of OA]...'. (P2, female, 68) |
| | | Missing the health professionals' real intention | 'The doctor told me: "You know that if I did not know that these x-rays belong to you, I would think that they belong to another person who is at least 30 years older than you"... but, I guess I did not feel as bad as he was describing me'. (P11, male, 65) |
| | Lack of willpower and fatigue in changing life habits | Losing the motivation with ageing | 'So it is that maybe when you are old, people back down, they lie on the couch,... Surely such a pain affecting someone who does not have that drive [motivation to stay fit] makes people unwilling to get up from the couch'. (P1, male, 49) |
| | | Laziness and fatigue in changing life habits | 'I think so, for laziness. Because if you want to, you are able to find the time. So it is, therefore, laziness'. (P5, female, 72) |
| | Exercise perceived useful only after surgery | Exercise useful only after undergoing surgery | 'But I imagine that someone can do this... let's call it preventive activity. Activity that can help with the recovery process following the intervention'. (P6, male, 55) |
| Exercise useless before surgery | | 'It is useless to start doing physiotherapy/exercise if I am undertaking surgery in a month'. (P9, female, 73) | |
| <i>Developing an uneasy relationship with food</i> | Diet as fatigue and deprivation | Eating to eat your own feelings | 'To follow a diet is a mental fatigue [...] and eating is an easy outlet to manage the stress of daily life'. (P4, female, 47) |
| | | Diet as deprivation | 'Think about it, if someone tells you something like "From tomorrow you will eat only these things [tasteless food]", I will only get a third of the satisfaction I normally get from eating...'. (P2, female, 68) |
| | Diet seen as useful only to lose weight | Diet to reduce the weight on the joints | 'Being overweight makes it worse, so obviously, the lighter I feel, the better I eat, and I also do my exercises, than [by doing so] I can see the difference'. (P7, female, 45) |
| | | Relationship between weight and OA | 'Of course, there is a relationship [between weight and OA]. The heavier the body, the more the knee suffers, it's a matter of physics'. (P2, female, 68) |

Abbreviation: OA, osteoarthritis.

Analysis of the interview data revealed seven main themes related to the OA care process (Figure 1): (1) *Experiencing a sense of uncertainty*; (2) *Establishing challenging relationships with the self and the other*; (3) *Being stuck in one's own or the health professionals' beliefs about the disease management; understanding*;

(4) *Dealing with one's own attitudes towards the disease; Understanding* (5) *the facilitators of and* (6) *the barriers to the adherence to therapeutic exercise*; (7) *Developing an uneasy relationship with food*. Hereafter, the different themes that stemmed from the synthesis of their related subthemes are discussed and explored.

TABLE 3 Participants' demographic and clinical characteristics

| Patient | Age | Gender | BMI | Retirement | Disease | Diagnosis |
|---------|-----|--------|-------|------------|-----------------|---------------|
| P1 | 49 | M | 26.28 | No | Hip OA | X-ray |
| P2 | 68 | F | 25.95 | Yes | Hip and knee OA | MRI |
| P3 | 73 | F | 27.34 | Yes | Knee OA | X-ray |
| P4 | 47 | F | 28.71 | No | Hip OA | MRI |
| P5 | 72 | F | 25.81 | No | Hip OA | X-ray and CT |
| P6 | 55 | M | 34.02 | No | Hip and knee OA | X-ray |
| P7 | 45 | F | 25.09 | No | Hip OA | X-ray and MRI |
| P8 | 66 | M | 28.70 | Yes | Knee OA | X-ray |
| P9 | 73 | F | 28.66 | Yes | Knee OA | X-ray |
| P10 | 65 | M | 24.69 | No | Hip OA | X-ray |
| P11 | 56 | M | 22.22 | No | Hip and knee OA | X-ray |

Abbreviations: BMI, body mass index; CT, computed tomography; F, female; M, male; MRI, magnetic resonance image; OA, osteoarthritis; P, person.



FIGURE 1 Themes and subthemes stemmed from the analysis of the interviews

Theme 1: Experiencing a Sense of Uncertainty

A general sense of *uncertainty* regarding the care process and, precisely, how to deal with OA and which treatments should have been taken, was a transversal perception among all the interviewees. They felt that the indications they had been given were not based on evidence and that health professionals' preferences and attitudes played a crucial role in the decision-making process of OA management.

There is an almost religious way of thinking about how to deal with the pathology. It is not an exact science; when you choose the

physicians, you choose the treatment. (P1, male, age 49)

Participants were typically unaware of the causes of their disease. However, in general, they considered it to be a consequence of overuse and wrong posture, and they tried to motivate it only through a biomechanical rationale. Moreover, the explanations provided to them by the health professionals seemed fuzzy and unclear.

I thought [OA] was a consequence of bad posture, as I've been using my leg wrongly after slipping on ice once. (P6, male, 55)

They tried to explain to me how OA works somehow, but I still don't have a clear idea of how it works. (P3, female, 73)

The interviewees felt doubtful because, on the one hand, they had not received any specific indication regarding the management of OA, and, on the other, the different and not coherent opinions retrieved from the various health professionals and acquaintances increased this sense of doubtfulness and worry about their condition.

I was worried because we, as patients, hear different opinions coming from our friends and acquaintances that give us their personal point of view on how they take care of their disease. (P9, female, 73)

Besides, this sense of uncertainty led the patients to retrieve information from a plethora of different professionals, word-of-mouth and also from the Internet.

In my experience, I've had to consult two or three physicians, unless the first two agree. (P5, female, 72)

No, the doctors did not explain it [OA] to me. But, eventually, I looked it up on the Internet and found answers to my questions. (P8, male, 66)

As a result, they felt frustrated and angry since they wanted to receive a straightforward treatment. The lack of clear indications and the presence of the conflicting opinions heard, eventually, led them not to take care of their disease or to base their care on their intuition.

It is very frustrating for a patient [not to have a precise indication] because you expect to have a disease, and a common one too, so the care process should be clear. (P1, male, 49)

Eventually, I did not do anything anymore, just nothing. (P5, female, 72)

...Yes, I would like to have a precise guideline, also regarding nutrition... It looks as if there are some things that are left to our intuition. (P2, female, 68)

Theme 2: Establishing Challenging Relationship with the Self and the Other

As far the relationship with the self and the other is concerned, common feelings across the interviewees revolve around a sense of shame and hopelessness that were described with two different nuances: the former was more concerned and present in front of their beloved ones, the latter was more evoked by the lack of hope conveyed by the

health professionals they consulted. Participants expressed a sense of shame caused by showing their conditions and limitations to others.

I felt it [shame] recently. I went to the beach with my granddaughter [...] she wanted me to be involved in her games, and she said 'Grandma come, sit down next to me'. I had to kneel down to play in the sand with her... I felt, how can I say... erm... like a piece of wood, like someone who can no longer manage their body. (P2, female, 68)

At the same time, they expressed a sense of hopelessness in regard to the received prognosis and the fact that physicians took for granted that this was a pathology that sees no other possibility apart from deteriorating.

Erm. Yes [I can only do surgery], because I dragged it on for too long and they told me that I have no other possibilities with other [non-surgical] interventions. (P10, male, 65)

Both of them were strictly related to a lack of empathy shown by both their acquaintances and health professionals. The former, since the interviewees reported that their acquaintances did not miss a chance to highlight some features related to OA (e.g., limping) that make them feel ashamed of their condition.

What annoys me the most is when people that know my condition ask me, 'What did you do? Why are you limping?' This makes me really upset because others see what I sometimes don't even notice. (P1, male, 49)

The latter, since the participants felt as if they were being considered as a mere number, rather than as human beings by the health professionals, they consulted, almost as if they were not worth their attention.

The orthopaedic surgeon did not give me much attention, and they told me that I have OA and that I have to live with it. (P2, female, 68)

Finally, some of the participants expressed their self-into-the-pathology through the use of relevant metaphors that helped them to understand their condition better.

I see it [the joint affected by OA] as a mountain which is crumbling. (P3, female, 73)

Theme 3: Being Stuck in One's Own or the Health Professionals' Beliefs about the Disease Management

The participants held some core beliefs that arose after consulting with their health professionals. In particular, the most

shared and common belief carried out was regarding the surgery that represented a sealed fate from which there is no escape, something that anyone with OA must eventually face.

It [OA surgery] is something you think about every day, something you try to resist, but that is your fate. (P1, male, 49)

And so when I went to see him [the physician] he said, 'no madam, your joint is ruined... try and get on with it for as long as you can, but sooner or later you will have to do it [surgery]'. (P9, female, 73)

Another shared belief was considering OA as a disease typical of old age. Having this belief made participants feel as if they were going through an unhealthy ageing process, even if some of them felt active and alive.

The doctor told me: 'You know that if I did not know that these x-rays belong to you, I would think that they belong to another person who is at least 30 years older than you'... but, I guess I did not feel as bad as he was describing me. (P11, male, 56)

In fact, according to the narratives collected, health professionals have been reported to be surprised when they saw a sign of OA radiographic findings in the younger interviewees. Besides, these radiographic findings were considered necessary to diagnose OA and to decide how to plan the care process, since interviewees reported that physicians were more focussed on radiographic findings than on the symptoms they were complaining about.

They told me: here we have the problem, and it is evident as we can see from the x-ray. (P11, male, 56)

Two other common beliefs emerged from the interviewees' encounters with their health professionals, concerned the management of OA. Firstly, the (ab)use of passive therapies that were recommended to postpone as much as possible the surgical interventions and, secondly, the low prescription of movement that was seen as dangerous as a risk factor for an anticipated surgery.

And he [the doctor] told me that I was too young for surgery, and he recommended I do this therapy, to put some ice on my joint. (P5, female, 72)

The doctor told me: 'You have to try to postpone surgery for as long as you can. So please stop [any physical exercise]'. (P1, male, 49)

Theme 4: Dealing with One's Own Attitudes Towards the Disease
Interviewees showed different attitudes towards the pathology that were linked by them to their age. The older patients perceive OA

as a sign of resignation towards becoming old, the younger ones as something they have to or can still fight against.

Maybe I am accepting my becoming old, what can I say... (P3, female, 73)

From a certain perspective, I took it [OA] positively since it is something I have to fight against. (P1, male, 49)

However, besides these differences, at a certain point, they all matured a sense of acceptance, as if OA is something they cannot change.

Besides, I am also a fatalist, things happen in life, and when they do, you face them. (P4, female, 47)

Both younger and older interviewees developed passive primarily coping strategies to manage their symptomatology. In particular, overuse of medications to control pain and the use of different physical therapies seemed to be the main traits of their care process, with partial room left to active therapies.

It's not an issue for me to take some pills not to feel any pain. (P4, female, 47)

I thought that by doing some thermal treatments [...] mud treatments [...] mesotherapy and other things... I thought that with them I would sort the disease out. (P5, female, 72)

Themes 5: Understanding the Facilitators of the Adherence to Therapeutic Exercise

Different facilitators of the adherence to therapeutic exercise were pointed out by the interviewees, among which the perceived benefit of doing exercises was the most shared one. In particular, being active was perceived as an intrinsic need of the body, allowing individuals to maintain a good level of functionality and as real and concrete support to the care process, something whose benefits could be seen in real-time compared to taking medications.

...The body has to move... (P3, female, 73)

It [physical exercise] is not like taking supplements with hyaluronic acid, those (supplements) you do not see what they do. (P1, male, 49)

Moreover, all the interviewees highlighted the importance of willpower, which was seen as either a facilitator or a barrier depending on its presence or lack thereof (see next theme). All interviewees agreed that their willingness to change their life habits was the key to sticking to the exercise plan and that willpower was a compulsory step to change life habits.

...Determination and willpower [to change life-habits]. (P7, female, 45)

Theme 6: Understanding the Barriers to the Adherence to Therapeutic Exercise

The barriers to the adherence to therapeutic exercise revolved around the cost of the therapy, the time needed, the lack of clear indications, the lack of willingness to change life habits and the perception of OA exercises useful only after surgery. The interviewees complained about the fact that exercise needs time, energy and money. Being absorbed in their jobs and family daily routines jeopardised their willingness to start doing physical exercises.

Yes... but also from an economic point of view [it is difficult to do supervised exercises]. (P2, female, 68)

We are trapped into a spiral in which work, we can say, takes up a lot of energy and a lot of time, and then that time is taken away from us... (P6, male, 55)

The lack of clear indications from specialists regarding the type of exercise to perform, as well as its intensity and frequency, were another barrier to starting any physical activity. The interviewees felt lost, as they perceived that they were missing precise guidelines to follow and that everything was left to their intuition.

My doctor told me to 'go for a walk', or maybe to 'move', but never specifically, something like 'it would be better in your case to do something more targeted'. (P11, male, 56)

The interviewees highlighted that the lack of willpower and the fatigue in changing life habits were some of the main reasons behind not sticking to the exercise plan together with losing the motivation with the ageing process.

I think so, for laziness... Because if you want to, you are able to find the time. So it is, therefore, laziness. (P5, female, 72)

Finally, feeling that exercise is useful only after surgery was another barrier to sticking to the exercise plan. The interviewees perceived that it was not valuable to invest time and energy in doing exercise if surgery was their final destiny.

It is useless to start doing physiotherapy/exercise if I am undertaking surgery in a month. (P9, female, 73)

The only preventive function that exercise seemed to have was to facilitate the recovery process after the surgery. The interviewees reported little to no appreciation of exercise as a treatment in its own right.

But I imagine that someone can do this... let's call it preventive activity. Activity that can help with the recovery process following the intervention. (P6, male, 55)

Theme 7: Developing an Uneasy Relationship with Food

Participants had developed an unhealthy relationship with food. As a matter of fact, they perceived diet as a deprivation, a sort of sacrifice in terms of time and mental fatigue, something that they forced themselves to follow. Moreover, for some of them, overeating was being used 'to eat your feelings'.

To follow a diet is a mental fatigue [...] and eating is an easy outlet to manage the stress of daily life. (P4, female, 47)

Furthermore, when it came to the management of OA, the participants considered following a diet only as a way to reduce weight on their joints, drawing a direct relationship between weight and joint load.

Of course, there is a relationship [between weight and OA]. The heavier the body, the more the knee suffers, it's a matter of physics. (P2, female, 68)

4 | DISCUSSIONS

The quality of the care process depends not only on the appropriateness of treatments delivered by health professionals (technical aspects) but also on relational and functional aspects.³⁷ From our results, several of these aspects were hindered during the OA care process, and this could be one of the possible explanations for why patients with this disease do not reach good levels of health-related outcomes and adherence towards first-line interventions.

As far as the technical aspects are concerned, patients did not receive first-line interventions. As a matter of fact, their care process was mainly based on passive therapies while waiting for surgery, with a scarce prescription of movement, seen not only as dangerous, but even as a possible risk factor for anticipated surgery, and valuable only after joint replacement. However, oral medications have been shown to reach similar effects to exercise therapy for improving functionality and pain relief in OA.³⁸ Nevertheless, oral medications, and in particular nonsteroidal anti-inflammatory drugs, have potential and well-documented side-effects, such as gastrointestinal toxicity, cardiovascular adverse effects and nephrotoxicity,³⁹ whereas exercise has been shown to have benefits that go beyond joint health, reducing the risk of developing a wide array of comorbidities and promoting a healthy lifestyle.⁴⁰ Moreover, evidence showed that individuals with OA are seeking

non-pharmacological and non-surgical treatments for their conditions, and they want more information about these treatments, which highlights the likelihood that they might be open to undergoing exercise and physiotherapy interventions if educated on their benefits.^{41,42} This was also retrieved in our interviews in which patients understood the importance of being active and saw exercise as a concrete support to their care, something that one can see while doing it, in contrast to medications and supplements that do not immediately show tangible effects.

According to the findings of this paper, there are several aspects that may facilitate or hinder patients' adherence towards physical exercise, and these should be taken into account by clinicians once it is prescribed. Regarding the obstacles, the lack of time and the cost seemed to be the main ones. The former was already known to be one of the more prominent causes of lack of adherence to physical exercise, in general.⁴³⁻⁴⁵ However, preliminary evidence showed that there is a gap between perceived and real lack of time⁴⁶ and that is why it is essential to address this issue while educating the patients, to help them reach a good level of awareness of the real-time they can devote to their own care, and to tailor self-management programmes according to their needs.⁴⁷ The latter, on the other hand, has to be tackled at a higher health-policy level. People in lower socioeconomic positions show a higher incidence of OA, more severe symptomatology and tend to experience lower benefits from OA interventions.^{48,49} However, despite Italy's public healthcare system, none of the participants who were being followed by professionals within the public system reported that exercise was included in their care process. There is a need to establish an effective care pathway for people with OA, to implement evidence-based treatments in the care routine employed in the public system, with the aim of reducing the healthcare inequalities faced by those patients who are not able to access alternative private systems.

Another barrier to the implementation of exercise into the patients' self-management routine was their unwillingness to change their habits. This was a shared barrier with the implementation of diet strategies into their care. It is already known that individuals do not engage in health-promoting behaviours, even though these can reduce mortality and contribute to their wellness.⁵⁰ However, our findings show that patients were not motivated by their health professionals to follow such treatments. As reported by Hardcastle et al.,⁵¹ there are several strategies that professionals can implement to foster patients' motivation, such as strategies targeting self-efficacy, outcome expectancies, effort and value beliefs, as well as motivational interviewing techniques, and these should be used and implemented for more effective communication. Moreover, as far as the diet is concerned, it was seen by the interviewees as applicable only to reduce weight on the joint. However, patients need to change their diet not only by focussing on losing weight but also because this has the potential to mitigate pro-inflammatory mediators, including cytokines, interleukins, histamine and free radicals that lead to increasing systematic inflammation.⁵²⁻⁵⁶ Patients should be made aware of the fundamental role of diet in their care process and should be guided to no longer see it as a liminal treatment.

The functional aspect can be defined as the basic expectations about how care is delivered, and it comprises, among others, the delivery of effective treatments by trusted professionals and the coordination and continuity of care.^{17,37} When considering the former, the interviewees did not perceive that the indications they had received were based on evidence, and this raised a general sense of uncertainty. Uncertainty is a poorly addressed and managed issue in healthcare, which can result in patients' poor or inefficient coping strategies and dysfunctional adaptation to illness, as well as in a conflicting relationship with health professionals.⁵⁷⁻⁵⁹ For what concerns the latter, the interviewees did not perceive the coordination and continuity of care as smooth and clear. In fact, the interviewees felt frustrated and angry, as they did not receive straightforward treatment. This was also powered by the fact that the different health professionals they had met had different and incoherent opinions, and therefore they perceived that indications about the OA decision-making process were based on health professionals' preferences and attitudes.

Patients desire empowerment and are keen to be actively involved in their own care process.⁴² If they do not find answers to their questions from a source (e.g., their health professional), they will seek them through other outlets, such as different professionals, acquaintances or the Internet.^{42,60} This is in line with what was observed in our study, where the sense of uncertainty experienced by the interviewees led them to seek information from different professionals, the Internet and word-of-mouth. However, as pointed out by Chou et al.,⁴² the patients' necessity to interrogate various sources can indeed stem from the dissatisfaction with the information retrieved from one source, but it can also arise from the patients' need to gather information from different and complementary sources to receive a tailored and holistic approach. In line with this, there rises a need to create ad hoc resources for people with OA whose content is based on solid evidence and that is tailored to their needs and expectations.

The relational aspect concerns all interactions between patients and health professionals.³⁷ The interviewees felt as if they were not being understood by both their relatives and health professionals, and they found a lack of empathy in the professionals they met. Empathy declines throughout medical formation, especially when it is not trained with specific interventions.^{61,62} Treatments devoid of empathy spoil the relationship between patients and health providers, and may lead to dissatisfied patients who are in turn discouraged to stick to the recommended interventions, resulting in poor health-related outcomes.⁶³ An empathetic communication style is a fundamental skill that health professionals need to learn and optimise because it can allow patients to reach good compliance towards treatments and, therefore, better outcomes. A strategy reported by patients for an efficient communication style was the use of metaphor to explain the pathology. Human beings rely on metaphor to understand the world around them. The use of relevant metaphors that tap into the patients' life and experience can ease the creation of a bond of trust between the health professional and the patient, leading to a truly therapeutic alliance,^{64,65} which is fundamental

when the care process of patients need a high level of patients' compliance, as in the case of OA.

In particular, this lack of empathy was found during the diagnosis and the prognosis of OA. The individuals interviewed extracted a sense of hopelessness from health professionals' words, which was derived from seeing OA as a pathology with a sealed faith—surgery—and a sign of their ageing process. This was also fostered by the reading of their radiographic findings that physicians found necessary to perform the diagnosis and commented with sentences such as "if I did not know that these x-rays belonged to you, I would think that they belong to someone who is at least 30 years older than you". However, radiographic findings are considered by CPGs, as complementary for the assessment of OA and should be considered only when other diseases are the suspected cause of the symptoms (e.g., infection, cancer, rheumatoid arthritis) or once the surgical intervention has been planned.⁶⁵ Besides, there is a weak association between the severity of radiographic findings, pain and disability levels^{28,66} and basing clinical decisions on imaging fosters the perception of OA as a wear-and-tear disease, which may, in turn, induce fear-avoidance behaviours.^{28,67} Finally, promoting surgery as the unique and real solution to OA may demotivate patients to change their life habits, considering the difficulty to follow diet and a physical activity programmes compared to the 'easy way out' of undergoing surgery.

Some limitations of this study need to be underlined. Firstly, the small sample size required to conduct a qualitative study limits the generalisability of the results.⁶⁸ Besides, all the interviewees lived in a similar geographical area (i.e., northern Italy), so that it is not possible to conclude that the results of this study may be transferrable to the other Italian regions. However, in Italy, there is a well-known negative gradient gap from North to South when it comes to the efficiency of the healthcare system.^{69,70} In light of this, it is possible that this study depicted the best-case scenario that patients with OA may experience. Moreover, the results gathered from this study may not be transferrable to other European countries, such as the ones comprised in the northern areas, due to the geopolitical differences between them.⁷¹ Conversely, they may be more applicable to other Mediterranean countries, considering the similarities in the health professionals' educational needs within the field of rheumatology.⁷¹ Secondly, the patients interviewed were at different stages of their care process, and their experience may change during the different stages. However, all patients agree with what emerged during the member checking phase. On the contrary, one of the strengths of this study is that it is the first one that takes into account the whole care process experienced by patients with OA from diagnosis to pre-surgery, highlighting some of the possible pitfalls that both patients and clinicians may encounter, and that can hinder the success of the intervention.

5 | CONCLUSIONS

This study highlights potential common themes in the experience of people with OA, with a focus on its care process, which should be taken into account to enhance the quality of it. People with hip

and knee OA seem to experience an uncertain care process. In particular, they experienced a lack of clear explanations, a lack of empathy, and a general, negative attitude towards first-line nonsurgical treatments. All those factors underline the importance of providing patients with adequate information through effective communication about the treatment options. By doing so, it will be possible to shift patients' beliefs and improve their awareness of the first-line treatments they should follow. This will enhance patient-centred treatments led by shared decision-making processes with patients, increasing their compliance towards first-line interventions and their skills to take care of their health and healthcare, with positive effects on their health-related outcomes and healthcare costs.

ACKNOWLEDGEMENTS

The researchers would like to thank Ugo Viora, Executive Manager of the Rheumatic Patients Association (Associazione Malati Reumatici, AMaR)—Piemonte, for his important insights during the creation of the interview guide and all the participants that joined the study, who dedicated their time to perform the interviews. This study was developed within the DINOGMI Department of Excellence framework of MIUR 2018–2022 (Legge 232 del 2016). This study is part of the project funded by the European Alliance of Associations for Rheumatology (EULAR) HPR: The EULAR health professionals research grant 2020.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

Simone Battista made substantial contributions to the conception or design of the work, the acquisition, analysis or interpretation of data. Mattia Manoni made substantial contributions to the conception or design of the work, the acquisition, analysis and interpretation of data. Andrea Dell'Isola made substantial contributions to the conception or design of the work, analysis, and interpretation of data. Mattia Manoni made substantial contributions to the conception or design of the work, and interpretation of data. Alvisa Palese made substantial contributions to the design of the work, the acquisition, analysis or interpretation of data. Marco Testa made substantial contributions to the conception and design of the work, the acquisition, analysis or interpretation of data. All authors drafted the work or revised it critically for important intellectual content. All authors approved the version to be published. All authors agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

ETHICS STATEMENT

Ethics approval was obtained from the Ethics Committee for University Research (CERA: Comitato Etico per la Ricerca di Ateneo), University of Genova (Approval date: 15 June 2020; CERA2020.07). The participants signed informed consent before participation.

DATA AVAILABILITY STATEMENT

Data are available upon reasonable request to the corresponding author.

ORCID

Simone Battista  <https://orcid.org/0000-0002-7471-1951>

REFERENCES

- Sebrant L, Jong M. What's the meaning of the concept of caring? a meta-synthesis. *Scand J Caring Sci.* 2020;35(2):353-365. doi:10.1111/scs.12850
- Mortari L, Saiani L. *Gestures and Thoughts of Caring: A Theory of Caring from the Voices of Nurses.* 1st ed. McGraw-Hill Education; 2014.
- Benedetti F. Placebo-induced improvements: how therapeutic rituals affect the patient's brain. *J Acupunct Meridian Stud.* 2012;5(3):97-103. doi:10.1016/j.jams.2012.03.001
- Testa M, Rossetti G. Enhance placebo, avoid nocebo: how contextual factors affect physiotherapy outcomes. *Man Ther.* 2016;24:65-74. doi:10.1016/j.math.2016.04.006
- Rossetti G, Carlino E, Testa M. Clinical relevance of contextual factors as triggers of placebo and nocebo effects in musculoskeletal pain. *BMC Musculoskelet Disord.* 2018;19(1):27. doi:10.1186/s12891-018-1943-8
- Hunter DJ, Schofield D, Callander E. The individual and socio-economic impact of osteoarthritis. *Nat Rev Rheumatol.* 2014;10(7):437-441. doi:10.1038/NRRHEUM.2014.44
- Jackson J, Iyer R, Mellor J, Wei W. The burden of pain associated with osteoarthritis in the hip or knee from the patient's perspective: a multinational cross-sectional study. *Adv Ther.* 2020;37(9):3985-3999. doi:10.1007/s12325-020-01445-4
- National Clinical Guidelines Centre. Osteoarthritis: Care and Management in Adults. *NICE Clin Guidelines.* 2014;177:1-498.
- Fernandes L, Hagen KB, Bijlsma JW, et al. EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. *Ann Rheum Dis.* 2013;72(7):1125-1135. doi:10.1136/annrheumdis-2012-202745
- Bannuru RR, Osani MC, Vaysbrot EE, et al. OARSJ guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis Cartilage.* 2019;27(11):1578-1589. doi:10.1016/j.joca.2019.06.011
- Katz JN, Arant KR, Loeser RF. Diagnosis and treatment of hip and knee osteoarthritis: a review. *JAMA.* 2021;325(6):568-578. doi:10.1001/jama.2020.22171
- Allen KD, Choong PF, Davis AM, et al. Osteoarthritis: models for appropriate care across the disease continuum. *Best Pract Res Clin Rheumatol.* 2016;30(3):503-535. doi:10.1016/j.berh.2016.09.003
- Swalthes L, Paskins Z, Dziedzic K, Finney A. Factors influencing the implementation of evidence-based guidelines for osteoarthritis in primary care: a systematic review and thematic synthesis. *Musculoskeletal Care.* 2020;18:1-10. doi:10.1002/msc.1452
- Basedow M, Esterman A. Assessing appropriateness of osteoarthritis care using quality indicators: a systematic review. *J Eval Clin Pract.* 2015;21(5):782-789. doi:10.1111/jep.12402
- Theis KA, Brady TJ, Sacks JJ. Where have all the patients Gone? Profile of US adults who report doctor-diagnosed arthritis but are not being treated. *J Clin Rheumatol.* 2018;25(8):341-347. doi:10.1097/rhu.0000000000000896
- Runciman WB, Hunt TD, Hannaford NA, et al. CareTrack: assessing the appropriateness of health care delivery in Australia. *Med J Aust.* 2012;197(2):100-105. doi:10.5694/MJA12.10510
- Chi-Lun-Chiao A, Chehata M, Broeker K, et al. Patients' perceptions with musculoskeletal disorders regarding their experience with healthcare providers and health services: an overview of reviews. *Arch Physiother.* 2020;10(1):17. doi:10.1186/s40945-020-00088-6
- Ferre F, de Belvis AG, Valerio L, et al. Italy: health system review. *Health Syst Transit.* 2014;16(4):1-168.
- Battista S, Salvioi S, Millotti S, Testa M, Dell'Isola A. Italian physiotherapists' knowledge of and adherence to osteoarthritis clinical practice guidelines: a cross-sectional study. *BMC Musculoskelet Disord.* 2021;22(1):380. doi:10.1186/s12891-021-04250-4
- Moustakas C. *Phenomenological Research Methods.* SAGE Publications, Inc.; 1994. doi:10.4135/9781412995658
- Spiegelberg H. *Doing Phenomenology: Essays on and in Phenomenology.* Springer; 1975.
- Matua GA, van der Wal DM. Differentiating between descriptive and interpretive phenomenological research approaches. *Nurse Res.* 2015;22(6):22-27. doi:10.7748/nr.22.6.22.e1344
- Korstjens I, Moser A. Series: practical guidance to qualitative research. Part 2: context, research questions and designs. *Eur J Gen Pract.* 2017;23(1):274-279. doi:10.1080/13814788.2017.1375090
- Moser A, Korstjens I. Series: practical guidance to qualitative research. Part 1: introduction. *Eur J Gen Pract.* 2017;23(1):271-273. doi:10.1080/13814788.2017.1375093
- Tong A, Sainsbury P, Craig J. Consolidated Criteria for Reporting Qualitative Research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care.* 2007;19(6):349-357. doi:10.1093/intqhc/mzm042
- Moser A, Korstjens I. Series: practical guidance to qualitative research. Part 3: sampling, data collection and analysis. *Eur J Gen Pract.* 2018;24(1):9-18. doi:10.1080/13814788.2017.1375091
- Egerton T, Diamond LE, Buchbinder R, Bennell KL, Slade SC. A systematic review and evidence synthesis of qualitative studies to identify primary care clinicians' barriers and enablers to the management of osteoarthritis. *Osteoarthritis Cartilage.* 2017;25(5):625-638. doi:10.1016/j.joca.2016.12.002
- Darlow B, Brown M, Thompson B, et al. Living with osteoarthritis is a balancing act: an exploration of patients' beliefs about knee pain. *BMC Rheumatol.* 2018;2(1):15. doi:10.1186/s41927-018-0023-x
- Hurley M, Dickson K, Hallett R, et al. Exercise interventions and patient beliefs for people with hip, knee or hip and knee osteoarthritis: a mixed methods review. *Cochrane Database Syst Rev.* 2018;2018(4):010842. doi:10.1002/14651858.CD010842.pub2
- Battista S, Dell'Isola A, Manoni M, Englund M, Palese A, Testa M. Experience of the COVID-19 pandemic as lived by patients with hip and knee osteoarthritis: an Italian qualitative study. *BMJ Open.* 2021;11(10):e053194. doi:10.1136/bmjopen-2021-053194
- Vaismoradi M, Turunen H, Bondas T. Content analysis and thematic analysis: implications for conducting a qualitative descriptive study. *Nurs Health Sci.* 2013;15(3):398-405. doi:10.1111/nhs.12048
- Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol.* 2006;3(2):77-101. doi:10.1191/1478088706qrp0630a
- Lindgren BM, Lundman B, Graneheim UH. Abstraction and interpretation during the qualitative content analysis process. *Int J Nurs Stud.* 2020;108:103632. doi:10.1016/j.jnurstu.2020.103632
- Korstjens I, Moser A. Series: practical guidance to qualitative research. Part 4: trustworthiness and publishing. *Eur J Gen Pract.* 2018;24(1):120-124. doi:10.1080/13814788.2017.1375092
- Birt L, Scott S, Cavers D, Campbell C, Walter F. Member checking: a tool to enhance trustworthiness or merely a nod to validation? *Qual Health Res.* 2016;26(13):1802-1811. doi:10.1177/1049732316654870

36. Baxter S, Clowes M, Muir D, Baird W, Broadway-Parkinson A, Bennett C. Supporting public involvement in interview and other panels: a systematic review. *Health Expect*. 2017;20(5):807-817. doi:10.1111/HEX.12491
37. Doyle C, Lennox L, Bell D. A systematic review of evidence on the links between patient experience and clinical safety and effectiveness. *BMJ Open*. 2013;3(1):e001570. doi:10.1136/BMJOPEN-2012-001570
38. Zhang W, Nuki G, Moskowitz RW, et al. OARSI recommendations for the management of hip and knee osteoarthritis: part III: changes in evidence following systematic cumulative update of research published through January 2009. *Osteoarthritis Cartilage*. 2010; 18(4):476-499. doi:10.1016/j.joca.2010.01.013
39. Wongrakpanich S, Wongrakpanich A, Melhado K, Rangaswami J. A comprehensive review of non-steroidal anti-inflammatory drug use in the elderly. *Aging Dis*. 2018;9(1):143-150. doi:10.14336/AD.2017.0306
40. Skou ST, Pedersen BK, Abbott JH, Patterson B, Barton C. Physical activity and exercise therapy benefit more than just symptoms and impairments in people with hip and knee osteoarthritis. *J Orthop Sports Phys Ther*. 2018;48(6):439-447. doi:10.2519/jospt.2018.7877
41. Vitaloni M, Botto-van Bemden A, Sciortino R, et al. A patients' view of OA: the Global Osteoarthritis Patient Perception Survey (GOAPPS), a pilot study. *BMC Musculoskelet Disord*. 2020;21(1):727. doi:10.1186/s12891-020-03741-0
42. Chou L, Ellis L, Papandony M. Patients' perceived needs of osteoarthritis health information: a systematic scoping review. *PLoS One*. 2018;13(4):e0195489. doi:10.1371/journal.pone.0195489
43. Escoto KH, Laska MN, Larson N, Neumark-Sztainer D, Hannan PJ. Work hours and perceived time barriers to healthful eating among young adults. *Am J Health Behav*. 2012;36(6):786-796. doi:10.5993/AJHB.36.6.6
44. Withall J, Jago R, Fox KR. Why some do but most don't. Barriers and enablers to engaging low-income groups in physical activity programmes: a mixed methods study. *BMC Public Health*. 2011;11:507. doi:10.1186/1471-2458-11-507
45. Herazo-Beltrán Y, Pinillos Y, Vidarte J, Crissien E, Suarez D, García R. Predictors of perceived barriers to physical activity in the general adult population: a cross-sectional study. *Braz J Phys Ther*. 2017; 21(1):44-50. doi:10.1016/j.bjpt.2016.04.003
46. Heesch K, Mässe L. Lack of time for physical activity: perception or reality for African American and Hispanic women? *Women Health*. 2004;39(3):45-62. doi:10.1300/J013V39N03_04
47. Jonas D, Ibuka Y, Russell L. How much time do adults spend on health-related self-care? Results from the American time use survey. *J Am Board Fam Med*. 2011;24(4):380-390. doi:10.3122/JABFM.2011.04.100260
48. Kiadaliri AA, Gerhardsson de Verdier M, Turkiewicz A, Lohmander LS, Englund M. Socioeconomic inequalities in knee pain, knee osteoarthritis, and health-related quality of life: a population-based cohort study in southern Sweden. *Scand J Rheumatol*. 2017; 46(2):143-151. doi:10.1080/03009742.2016.1181203
49. Reyes C, García-Gil M, Elorza JM, et al. Socio-economic status and the risk of developing hand, hip or knee osteoarthritis: a region-wide ecological study. *Osteoarthritis Cartilage*. 2015;23(8):1323-1329. doi:10.1016/j.joca.2015.03.020
50. Ford E, Zhao G, JTsai J, Li C. Low-risk lifestyle behaviors and all-cause mortality: findings from the National Health and Nutrition Examination Survey III Mortality Study. *Am J Public Health*. 2011; 101(10):1922-1929. doi:10.2105/AJPH.2011.300167
51. Hardcastle SJ, Hancox J, Hattar A, Maxwell-Smith C, Thøgersen-Ntoumani C, Hagger MS. Motivating the unmotivated: how can health behavior be changed in those unwilling to change? *Front Psychol*. 2015;6:835. doi:10.3389/fpsyg.2015.00835
52. Philpot U, Johnson MI. Diet therapy in the management of chronic pain: better diet less pain? *Pain Manag*. 2019;9(4):335-338. doi:10.2217/pmt-2019-0014
53. Philippou E, Petersson SD, Rodomar C, Nikiphorou E. Rheumatoid arthritis and dietary interventions: systematic review of clinical trials. *Nutr Res*. 2021;79(4):410-428. doi:10.1093/NUTRIT/NUAA033
54. Nikiphorou E, Santos EJF, Marques A, et al. 2021 EULAR recommendations for the implementation of self-management strategies in patients with inflammatory arthritis. *Ann Rheum Dis*. 2021;80(0):1285. doi:10.1136/ANNRHEUMDIS-2021-220249
55. Rondanelli M, Faliva MA, Miccono A, et al. Food pyramid for subjects with chronic pain: foods and dietary constituents as anti-inflammatory and antioxidant agents. *Nutr Res Rev*. 2018;31(1):131-151. doi:10.1017/S0954422417000270
56. Runhaar J, Beavers MJ, Miller GD, et al. Inflammatory cytokines mediate the effects of diet and exercise on pain and function in knee osteoarthritis independent of BMI. *Osteoarthritis Cartilage*. 2019; 27(8):1118-1123. doi:10.1016/j.joca.2019.04.009
57. Lillrank A. Back pain and the resolution of diagnostic uncertainty in illness narratives. *Soc Sci Med*. 2003;57(6):1045-1054. doi:10.1016/S0277-9536(02)00479-3
58. Lütze U, Archenholtz B. The impact of arthritis on daily life with the patient perspective in focus. *Scand J Caring Sci*. 2007;21(1):64-70. doi:10.1111/j.1471-6712.2007.00443.x
59. Etkind SN, Bristowe K, Bailey K, Selman LE, Murtagh FE. How does uncertainty shape patient experience in advanced illness? A secondary analysis of qualitative data. *Palliat Med*. 2017;31(2):171-180. doi:10.1177/0269216316647610
60. Fedutes B, Starz T, Vogt M, Weber R, Ansani N. Gauging patients' interest in an arthritis Web site. *Am J Health Syst Pharm*. 2004; 61(13):1338-1339. doi:10.1093/AJHP/61.13.1338
61. Hojat M, Vergare MJ, Maxwell K, et al. The devil is in the third year: a longitudinal study of erosion of empathy in medical school. *Acad Med*. 2009;84(9):1182-1191. doi:10.1097/ACM.0B013E3181B17E55
62. Nunes P, Williams S, Sa B, Stevenson K. A study of empathy decline in students from five health disciplines during their first year of training. *Int J Med Educ*. 2011;2:12-17. doi:10.5116/IJME.4D47.DDB0
63. Riess H. The science of empathy. *J Patient Exp*. 2017;4(2):74-77. doi:10.1177/2374373517699267
64. Hanne M. Diagnosis and metaphor. *Perspect Biol Med*. 2015;58(1):35-52. doi:10.1353/PBM.2015.0010
65. Chichirez C, Purcărea V. Interpersonal communication in healthcare. *J Med Life*. 2018;11(2):119-122.
66. Helminen EE, Sinikallio SH, Valjakka AL, Väisänen-Rouvali RH, Arokoski JP. Determinants of pain and functioning in knee osteoarthritis: a one-year prospective study. *Clin Rehabil*. 2016; 30(9):890-900. doi:10.1177/0269215515619660
67. Gunn AH, Schwartz TA, Arbeeva LS, et al. Fear of movement and associated factors among adults with symptomatic knee osteoarthritis. *Arthritis Care Res*. 2017;69(12):1826-1833. doi:10.1002/acr.23226
68. Sim J, Saunders B, Waterfield J, Kingstone T. Can sample size in qualitative research be determined a priori? *Int J Soc Res Methodol*. 2018;21(5):619-634. doi:10.1080/13645579.2018.1454643
69. Toth F. How health care regionalisation in Italy is widening the North-South gap. *Health Econ Policy Law*. 2014;9(3):231-249. doi:10.1017/S1744133114000012
70. Cutolo R, Battista S, Testa M. Actual and perceived level of scientific English across Italian physiotherapy courses: a cross-sectional study. *Healthcare*. 2021;9(9):1135. doi:10.3390/HEALTHCARE9091135

71. Vliet Vlieland TP, van den Ende CH, Alliot-Launois F, et al. Educational needs of health professionals working in rheumatology in Europe. *RMD Open*. 2016;2(2):e000337. doi:10.1136/rmdopen-2016-000337

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Battista S, Manoni M, Dell'Isola A, Englund M, Palese A, Testa M. Giving an account of patients' experience: a qualitative study on the care process of hip and knee osteoarthritis. *Health Expect*. 2022;1-17. doi:10.1111/hex.13468

Study 3

Factors Associated With Adherence to a Supervised Exercise Intervention for Osteoarthritis: Data From the Swedish Osteoarthritis Registry

Simone Battista,¹ Ali Kiadaliri,² Thérèse Jönsson,² Kristin Gustafsson,³ Martin Englund,² Marco Testa,⁴ and Andrea Dell'Isola²

Objective. To explore how lifestyle and demographic, socioeconomic, and disease-related factors are associated with supervised exercise adherence in an osteoarthritis (OA) management program and the ability of these factors to explain exercise adherence.

Methods. A cohort register-based study on participants from the Swedish Osteoarthritis Registry who attended the exercise part of a nationwide Swedish OA management program. We ran a multinomial logistic regression to determine the association of exercise adherence with the abovementioned factors. We calculated their ability to explain exercise adherence with the McFadden R^2 .

Results. Our sample comprises 19,750 participants (73% female, mean \pm SD age 67 ± 8.9 years). Among them, 5,862 (30%) reached a low level of adherence, 3,947 (20%) a medium level, and 9,941 (50%) a high level. After a listwise deletion, the analysis was run on 16,685 participants (85%), with low levels of adherence as the reference category. Some factors were positively associated with high levels of adherence, such as older age (relative risk ratio [RRR] 1.01 [95% confidence interval (95% CI) 1.01–1.02] per year), and the arthritis-specific self-efficacy (RRR 1.04 [95% CI 1.02–1.07] per 10-point increase). Others were negatively associated with high levels of adherence, such as female sex (RRR 0.82 [95% CI 0.75–0.89]), having a medium (RRR 0.89 [95% CI 0.81–0.98]) or a high level of education (RRR 0.84 [95% CI 0.76–0.94]). Nevertheless, the investigating factors could explain 1% of the variability in exercise adherence ($R^2 = 0.012$).

Conclusion. Despite the associations reported above, the poorly explained variability suggests that strategies based on lifestyle and demographic, socioeconomic, and disease-related factors are unlikely to improve exercise adherence significantly.

INTRODUCTION

In osteoarthritis (OA), exercise is considered a first-line intervention by international clinical practice guidelines (1,2) due to its ability to improve symptoms and levels of functionality (3,4). Exercise positively affects body weight, lipid metabolism, glycemic control, and systemic inflammation, preventing and treating OA-related chronic diseases (5). Despite these benefits, adherence to exercise in OA is suboptimal (6,7).

Adherence is described by the World Health Organization (WHO) as “the extent to which a person’s behavior, taking medication, following a diet, and/or executing lifestyle changes,

corresponds with agreed recommendations from a health care provider” (8). Poor adherence to exercise can severely compromise its long-term effectiveness, limiting its benefits (9). Considering the rising prevalence (10) and economic burden of OA (11), identifying factors associated with exercise adherence is fundamental to creating specific interventions to improve it.

Several elements have been hypothesized to be associated with exercise adherence, including lifestyle and demographic, socioeconomic, and disease-related factors (12–17). However, evidence on this topic arises mainly from other chronic conditions than OA, qualitative studies whose aims are not to generalize knowledge, as well as studies with small samples (12–19).

¹Simone Battista, MSc: Lund University, Lund, Sweden, and University of Genoa, Campus of Savona, Savona, Italy; ²Ali Kiadaliri, PhD, Thérèse Jönsson, PhD, Martin Englund, PhD, Andrea Dell'Isola, PhD: Lund University, Lund, Sweden; ³Kristin Gustafsson, MSc: Ryhov County Hospital Jönköping, Jönköping, and Linköping University, Linköping, Sweden; ⁴Marco Testa, PhD: University of Genoa, Campus of Savona, Savona, Italy.

Author disclosures are available online at <https://onlinelibrary.wiley.com/doi/10.1002/acr.25135>.

Address correspondence via email to Marco Testa, PhD, marco.testa@unige.it.

Submitted for publication November 17, 2022; accepted in revised form April 13, 2023.

SIGNIFICANCE & INNOVATIONS

- Though exercise is a first-line intervention in osteoarthritis (OA), levels of exercise adherence among people with OA are suboptimal. Several elements have been hypothesized to be associated with exercise adherence, including lifestyle and demographic, socioeconomic, and disease-related factors in conditions other than OA.
- Analyzing real-world data from a first-line intervention provided nationwide in Swedish primary care, we found that high levels of adherence were positively associated with increased age, frequent pain, walking difficulties, and higher levels of self-efficacy. Conversely, high levels of adherence were negatively associated with female sex, higher body mass index, and high socioeconomic positions. However, these factors could explain 1% of the exercise variability.
- In OA, strategies based on lifestyle and demographic, socioeconomic, and disease-related factors are unlikely to improve exercise adherence significantly. Therefore, to improve adherence significantly, we need to consider other elements.

Moreover, the WHO has stated that the combination of different factors, rather than a single one, determines adherence (8). In contrast, the abovementioned studies focused primarily on single factors and their measures of mean association with adherence (e.g., odds ratio). Relying just on measures of association corresponds to an abstraction that does not take into account the variability of individual-level effects (20).

Therefore, we aimed to investigate the associations between lifestyle and demographic, socioeconomic, and disease-related factors with adherence to supervised exercise as a part of an OA management program delivered nationwide in Swedish primary care. Furthermore, we aimed to investigate these factors' ability to explain exercise adherence variability.

MATERIALS AND METHODS

Study design and setting. This study is a cohort register-based study on individual-level data retrieved from the Swedish Osteoarthritis Registry (SOAR; for data on the OA management program) and the Longitudinal Integration Database for Health Insurance and Labour Market Studies (LISA) administered by Statistics Sweden (for data on socioeconomic positions). These data sets were merged using personal identity numbers unique to all citizens in Sweden.

SOAR includes data from approximately 195,000 people with OA who attended an OA management program provided nationwide by the Swedish health care system (21,22). This program has already been thoroughly described elsewhere (23,24). Briefly, it is composed of 2 parts: education and exercise. The

education part is mandatory, while the exercise part is optional. The education part is based on 3 sessions that revolve around the pathophysiology of the disease and its self-care management. The first 2 sessions are mandatory and held by a physiotherapist. The third is optional and held by a person with OA, trained as an OA communicator. The exercise (optional) part starts with an individual encounter with a physiotherapist to tailor the exercise program to the participants' needs and characteristics. At this point, participants can decide whether to exercise at home or with a physiotherapist. Those who decide to exercise with a physiotherapist are offered the opportunity to attend 12 sessions over 6 to 8 weeks (2 sessions/week) following OA Swedish clinical practice guidelines (25). LISA provides socioeconomic data such as cohabitation, institutionally based education level, employment, income, and residential area (26). The research was conducted in respect of the Declaration of Helsinki and reported following the Strengthening the Reporting of Observational studies in Epidemiology guidelines. Ethical approval was obtained from the Swedish Ethics Committee (Dnr: 2019-02570).

Population. The study cohort comprises all the participants in the SOAR with a first registration (baseline) between 2012 and 2015. We included only those who started the exercise group sessions supervised by the physiotherapists after the initial encounter with them. We selected participants with knee or hip OA who were recorded in the SOAR only once.

Variables. The level of adherence to the supervised exercise part, reported in the SOAR, is the dependent variable of this study. This is a predetermined categorical variable recorded by the physiotherapists and stratified on the number of sessions participants attended (low levels of adherence: 1–6 training sessions; medium levels of adherence: 7–9 training sessions; or high levels of adherence: 10–12 sessions). In this study, high levels of adherence represent >80% of the adherence with the recommended interventions (12 sessions) (25), which is typically considered a satisfactory level of adherence (27). The collected independent variables are reported hereafter and divided as demographic and lifestyle characteristics, socioeconomic characteristics, and disease-related characteristics.

Demographic and lifestyle characteristics. Participants' demographic and lifestyle characteristics were reported by the participants themselves at the baseline and recorded in the SOAR. These characteristics were assigned sex at birth (binary variable: male/female), age (continuous variable), body mass index (BMI; continuous variable computed from self-reported height and weight), weekly physical activity (continuous variable: hours) that was assessed with the question "How active are you during a regular, typical week?" (21), and health-related quality of life (HRQoL; continuous variable: EuroQoL 5-domain instrument visual analog scale [EQ-5D VAS]). In the EQ-5D VAS,

the respondents reported their perceived HRQoL on a VAS scale that scored from 0 (the worst possible) to 100 (the best possible). The EQ-5D VAS is part of the EQ-5D scale (28).

Socioeconomic characteristics. Each socioeconomic position indicator from the year before the enrollment to the SOAR was considered for the analysis. In particular, the following socioeconomic position factors were retrieved and categorized: living alone (binary variable: living alone/living with someone), institutionally based education level (categorical variable: low [primary school: 0–9 years], medium [secondary school up to postsecondary education <3 years: 10–14 years], or high [postsecondary education: ≥15 years]), employment (binary variable: employed/retired-unemployed), residential area (categorical variable: rural/suburban/urban) and the net income.

Residential area was classified based on the Swedish Association of Local Authorities and Regions classification of Swedish municipalities. Specifically, rural areas are smaller towns/urban areas and rural municipalities, suburban areas are medium-sized towns (≥40,000 inhabitants) and municipalities near medium-sized towns, and urban areas are large cities (≥200,000 inhabitants) and municipalities near large cities (29). The individual yearly net income was categorized into quartiles based on the sample income distribution: lowest income quartile (<146,500 Swedish krona [SEK]), second income quartile (146,501–198,100 SEK), third income quartile (198,101–278,800 SEK), and highest income quartile (>278,800 SEK) (29).

Disease-related characteristics. The physiotherapists recorded the index joint (categorical variable: hip or knee) (21), namely, the joint with OA. They assessed this variable based on the participant's medical history, symptoms, and clinical assessment. In the case of multiple joints with OA, the most symptomatic joint was considered the index joint for the treatment. The participants self-recorded the numbers of painful joints (continuous variable); their desire for surgery (binary variable: yes/no) that was assessed by asking them: "Are your knee/hip symptoms so severe that you wish to undergo surgery?" (21); their pain intensity (ordinal variable: 0–10 on a numeric rating scale [NRS] (30)) in their index joint; their pain frequency (binary variable: infrequent pain [less than every week], frequent pain [almost every day]) that was assessed with the question: "How often do you have pain in your knee/hip" (21); their fear of movement (binary variable: yes/no) that was assessed with the question "Are you afraid your joints will be injured by physical training/activity?"; the Charnley score (categorical variable: A = unilateral hip or knee OA, B = bilateral hip or knee OA, C = multiple joint OA or some other condition) that categorizes people with OA into 3 classes based on the diseases that affect walking ability (31); and arthritis-specific self-efficacy (continuous variable: 10–100, pain and symptoms on the Arthritis Self-Efficacy Scale [ASES], using the Swedish version of the scale) (32). The ASES scale is a reliable instrument that

assesses patients' arthritis-specific self-efficacy, namely, their beliefs about their ability to perform a specific task and cope with OA (33). The full version is composed of 3 subscales: 1) self-efficacy pain scale (5 items), 2) function scale (9 items), and 3) other symptoms scale (6 items). Participants indicate to what extent they feel confident they can do the tasks reported in the items from 10 (very uncertain) to 100 (very certain). In the SOAR, only 1) and 3) were adopted and combined as suggested in the scale instruction (33).

Statistical analysis. Descriptive statistics are reported as mean ± SD and absolute and percentage frequencies. A multivariable exploratory analysis was performed to identify which independent variables were independently associated with exercise adherence in the SOAR (34). Multivariable exploratory analyses detect patterns and identify relationships between the independent variables and the outcome (34–36).

Since the proportional odds assumption was not met, an ordered logistic regression could not be performed. Hence, we ran a multinomial logistic regression with a listwise deletion (Stata function `mlogit`) to determine the association between the independent variables and the adherence to exercise. No missing data were reported in the outcome (adherence). Less than 1% of the data on socioeconomic characteristics was missing, primarily due to an error during the data upload process in LISA. Missing data on demographic and lifestyle and disease-related characteristics in the SOAR are most likely a result of a mistake by the physiotherapists responsible for uploading the data at the local unit. Hence missing data in both registers could be considered missing completely at random, introducing no or minimal bias in our analysis.

The selection of the variables in the model was informed by previous literature on exercise adherence in other chronic pain conditions (12–17) and the evidence for action on adherence by the WHO (8). Then, the variables were clustered in demographic and lifestyle, socioeconomic, and disease-related groups, following the dimensions proposed by the WHO (8). The multicollinearity assumption between continuous variables was tested, and none of the continuous variables was highly correlated. The relative risk ratio (RRR) of being in medium level of adherence or high level of adherence with respect to low level of adherence and 95% confidence intervals (95% CIs) were estimated for each covariate in the model. For the variables HRQoL and arthritis-specific self-efficacy, the RRR is presented as a 10-point change in their scales.

Finally, the ability of the models to explain the variability of exercise adherence was calculated with the McFadden R^2 statistic (Stata function `fitstat`). McFadden R^2 measures the ability of a model to explain the variance of dependent variables on a convenient 0–100% scale. In particular, this value highlights how much of the variance in the dependent variable (adherence) can be explained by the independent variables collectively. We calculated McFadden R^2 for the model with all variables included (full model).

Afterward, we excluded 1 set of variables from the model and calculated the difference between McFadden R^2 with the full model. A higher difference would indicate a higher contribution of the variables set into the explanatory power of the full model. The analysis was done through Stata 17.

RESULTS

Between January 1, 2012 and December 31, 2015, 46,905 people with OA were recorded in the SOAR. However, we excluded 7 participants who had joints other than hip and knee as their first cause of pain, 27,147 who did not perform any supervised exercise sessions, and 1 for attending the program more than once. Hence, 19,750 participants with knee (69%) and hip (31%) OA were included in this study (73% female, mean \pm SD age 67 ± 8.9 years). Figure 1 shows the participants' selection process. Table 1 presents the characteristics of the entire sample and stratified by the levels of adherence. Specifically, 5,862 (30%) reached a low level of adherence, 3,947 (20%) a medium level, and 9,941 (50%) a high level.

After the listwise deletion, the multinomial logistic regression was run on 16,685 individuals (85%), using low levels of

adherence as the reference category (Table 2). Overall, excluded participants ($n = 3,065$) had similar characteristics to the ones included in the analysis (see Supplementary Table 1, available on the *Arthritis Care & Research* website at <http://onlinelibrary.wiley.com/doi/10.1002/acr.25135>). We found that female sex (RRR 1.13 [95% CI 1.02–1.27]), living with someone (RRR 1.21 [95% CI 1.10–1.32]), and an increase of 1 number of joints with OA (RRR 1.06 [95% CI 1.01–1.10]) were positively associated with achieving medium levels of adherence. Conversely, an increase in an hour of weekly physical activity (RRR 0.98 [95% CI 0.96–0.99]), living in an urban area (RRR 0.87 [95% CI 0.78–0.98]), and being employed (RRR 0.82 [95% CI 0.72–0.93]) were negatively associated with achieving medium levels of adherence.

An increase of 1 year in age (RRR 1.01 [95% CI 1.01–1.02]), having frequent pain (RRR 1.13 [95% CI 1.02–1.25]), having walking difficulties (RRR 1.12 [95% CI 1.01–1.24]), and having a 10-point increase on the ASES (RRR 1.04 [95% CI 1.02–1.07]) were positively associated with high levels of adherence. By contrast, female sex (RRR 0.82 [95% CI 0.75–0.89]), an increase of 1 point in BMI (RRR 0.99 [95% CI 0.98–0.99]), living in a suburban (RRR 0.79 [95% CI 0.73–0.86]) or an urban area (RRR 0.78 [95% CI 0.71–0.86]), being employed (RRR 0.71 [95% CI 0.64–0.78]),




Figure 1. Selection of the study population. SOAR = Swedish Osteoarthritis Registry.

Table 1. Descriptive statistics*

| Variables | Total sample (n = 19,750) | Level of adherence | | |
|--|------------------------------|--------------------|-----------------------|---------------------|
| | | Low (n = 5,862) | Medium (n = 3,947) | High (n = 9,941) |
| Demographic and lifestyle characteristics | | | | |
| Assigned sex at birth | n = 19,750 | n = 5,862 | n = 3,947 | n = 9,941 |
| Male | 5,421 (27.45) | 1,519 (25.91) | 925 (23.44) | 2,977 (29.95) |
| Female | 14,329 (72.55) | 4,343 (74.09) | 3,022 (76.65) | 6,964 (70.05) |
| Age | n = 19,750 | n = 5,862 | n = 3,947 | n = 9,941 |
| Mean \pm SD | 66.86 \pm 8.94 | 65.87 \pm 9.39 | 66.47 \pm 9.01 | 67.60 \pm 8.57 |
| Body mass index | n = 19,381 | n = 5,735 | n = 3,867 | n = 9,779 |
| Mean \pm SD | 27.56 \pm 4.76 | 27.73 \pm 4.90 | 27.75 \pm 4.89 | 27.43 \pm 4.63 |
| HRQoL (EQ-5D VAS, 0–100) | n = 17,933 | n = 5,317 | n = 3,592 | n = 9,024 |
| Mean \pm SD | 65.82 \pm 19.22 | 65.84 \pm 19.37 | 65.74 \pm 19.35 | 65.85 \pm 19.07 |
| Weekly physical activity, hours | n = 18,050 | n = 5,364 | n = 3,606 | n = 9,080 |
| Mean \pm SD | 4.11 \pm 2.53 | 4.14 \pm 2.53 | 4.03 \pm 2.49 | 4.13 \pm 2.54 |
| Socioeconomic characteristics | | | | |
| Institutionally based education level | n = 19,699 | n = 5,862 | n = 3,938 | n = 9,918 |
| Low | 4,331 (21.99) | 1,170 (20.02) | 795 (20.19) | 2,366 (23.86) |
| Medium | 9,843 (49.97) | 2,962 (50.69) | 2,007 (50.96) | 4,874 (49.14) |
| High | 5,525 (28.05) | 1,711 (29.28) | 1,136 (28.85) | 2,678 (27.00) |
| Income quartile | n = 19,738 | n = 5,858 | n = 3,945 | n = 9,935 |
| Lowest | 4,942 (25.04) | 1,345 (22.96) | 1,022 (25.91) | 2,575 (25.92) |
| Second | 4,936 (25.01) | 1,393 (23.78) | 982 (24.89) | 2,561 (25.78) |
| Third | 4,929 (24.97) | 1,517 (25.90) | 976 (24.74) | 2,436 (24.52) |
| Highest | 4,931 (24.98) | 1,603 (27.36) | 965 (24.46) | 2,363 (23.78) |
| Area of living | n = 19,738 | n = 5,858 | n = 3,945 | n = 9,935 |
| Rural | 6,047 (30.64) | 1,667 (28.46) | 1,180 (29.91) | 3,200 (32.21) |
| Suburban | 8,252 (41.81) | 2,435 (41.57) | 1,708 (43.30) | 4,109 (41.36) |
| Urban | 5,439 (27.56) | 1,756 (29.98) | 1,057 (26.79) | 2,626 (26.43) |
| Employment | n = 19,738 | n = 5,858 | n = 3,945 | n = 9,935 |
| Unemployed | 12,244 (62.03) | 3,275 (55.91) | 2,394 (60.68) | 6,575 (66.18) |
| Employed | 7,494 (37.97) | 2,583 (44.09) | 1,551 (39.32) | 3,360 (33.82) |
| Living alone | n = 19,738 | n = 5,858 | n = 3,945 | n = 9,935 |
| Living alone | 7,754 (39.28) | 2,411 (41.16) | 1,457 (36.93) | 3,886 (39.11) |
| Living with someone | 11,984 (60.72) | 3,447 (58.84) | 2,488 (63.07) | 6,049 (60.89) |
| Disease-related characteristics | | | | |
| Worst joint | n = 19,750 | n = 5,862 | n = 3,947 | n = 9,941 |
| Hip | 6,049 (30.63) | 1,708 (29.14) | 1,188 (30.10) | 3,153 (31.72) |
| Knee | 13,701 (69.37) | 4,154 (70.86) | 2,759 (69.90) | 6,788 (68.28) |
| Pain intensity (NRS 0–10) | n = 19,686 | n = 5,843 | n = 3,935 | n = 9,908 |
| Mean \pm SD | 5.25 \pm 1.83 | 5.23 \pm 1.85 | 5.24 \pm 1.87 | 5.26 \pm 1.80 |
| Pain frequency | n = 19,700 | n = 5,842 | n = 3,940 | n = 9,918 |
| Infrequent | 3,436 (17.44) | 1,100 (18.83) | 723 (18.35) | 1,613 (16.26) |
| Frequent | 16,264 (82.56) | 4,742 (81.17) | 3,217 (81.65) | 8,305 (83.74) |
| Number of painful joints | n = 19,750 | n = 5,862 | n = 3,947 | n = 9,941 |
| Mean \pm SD | 1.94 \pm 1.29 | 1.95 \pm 1.28 | 2.00 \pm 1.32 | 1.91 \pm 1.27 |
| Charnley score | n = 19,735 | n = 5,855 | n = 3,946 | n = 9,934 |
| A | 6,814 (34.53) | 2,000 (34.16) | 1,340 (33.96) | 3,474 (34.97) |
| B | 3,437 (17.42) | 1,009 (17.23) | 686 (17.38) | 1,742 (17.54) |
| C | 9,484 (48.06) | 2,946 (48.61) | 1,920 (48.66) | 4,718 (47.49) |
| Walking difficulties | n = 19,651 | n = 5,835 | n = 3,932 | n = 9,884 |
| No | 3,472 (17.67) | 1,105 (18.94) | 731 (18.59) | 1,636 (16.55) |
| Yes | 16,179 (82.33) | 4,730 (81.06) | 3,201 (81.41) | 8,248 (83.45) |
| Fear of movement | n = 19,651 | n = 5,821 | n = 3,928 | n = 9,902 |
| No | 16,562 (84.28) | 4,871 (83.68) | 3,303 (84.09) | 8,388 (84.71) |
| Yes | 3,089 (15.72) | 950 (16.32) | 625 (15.91) | 1,514 (15.29) |
| Desire for surgery | n = 19,558 | n = 5,798 | n = 3,906 | n = 9,854 |
| No | 14,936 (76.37) | 4,441 (76.60) | 3,017 (77.24) | 7,478 (75.89) |
| Yes | 4,622 (23.63) | 1,357 (23.40) | 889 (22.76) | 2,376 (24.11) |
| ASES pain and symptoms (0–100) | n = 19,149 | n = 5,660 | n = 3,834 | n = 9,655 |
| Mean \pm SD | 65.54 \pm 16.43 | 65.44 \pm 16.54 | 65.51 \pm 16.62 | 65.61 \pm 16.28 |

* Values are the number (%) unless indicated otherwise. To calculate the missing values, subtract the number of participants listed in the second column (Total sample) from the total sample size of 19,750. ASES = Arthritis Self-Efficacy Scale; EQ-5D VAS = EuroQol 5-domain instrumental visual analog scale; HRQoL = health-related quality of life; NRS = numeric rating scale.

Table 2. Association between exercise adherence and investigated factors (n = 16,685)*

| Variables | P | RRR (95% CI for EXP[B]) |
|---------------------------------------|-------|-------------------------|
| Medium levels of adherence | | |
| Assigned sex at birth | | |
| Male (base category) | – | – |
| Female | 0.03 | 1.13 (1.02–1.27) |
| Age | 0.14 | 1.00 (0.99–1.01) |
| Body mass index | 0.37 | 0.99 (0.99–1.01) |
| HRQoL (EQ-5D VAS, 0–100)† | 0.57 | 0.99 (0.97–1.02) |
| Weekly physical activity, hours | 0.02 | 0.98 (0.96–0.99) |
| Institutionally based education level | | |
| Low (base category) | – | – |
| Medium | 0.88 | 0.99 (0.88–1.12) |
| High | 0.63 | 0.97 (0.84–1.11) |
| Income quartile | | |
| Lowest (base category) | – | – |
| Second | 0.71 | 0.98 (0.86–1.11) |
| Third | 0.63 | 0.97 (0.84–1.11) |
| Highest | 0.41 | 0.94 (0.81–1.09) |
| Area of living | | |
| Rural (base category) | – | – |
| Suburban | 0.27 | 0.94 (0.85–1.05) |
| Urban | 0.02 | 0.87 (0.78–0.98) |
| Employment | | |
| Unemployed (base category) | – | – |
| Employed | <0.01 | 0.82 (0.72–0.93) |
| Living alone | | |
| Living alone (base category) | – | – |
| Living with someone | <0.01 | 1.21 (1.10–1.32) |
| Worst joint | | |
| Hip (base category) | – | – |
| Knee | 0.35 | 0.95 (0.86–1.05) |
| Pain intensity (NRS 0–10) | 0.49 | 1.01 (0.98–1.04) |
| Pain frequency | | |
| Infrequent (base category) | – | – |
| Frequent | 0.80 | 0.98 (0.87–1.11) |
| Number of painful joints | 0.01 | 1.06 (1.01–1.10) |
| Charnley score | | |
| A (base category) | – | – |
| B | 0.99 | 0.99 (0.97–1.15) |
| C | 0.13 | 0.91 (0.81–1.03) |
| Walking difficulties | | |
| No (base category) | – | – |
| Yes | 0.93 | 0.99 (0.88–1.13) |
| Fear of movement | | |
| No (base category) | – | – |
| Yes | 0.49 | 1.04 (0.92–1.18) |
| Desire for surgery | | |
| No (base category) | – | – |
| Yes | 0.26 | 0.94 (0.83–1.05) |
| ASES pain and symptoms (0–100)† | 0.29 | 1.02 (0.99–1.05) |
| High levels of adherence | | |
| Assigned sex at birth | | |
| Male (base category) | – | – |
| Female | <0.01 | 0.82 (0.75–0.89) |
| Age | <0.01 | 1.01 (1.01–1.02) |
| Body mass index | 0.01 | 0.99 (0.98–0.99) |
| HRQoL (EQ-5D VAS, 0–100)† | 0.18 | 0.98 (0.96–1.01) |
| Weekly physical activity, hours | 0.79 | 0.99 (0.98–1.01) |
| Institutionally based education level | | |
| Low (base category) | – | – |
| Medium | 0.02 | 0.89 (0.81–0.98) |

(Continued)

Table 2. (Cont'd)

| Variables | P | RRR (95% CI for EXP[B]) |
|---------------------------------|-------|-------------------------|
| High | <0.01 | 0.84 (0.76–0.94) |
| Income quartile | | |
| Lowest (base category) | – | – |
| Second | 0.79 | 1.01 (0.91–1.13) |
| Third | 0.61 | 1.03 (0.92–1.15) |
| Highest | 0.95 | 1.00 (0.89–1.14) |
| Area of living | | |
| Rural (base category) | – | – |
| Suburban | <0.01 | 0.79 (0.73–0.86) |
| Urban | <0.01 | 0.78 (0.71–0.86) |
| Employment | | |
| Unemployed (base category) | – | – |
| Employed | <0.01 | 0.71 (0.64–0.78) |
| Living alone | | |
| Living alone (base category) | – | – |
| Living with someone | 0.29 | 1.04 (0.97–1.12) |
| Worst joint | | |
| Hip (base category) | – | – |
| Knee | 0.03 | 0.92 (0.85–0.99) |
| Pain intensity (NRS 0–10) | 0.12 | 1.02 (0.99–1.04) |
| Pain frequency | | |
| Infrequent (base category) | – | – |
| Frequent | 0.02 | 1.13 (1.02–1.25) |
| Number of painful joints | 0.50 | 1.01 (0.98–1.05) |
| Charnley score | | |
| A (base category) | – | – |
| B | 0.74 | 1.02 (0.91–1.14) |
| C | 0.11 | 0.93 (0.84–1.02) |
| Walking difficulties | | |
| No (base category) | – | – |
| Yes | 0.03 | 1.12 (1.01–1.24) |
| Fear of movement | | |
| No (base category) | – | – |
| Yes | 0.93 | 1.00 (0.91–1.11) |
| Desire for surgery | | |
| No (base category) | – | – |
| Yes | 0.44 | 0.96 (0.88–1.06) |
| ASES pain and symptoms (0–100)† | <0.01 | 1.04 (1.02–1.07) |

* Low levels of adherence are the reference. 95% CI = 95% confidence interval; ASES = Arthritis Self-Efficacy Scale; EQ-5D VAS = EuroQoL 5-domain instrument visual analog scale; HRQoL = health-related quality of life; NRS = numeric rating scale; RRR = relative risk ratio.

† RRR is reported as an increase of 10 points in the scale.

having a medium (RRR 0.89 [95% CI 0.81–0.98]) or a high level of institutionally based education (RRR 0.84 [95% CI 0.76–0.94]), and having the knee as the worst joint (RRR 0.92 [95% CI 0.85–0.99]) were negatively associated with high levels of adherence.

Finally, the McFadden R^2 of the full model suggested that participants' demographic and lifestyle characteristics, socioeconomic characteristics, and disease-related characteristics can explain approximately 1.2% of the variation in adherence. After we removed participants' demographic and lifestyle characteristics, socioeconomic characteristics, and disease-related characteristics alternatively, there was a difference in the McFadden R^2 with respect to the full model of 0.3%, 0.4%, and 0.2%, respectively. Disease-related characteristics had the most

explanatory power, albeit the total explanatory ability of the full model was very small.

DISCUSSION

This study is the first to try to understand the relationship between demographic and lifestyle, socioeconomic, and disease-related factors, with the level of adherence to a face-to-face supervised exercise program for OA in a large sample of participants with this disease. Of the total sample, approximately 30% had low adherence levels, 20% had medium adherence levels, and 50% had high adherence levels. The distribution of adherence levels in our sample is consistent with that of participants in a similar Danish intervention (37) but differs from the distribution observed in an online version of the same intervention, which had a higher proportion of people with high levels of adherence than our sample (38). While several factors were associated with adherence, the full model could explain only 1% of the variability, which suggests that these factors are unlikely to have a tangible impact on adherence.

Regarding demographic and lifestyle factors, female sex was negatively associated with a high level of adherence. Previous evidence has indicated that women (with or without OA) might face societal expectations of household and caregiving responsibilities, experiencing greater difficulty finding time to exercise (39–43). However, in the digital version of this intervention, female sex suggested a positive association with high levels of exercise adherence (38), suggesting that digital interventions may be more convenient for females. Despite these findings, addressing the root causes of these disparities in exercise adherence is crucial, rather than focusing on exercise delivery mode to reduce this sex gap. However, our study only collected information on participants' assigned sex at birth, limiting the generalizability of our results to those individuals who are not cisgender. Therefore, further research is needed to explore the relationship between gender identity, sex, and exercise adherence in individuals with OA. In addition, participants' older age was positively associated with reaching a high level of adherence. Considering how exercise is delivered in this program, our result aligns with previous evidence where older adults adhered more to self-paced rather than moderate-intensity exercise (44). Finally, BMI was negatively associated with reaching high levels of adherence, which is consistent with previous evidence where people with high BMI are less keen on engaging in physical exercise (38,45).

Among the socioeconomic factors, people who lived in an urban or suburban area, were employed, and had medium or high levels of institutionally based education tended to exercise less than their counterparts. Similar results were found in the digital version of this intervention, where lower institutionally based education and living outside the largest Swedish cities were associated with higher adherence (38). These results contrast with the previous literature, where socioeconomic categories typically representing higher socioeconomic positions tended to adhere

more to exercise (46,47). However, it is essential to consider that most of the data on adherence were retrieved from secondary analyses of randomized controlled trials (RCTs) (48). First, these studies were not designed to study adherence. RCTs per se tend to enhance adherence to treatment, which might create an over-estimation of the factors related to adherence (49).

Second, in RCTs, people are volunteers who are selected following specific inclusion and exclusion criteria, which may fail to mirror the socioeconomic variability of the underlying population from which the sample is drawn (50). Moreover, we might not have reached the more socioeconomically disadvantaged groups, considering the higher socioeconomic positions of the SOAR sample compared to the general Swedish population (29). Finally, another explanation of this tendency is that people in lower socioeconomic positions seemed exposed to a more detrimental OA-disease burden than their higher counterparts (51). Severe symptoms can act as a motivator and drive exercise adherence (46,52). Those who experience a higher disease burden might be more motivated to follow exercise regimens. This phenomenon was also highlighted in our study when looking at the disease-related factors, as having frequent pain and walking difficulties were associated with high levels of adherence.

Moreover, self-efficacy was associated with exercise adherence, as per previous evidence (53), but with a modest RRR. Self-efficacy is characterized by a curvilinear (U-shaped) relationship between this construct and task accomplishment (54). People with low self-efficacy are likely to doubt their chance to accomplish a task, and those with a high-self efficacy might be characterized by complacency, inadequate preparation, and a focus on achieving task-related targets (54). Therefore, low and high levels of self-efficacy can lead to a similar outcome, namely, low adherence to a task (e.g., exercise). Considering the large cohort of our study, the effect of self-efficacy might be diluted due to the high variety of our population.

However, our model could explain just 1% of the variability, as indicated by the McFadden R^2 . Thus, if we wanted to design an exercise intervention and understand which strategies to adopt to increase adherence, we should accept that demographic and lifestyle, socioeconomic, and disease-related factors are unlikely to improve adherence significantly, considering how little they explain adherence variability. This conclusion is further supported by the limited ability of similar factors to explain exercise adherence in the digital version of the intervention (38). Therefore, other factors should be taken into account.

The SOAR gathers real-world data from >500 different units throughout Sweden, with considerable variability among them. These contexts are characterized by specific contextual factors (e.g., structures' facilities, clinicians' communication style and ability to motivate patients, etc.) that affect people's outcomes via a placebo (or nocebo) response if positively (placebo) or negatively (nocebo) encoded by the brain via the so-called "mindsets" (55). Mindsets are "core assumptions about a domain or category that

orient individuals to a particular set of attributions, expectations, and goals" (56,57). Preliminary evidence indicated that improving mindsets about exercise increased its adherence (57). Moreover, booster sessions, reminders, and behavioral change techniques can improve exercise adherence by increasing motivation to partake in exercise (58,59). These strategies seem to ground their efficacy on contextual factors as well (e.g., communication with the clinicians, feeling taken care of by them, etc.). Therefore, we can argue that contextual factors and the mindsets responsible for interpreting them are worth exploring in future studies to understand their relationship with exercise adherence.

Some limitations of this study need to be discussed. First, the observational nature of the study does not allow us to establish causality and draw any definitive conclusion on the relationship between exercise adherence and the investigated factors. Second, a few variables were not reported. However, as explained in the methods section, the missingness of our data could be considered to be completely at random, primarily due to an error during the data upload process in the registers, introducing no or minimal bias in our results. However, we recommend interpreting our results cautiously, as we could not verify the reason for the data missingness. Third, our results might not be reliably applied to other forms of exercise (e.g., unsupervised home exercise) due to the specific research question of our study. Finally, physical activity hours, the number of painful joints, and living alone were found to be associated with medium but not high levels of adherence. However, this result may be influenced by chance and could also be attributed to the ad hoc adherence categorization adopted in the SOAR. Bearing in mind the limits of this study, it is worth highlighting that we reported the results of roughly 20,000 people with OA, followed by physiotherapists in the Swedish national health care system who tailored their intervention to patients' needs and characteristics. The size and data quality of our study strengthen its clinical importance and relevance for research.

To conclude, strategies based on demographic and lifestyle, socioeconomic, and disease-related factors are unlikely to improve exercise adherence significantly. Other elements, such as mindsets and contextual factors, need to be investigated. Moreover, as booster sessions, reminders, and behavioral-change techniques seem to improve exercise adherence (58,59), we should also understand how they motivate people to partake in exercise. Considering the complexity of adherence and the types of treatments that have succeeded in improving it so far, there is a call for solutions that go beyond a one-size-fits-all approach, to accept human variability and uncertainty, and to foster tailored interventions for individuals.

AUTHOR CONTRIBUTIONS

All authors were involved in drafting the article or revising it critically for important intellectual content, and all authors approved the final version to be submitted for publication. Dr. Testa had full access to all of

the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study conception and design. Battista, Kiadaliri, Jönsson, Dell'Isola.

Acquisition of data. Battista, Jönsson, Dell'Isola.

Analysis and interpretation of data. Battista, Kiadaliri, Gustafsson, Englund, Testa, Dell'Isola.

REFERENCES

- Bannuru RR, Osani MC, Vaysbrot EE, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis Cartilage* 2019;27:1578–89.
- National Institute for Health Care Excellence. Osteoarthritis in over 16s: diagnosis and management. NICE Guideline 226. London: National Institute for Health Care Excellence; 2022.
- Zeng CY, Zhang ZR, Tang ZM, et al. Benefits and mechanisms of exercise training for knee osteoarthritis. *Front Physiol* 2021;12:794062.
- Raposo F, Ramos M, Lúcia Cruz A. Effects of exercise on knee osteoarthritis: a systematic review. *Musculoskeletal Care* 2021;19:399–435.
- Skou ST, Pedersen BK, Abbott JH, et al. Physical activity and exercise therapy benefit more than just symptoms and impairments in people with hip and knee osteoarthritis. *J Orthop Sports Phys Ther* 2018;48:439–47.
- De Groot IB, Bussmann JB, Stam HJ, et al. Actual everyday physical activity in patients with end-stage hip or knee osteoarthritis compared with healthy controls. *Osteoarthritis Cartilage* 2008;16:436–42.
- Dunlop DD, Song J, Semanik PA, et al. Physical activity levels and functional performance in the Osteoarthritis Initiative: a graded relationship. *Arthritis Rheum* 2011;63:127–36.
- World Health Organization. Adherence to long term therapies: evidence for action. Geneva: World Health Organization; 2003. URL: <https://apps.who.int/iris/bitstream/handle/10665/42682/9241545992.pdf;jsessionid=159CA36674ED7E190879E9A5521BB C3F?sequence=1>.
- Pisters MF, Veenhof C, Schellevis FG, et al. Exercise adherence improving long-term patient outcome in patients with osteoarthritis of the hip and/or knee. *Arthritis Care Res (Hoboken)* 2010;62:1087–94.
- Hawker GA. Osteoarthritis is a serious disease [review]. *Clin Exp Rheumatol* 2019;37 Suppl 120:3–6.
- Safiri S, Kolahi AA, Smith E, et al. Global, regional and national burden of osteoarthritis 1990–2017: a systematic analysis of the Global Burden of Disease Study 2017. *Ann Rheum Dis* 2020;79:819–28.
- Jefferis BJ, Sartini C, Lee IM, et al. Adherence to physical activity guidelines in older adults, using objectively measured physical activity in a population-based study. *BMC Public Health* 2014;14:382.
- Hemmingsson E, Ekelund U. Is the association between physical activity and body mass index obesity dependent? *Int J Obes (Lond)* 2007;31:663–8.
- Marmot M. Social determinants of health inequalities. *Lancet* 2005;365:1099–104.
- Austin S, Qu H, Shewchuk RM. Association between adherence to physical activity guidelines and health-related quality of life among individuals with physician-diagnosed arthritis. *Qual Life Res* 2012;21:1347–57.
- Hunter DJ, Schofield D, Callander E. The individual and socioeconomic impact of osteoarthritis [review]. *Nat Rev Rheumatol* 2014;10:437–41.
- Taulaniemi A, Kankaanpää M, Rinne M, et al. Fear-avoidance beliefs are associated with exercise adherence: secondary analysis of a randomised controlled trial (RCT) among female healthcare workers with recurrent low back pain. *BMC Sports Sci Med Rehabil* 2020;12:28.

18. Battista S, Manoni M, Dell'Isola A, et al. Giving an account of patients' experience: a qualitative study on the care process of hip and knee osteoarthritis. *Health Expect* 2022;25:1140–56.
19. Battista S, Dell'Isola A, Manoni M, et al. Experience of the COVID-19 pandemic as lived by patients with hip and knee osteoarthritis: an Italian qualitative study. *BMJ Open* 2021;11:e053194.
20. Merlo J, Mulinari S, Wemrell M, et al. The tyranny of the averages and the indiscriminate use of risk factors in public health: the case of coronary heart disease. *SSM Popul Health* 2017;3:684–98.
21. Dell'Isola A, Jönsson T, Nero H, et al. Factors associated with the outcome of a first-line intervention for patients with hip or knee osteoarthritis or both: data from the BOA register. *Phys Ther* 2020;100:1771–81.
22. Svenska Artrosregistret. The Swedish Osteoarthritis registry. URL: https://boa.registercentrum.se/in-english/the-swedish-osteoarthritis-registry/p/By_o8GxVg.
23. Jönsson T, Eek F, Dell'Isola A, et al. The Better Management of Patients with Osteoarthritis Program: outcomes after evidence-based education and exercise delivered nationwide in Sweden. *PLoS One* 2019;14:e0222657.
24. Dell'Isola A, Jönsson T, Ranstam J, et al. Education, home exercise, and supervised exercise for people with hip and knee osteoarthritis as part of a nationwide implementation program: data from the Better Management of Patients With Osteoarthritis Registry. *Arthritis Care Res (Hoboken)* 2020;72:201–7.
25. Socialstyrelsen. Nationella riktlinjer för rörelseorganens sjukdomar: reumatoid artrit, axial spondylartrit, psoriasisartrit, artros och osteoporos. 2021. URL: <https://www.socialstyrelsen.se/globalassets/sharepoint-dokument/artikelkatalog/nationella-riktlinjer/2021-1-7137.pdf>.
26. Statistics Sweden. Longitudinal integration database for health insurance and labour market studies (LISA). URL: <https://pubmed.ncbi.nlm.nih.gov/30929112/>.
27. Bailey DL, Holden MA, Foster NE, et al. Defining adherence to therapeutic exercise for musculoskeletal pain: a systematic review. *Br J Sports Med* 2020;54:326–31.
28. Balestroni G, Bertolotti G. EuroQol-5D (EQ-5D): an instrument for measuring quality of life. *Monaldi Arch Chest Dis* 2012;78:155–9. In Italian.
29. Gustafsson K, Kvist J, Eriksson M, et al. Socioeconomic status of patients in a Swedish national self-management program for osteoarthritis compared with the general population: a descriptive observational study. *BMC Musculoskelet Disord* 2020;21:10.
30. Hawker GA, Mian S, Kendzerska T, et al. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). *Arthritis Care Res (Hoboken)* 2011;63:S240–52.
31. Bjorgul K, Novicoff WM, Saleh KJ, et al. Evaluating comorbidities in total hip and knee arthroplasty: available instruments [review]. *J Orthop Traumatol* 2010;11:203–9.
32. Lomi C. Evaluation of a Swedish version of the Arthritis Self-Efficacy Scale. *Scand J Caring Sci* 1992;6:131–8.
33. Lorig K, Chastain RL, Ung E, et al. Development and evaluation of a scale to measure perceived self-efficacy in people with arthritis. *Arthritis Rheum* 1989;32:37–44.
34. Valveny N, Gilliver S. How to interpret and report the results from multivariable analyses. *Med Writ* 2016;25:37–42.
35. Neely JG, Paniello RC, Cho Lieu JE, et al. Practical guide to understanding multivariable analyses: part A. *Otolaryngol Head Neck Surg* 2013;148:185–90.
36. Komorowski M, Marshall DC, Saliccioli JD, et al. Exploratory data analysis. In: *Secondary analysis of electronic health records*. Springer; 2016. p. 185–203.
37. Skou ST, Roos EM. Good Life with osteoArthritis in Denmark (GLA:DTM): evidence-based education and supervised neuromuscular exercise delivered by certified physiotherapists nationwide. *BMC Musculoskelet Disord* 2017;18:72.
38. Kiadaliri A, Dell'Isola A, Lohmander LS, et al. Assessing the importance of predictors of adherence to a digital self-management intervention for osteoarthritis. *J Orthop Surg Res* 2023;18:97.
39. Gommans LN, Scheltinga MR, van Sambeek MR, et al. Gender differences following supervised exercise therapy in patients with intermittent claudication. *J Vasc Surg* 2015;62:681–8.
40. Oosenbrug E, Marinho RP, Zhang J, et al. Sex differences in cardiac rehabilitation adherence: a meta-analysis [review]. *Can J Cardiol* 2016;32:1316–24.
41. McArthur D, Dumas A, Woodend K, et al. Factors influencing adherence to regular exercise in middle-aged women: a qualitative study to inform clinical practice. *BMC Womens Health* 2014;14:49.
42. Verhoef M, Love EJ, Rose SA. Women's social roles and their exercise participation. *Women Health* 1992;19:15–29.
43. Cheung C, Wyman JF, Peden-McAlpine C. Long-term yoga and aerobic/strength exercise adherence in older women with knee osteoarthritis: a mixed methods approach. *Int J Yoga Therap* 2022;32: article 4.
44. Lee HH, Dunsiger S, Connell Bohlen L, et al. Age moderates the effect of self-paced exercise on exercise adherence among overweight adults. *J Aging Health* 2020;32:154–61.
45. Baillet A, Chenail S, Polita NB, et al. Physical activity motives, barriers, and preferences in people with obesity: a systematic review. *PLoS One* 2021;16:e0253114.
46. Tuaki-Wosornu YA, Selzer F, Losina E, et al. Predictors of exercise adherence in patients with meniscal tear and osteoarthritis. *Arch Phys Med Rehabil* 2016;97:1945–52.
47. Areerak K, Waongenngarm P, Jarwananakul P. Factors associated with exercise adherence to prevent or treat neck and low back pain: a systematic review. *Musculoskelet Sci Pract* 2021;52:102333.
48. Kahan BC, Rehal S, Cro S. Risk of selection bias in randomised trials [review]. *Trials* 2015;16:405.
49. Van Onzenoort HA, Menger FE, Neef C, et al. Participation in a clinical trial enhances adherence and persistence to treatment: a retrospective cohort study. *Hypertension* 2011;58:573–8.
50. Collado-Mateo D, Lavin-Pérez AM, Peñacoba C, et al. Key factors associated with adherence to physical exercise in patients with chronic diseases and older adults: an umbrella review. *Int J Environ Res Public Health* 2021;18:2023.
51. Unevik E, Abbott A, Fors S, et al. Sociodemographic inequalities in outcomes of a Swedish nationwide self-management program for osteoarthritis: results from 22,741 patients between years 2008–2017. *J Clin Med* 2020;9:2294.
52. Essery R, Geraghty AW, Kirby S, et al. Predictors of adherence to home-based physical therapies: a systematic review. *Disabil Rehabil* 2017;39:519–34.
53. Daniali SS, Darani FM, Eslami AA, et al. Relationship between self-efficacy and physical activity, medication adherence in chronic disease patients. *Adv Biomed Res* 2017;6:63.
54. Bachrach DG, Rapp TL, Rapp AA, et al. “Too much” self-efficacy? Understanding the curvilinear consequences of between-person self-efficacy through a moderated-mediation model of perceived proximity and employee effort. *Group Organ Manag* 2022;2022:1–38.

55. Testa M, Rossetini G. Enhance placebo, avoid nocebo: how contextual factors affect physiotherapy outcomes [review]. *Man Ther* 2016; 24:65–74.
56. Crum AJ, Salovey P, Achor S. Rethinking stress: the role of mindsets in determining the stress response. *J Pers Soc Psychol* 2013;104: 716–33.
57. Boswell MA, Evans KM, Zion SR, et al. Mindset is associated with future physical activity and management strategies in individuals with knee osteoarthritis. *Ann Phys Rehabil Med* 2022;65:101634.
58. Cinthuja P, Krishnamoorthy N, Shivapatham G. Effective interventions to improve long-term physiotherapy exercise adherence among patients with lower limb osteoarthritis: a systematic review. *BMC Musculoskelet Disord* 2022;23:1–11.
59. Willett M, Duda J, Gautrey C, et al. Effectiveness of behavioural change techniques in physiotherapy interventions to promote physical activity adherence in patients with hip and knee osteoarthritis: a systematic review protocol. *BMJ Open* 2017;7: e015833.

Study 4

Various cultural, sociodemographic, and economic elements can influence osteoarthritis (OA) care. This thesis examines physiotherapists' knowledge of and adherence to OA clinical practice guidelines, the sociodemographic and economic factors of people with OA, and their experience and beliefs about OA care. The investigated cohort of physiotherapists showed knowledge but an inconsistent implementation of OA first-line interventions. The interviewed people with OA experienced an uncertain care process and had a negative attitude towards first-line interventions. Once exploring factors associated with exercise adherence in OA, we found an association with age, sex, socioeconomic status, disease severity, and self-efficacy. However, these factors explained only a small portion of exercise adherence variability. Income inequalities were identified among the outcomes of the participants in an OA first-line intervention. Improving OA care requires bridging the gap between knowledge of and adherence to clinical practice guidelines, empowering individuals with OA through precise explanations and support, further research on exercise adherence factors, and addressing income inequalities.



SIMONE BATTISTA is a physiotherapist specialised in Rheumatic and Musculoskeletal Diseases Rehabilitation with a Master's degree in 'Health Professional Rehabilitation Sciences'. Simone Battista is interested in the rehabilitation and management of chronic rheumatic and musculoskeletal diseases and the impact that the socioeconomic, cultural, and psychological factors specific to patients and healthcare professionals can have on the care process of these diseases. His goal is to study and improve the care quality of rheumatic and musculoskeletal conditions, considering the challenges and needs of those with these diseases.



FACULTY OF
MEDICINE

University of Genova, School of Medical and
Pharmaceutical Sciences, Doctoral Dissertation

Lund University, Faculty of Medicine,
Doctoral Dissertation Series 2023:92

ISSN 1652-8220

ISBN 978-91-8021-432-2

