

Cardiovascular magnetic resonance derived pressure volume loop variables in patients with ST-elevation myocardial infarction provide physiological information beyond ejection fraction

Lav, Theodor; Nordlund, David; Jablonowski, Robert; Khoshnood, Ardavan M.; Ekelund, Ulf; Atar, Dan; Erlinge, David; Engblom, Henrik; Arheden, Håkan

2023

Document Version: Publisher's PDF, also known as Version of record

Link to publication

Citation for published version (APA):

Lav, T., Nordlund, D., Jablonowski, R., Khoshnood, A. M., Ekelund, U., Atar, D., Erlinge, D., Engblom, H., & Arheden, H. (2023). *Cardiovascular magnetic resonance derived pressure volume loop variables in patients with ST-elevation myocardial infarction provide physiological information beyond ejection fraction.* Poster session presented at Svensk förening för klinisk fysiologi - Höstmötet 2023, Västerås, Sweden.

Total number of authors:

Creative Commons License: Unspecified

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
 You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00

Download date: 04. Jul. 2025

Word maximum: 300

Word count: 251

Cardiovascular magnetic resonance derived pressure volume loop variables in

patients with ST-elevation myocardial infarction provide physiological

information beyond ejection fraction

Theodor Lava, David Nordlund MD PhDa, Robert Jablonowski MD PhDa, Ardavan Khoshnood

MD PhDb, Ulf Ekelund MD PhDc, Dan Atar MD PhDd, David Erlinge MD PhDf, Henrik

Engblom MD PhDa, Håkan Arheden MD PhDa

a. Clinical Physiology, Department of Clinical Sciences Lund, Lund University and Skane

University Hospital, Lund, Sweden

b. Emergency Medicine, Department of Clinical Sciences Malmö, Lund University, Skane

University Hospital, Malmö, Sweden

c. Emergency Medicine, Department of Clinical Sciences Lund, Lund University and

Skane University Hospital, Lund, Sweden

d. Dept. of Cardiology, Oslo University Hospital Ulleval, Oslo, Norway

e. Institute of Clinical Medicine, University of Oslo, Norway

f. Cardiology, Department of Clinical Sciences Lund, Lund University and Skane

University Hospital, Lund, Sweden

Background

A novel non-invasive method for generation of pressure volume loops (PV-loops) using brachial blood pressure and cardiovascular magnetic resonance (CMR) imaging has recently been presented and validated (1). The aim of the present study was to investigate if PV-loop variables such as stroke work, contractility and ventriculoarterial coupling could provide incremental diagnostic information beyond conventional measurements such as cardiac dimensions and ejection fraction in patients with acute myocardial infarction (MI).

Method

100 patients with ST-elevation MI and 75 healthy volunteers were included in the study and underwent a CMR examination. All patients were examined 2-6 days after MI. Non-invasive PV-loops were generated by combining volumetric CMR data and brachial sphygmomanometric pressure measurements by using a recently validated method (1). Maximal elastance (E_{max} , translated to contractility), stroke work and ventriculoarterial coupling (E_a/E_{max}) were measured from the PV-loops. Examples of the delineations and PV-loops are shown in Figure 1. Infarct size and myocardium at risk were assessed using late gadolinium enhancement and contrast-enhanced steady state free precession images, respectively.

Results

All PV-loop variables differed significantly in patients with acute myocardial infarction compared to healthy volunteers (Table 1). Furthermore, contractility, stroke work and ventriculoarterial coupling correlated to infarct size (E_{max} : r^2 =0.29, E_a / E_{max} : r^2 =0.41, stroke

work: r^2 =0.25) and myocardium at risk (E_{max} : r^2 =0.25, E_a/E_{max} : r^2 =0.36, stroke work: r^2 =0.21) as shown in Figure 2.


Conclusion

Non-invasive cardiovascular magnetic resonance derived PV-loop variables such as contractility, stroke work and ventriculoarterial coupling provide incremental diagnostic information beyond cardiac dimensions and ejection fraction early after acute myocardial infarction.

References

(1) Seemann F., Arvidsson P., Nordlund D., et al. Noninvasive Quantification of Pressure-Volume Loops From Brachial Pressure and Cardiovascular Magnetic Resonance. Circ Cardiovasc Imaging 2019;12(1). Doi: 10.1161/CIRCIMAGING.118.008493.

Figure 1. Example of PV-loops in patients with myocardial infarction.






Table 1. PV-loop variables in patients with myocardial infarction and healthy volunteers.

Variables	Myocardial infarction	Healthy volunteers	p-value
Contractility, mmHg/ml	1.34±0.48	1.50±0.41	0.024
Ventricular arterial coupling	1.27±0.61	0.73±0.17	<0.001
Stroke work, J	0.96±0.32	1.38±0.32	<0.001
EDV, ml	166.5±34.0	174.3±32.9	0.131
EF, %	48.6±10.0	61.0±5.9	<0.001

Figure 2. Stroke work, contractility and ventriculoarterial coupling versus myocardium at risk (left column) and infarct size (right column).

