



LUND UNIVERSITY

Three new epilithic Xanthoria species from the southwestern Baltic Sea region.

Kondratiuk, Serhii; Schiefelbein, Ulf; Kärnefelt, Ingvar; Hansson, Mats; Thell, Arne

2024

Document Version:

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Kondratiuk, S., Schiefelbein, U., Kärnefelt, I., Hansson, M., & Thell, A. (2024). *Three new epilithic Xanthoria species from the southwestern Baltic Sea region..* Poster session presented at Systematikdagarna 2024, Gothenburg, Sweden.

Total number of authors:

5

Creative Commons License:

Unspecified

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



LUND
UNIVERSITY

Three new epilithic *Xanthoria* species from the southwestern Baltic Sea region

SERHII Y. KONDRATYUK^{1&2}, ULF SCHIEFELBEIN³, INGVAR KÄRNEFELT⁴, MATS HANSSON² & ARNE THELL⁴

¹ M. H. KHOLODNY INSTITUTE OF BOTANY, TERESHCHENKIVSKA STR. 2, 01004 KYIV, UKRAINE

² MOLECULAR CELL BIOLOGY UNIT, LUND UNIVERSITY, SÖLVEGATAN 35A, 22362 LUND, SWEDEN

³ UNIVERSITY OF ROSTOCK, BOTANICAL GARDEN, SCHWAANSCH STRASS 2, 18055 ROSTOCK, GERMANY

⁴ BIOLOGICAL MUSEUM, LUND UNIVERSITY BOX 117, SE-22100 LUND, SWEDEN

Abstract

Three species new to science in the genus *Xanthoria* (Teloschistaceae), *X. pedersenii*, *X. pylyporlykii* and *X. wennergrenii*, from the southwestern Baltic Sea region are presented and compared with related species.

Introduction

Xanthoria (Teloschistaceae) is a genus of lichenized fungi with a worldwide distribution. Before the molecular era, the genus included c. 50 species (Kärnefelt, 1989), but today only 13 species remain in *Xanthoria* in the strict sense (Kondratyuk et al., 2022a). Three or four species of this genus were hitherto reported from the southwestern Baltic Sea region.

Material & Methods

A method of collecting epilithic lichens thalli without damaging the rock surface with water spray pretreatment was developed. Approximately 2500 specimens were collected at 110 localities (Fig. 1–3). At least 50 ascospores outside of apothecium sections in separate specimens were measured. Conclusions about new species were based on correlation of morphological, anatomical and molecular (nrITS phylogeny) data. The specimens were studied and determined microscopically at the unit of Molecular Cell Biology, Department of Biology, Lund University.




Fig. 1. Investigated sites with epilithic *Xanthoria* species in the southwestern Baltic Sea region




Fig. 2. Investigated site: Tofta church, Skåne, Sweden. 02.04.2023



Fig. 3. Tiles of cemetery wall with *Xanthoria pylyporlykii*. Tofta church, Skåne, Sweden.

Results & Discussion

Molecular data for specimens of *Xanthoria ectaneoides*, identical with Nylander's type specimen, is presented. The species is positioned in a separate subclade – not in the *Xanthoria calcicola* subclade (Fig. 4). *Xanthoria aureola* and *X. calcicola* are found to be rather rare species in the investigated region, while *X. ectaneoides* and *X. pylyporlykii* are more common. Three species new to science, *Xanthoria pedersenii*, *X. pylyporlykii* and *X. wennergrenii* are described and illustrated.



Fig. 5 *Xanthoria pedersenii*.



Fig. 6 *Xanthoria pylyporlykii*.



Fig. 7 *Xanthoria wennergrenii*.

Xanthoria pedersenii (ined. Kondratyuk et al. 2024b) (Fig. 5). This species is similar to *Xanthoria calcicola*, but differs in having thin, paper-like thalline lobes, in having smaller knob-like warts, which may resemble coarse isidia, which never fall off, in having much narrower ascospores and much wider range of variation of ascospore septa, and in having slightly shorter but wider conidia.

Etymology: This species is named after **Christiern Pedersen** (ca 1480–16 January 1554), a Danish canon, humanist, writer, printer and publisher, who is best known for the new edition of Saxo's chronicle which thus saved the early Danish history for posterity and the translation of the Bible into Danish, known as 'Christian III's Bible'. His role in Denmark can be compared with the humanist and theologian Erasmus Rotterdam in the Netherlands.




Fig. 4. Consensus MP tree based on nrITS of the members of the genus *Xanthoria*. *data are submitted to GenBank as *Xanthoria parietina*, **as *Xanthoria* sp., and ***as *Xanthoria ectaneoides*.

Xanthoria pylyporlykii S.Y.Kondr., Kärnefelt & A.Thell in Kondratyuk et al. (2024a) (Fig. 6). The species is similar to *Xanthoria ectaneoides*, but differs in having wider thalline lobes, well-developed overlapping and irregularly orientated in the centre of the thallus, shorter ascospores and narrower ascospore septa.

Etymology: It is named after **Pylyp Orlyk** (21 Oct 1672–26 May 1742), in-exile Hetman of Ukraine and the author of the Constitution of Pylyp Orlyk (the first constitution of Ukraine). In 1716–1719 Pylyp Orlyk lived in Kristianstad with his family after an official invitation from the Swedish king Karl XII. The Latin-language original of the Constitution of Pylyp Orlyk with his signature is kept in the National Archives of Sweden.

Xanthoria wennergrenii (ined. Kondratyuk et al. 2024b) (Fig. 7). The species is similar to *Xanthoria calcicola*, but differs in having smaller thalline lobes, well developed in the peripheral zone only, pustule-like isidia and a bulky centre of the thallus, longer ascospores and wider septa.

Etymology: The new species is named after **Axel Wenner-Gren** (originally Wennergren, 5 June 1881 – 24 November 1961), a Swedish industrialist and philanthropist, who established foundations to support research including the fellowship received by the first author for this taxonomic revision of the *Xanthoria calcicola* group.

References

Kondratyuk et al. (2024a) Acta Botanica Hungarica 66(1–2): 47–77.

Kondratyuk et al. (2024b) Acta Botanica Hungarica 66(3–4): (accepted).

ACKNOWLEDGEMENTS

This work was supported by the Ove Almborn Foundation for AT, the Wenner-Gren Foundations and Erasmus+ for SK.

