



LUND UNIVERSITY

Kissing knees - factors behind the attraction. Knee abduction in individuals with anterior cruciate ligament injury.

Cronström, Anna

2017

Document Version:

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Cronström, A. (2017). *Kissing knees - factors behind the attraction. Knee abduction in individuals with anterior cruciate ligament injury*. [Doctoral Thesis (compilation)]. Lund University: Faculty of Medicine.

Total number of authors:

1

Creative Commons License:

Unspecified

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal



Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



Kissing knees – factors behind the attraction

Knee abduction in individuals with anterior cruciate ligament injury

ANNA CRONSTRÖM

MUSCULOSKELETAL FUNCTION | DEPARTMENT OF HEALTH SCIENCES | LUND UNIVERSITY





Individuals with anterior cruciate ligament (ACL) injury often exhibit poor movement quality, in the form of increased knee abduction during activity, which may affect function and increase the risk of re-injury. The results from this thesis indicate that sensorimotor function, such as proprioception and muscle activation patterns should be considered when designing training regimes, aiming at reducing knee abduction to possibly prevent further injuries in this population. Given that healthy women perform weight-bearing activities with increased knee abduction compared to men, there may need to be a greater focus on this movement pattern during prevention and rehabilitation training in women, whereas other factors may be more relevant in men.



Kissing knees – factors behind the attraction

Knee abduction in individuals with anterior cruciate ligament injury

Anna Cronström



LUND
UNIVERSITY


DOCTORAL DISSERTATION

by due permission of the Faculty of Medicine, Lund University, Sweden.
To be defended at H01, HSC, baravägen 3, Lund on 10 February 2017 at 9.00 am

Faculty opponent
Professor Joanna Kvist

Organization LUND UNIVERSITY, Sweden Department of Health Sciences Research group, Musculoskeletal function Author Anna Cronström	Document name DOCTORAL DISSERTATION	
	Date of issue February 10, 2017	
Title and subtitle Kissing knees – factors behind the attraction. Knee abduction in individuals with anterior cruciate ligament injury		
Abstract Anterior cruciate ligament (ACL) injury and patellofemoral pain (PFP) are common sports-related knee injuries. Their consequences include compromised health of the effected individual and substantial financial costs for society. Increased knee abduction or a knee medial to foot position (KMFP), so called "kissing knees", during weight-bearing activities is reported to be more common in patients with ACL injury or PFP than in non-injured individuals and is also reported to be associated with greater pain and worse function in these patients. Furthermore, increased knee abduction during activity is suggested to increase the risk of sustaining both primary and subsequent ACL injuries. However, the contributing factors for this altered movement pattern are not well understood. Therefore, the primary aim of this thesis was to investigate the association between sensorimotor factors, modifiable by training, and knee abduction in patients with ACL injury. Two systematic reviews with meta-analyses were conducted (papers I and II). These included patients with ACL injury or PFP and healthy individuals and investigated the association between sex, muscle strength, muscle activation, sensory function and joint range of motion (ROM), respectively, and knee abduction. The association between sensory function (n=51, 23 women), muscle strength and muscle activation (n=29, 11 women), respectively, and a knee abduction/KMFP in patients with ACL injury were investigated in three cross-sectional studies (papers III – V). This thesis shows that non-injured women and those with PFP perform weight-bearing activities with greater knee abduction compared to their male counterparts. In healthy individuals, lower trunk lateral flexion strength, lower gluteus maximus activation, increased hip external rotation ROM and reduced ankle dorsi-flexion range of motion were associated with increased knee abduction/KMFP. In individuals with ACL injury, reduced sensory function was associated with a KMFP, and lower vastus medialis (VM) activation and iliocostalis activation on the non-injured side but higher semitendinosus (ST) activation and ST/VM ratio were associated with increased 3D knee abduction during weight-bearing activities. Hip and knee strength was not, or only weakly, associated with knee abduction in healthy individuals and in those with knee injury. The result from this thesis indicates that sensorimotor function, such as proprioception and muscle activation patterns should be considered in the rehabilitation after ACL injury. Given that healthy women perform weight-bearing activities with increased knee abduction compared to men, there may need to be a greater focus on this movement pattern during prevention and rehabilitation training in women, whereas other factors may be more relevant in men. Further investigations are needed to confirm these results and to establish the factors, both structural and neuromuscular, that contribute to knee abduction in men and women with ACL injury.		
Key words ACL injury, sensorimotor function, postural orientation, knee abduction		
Supplementary bibliographical information		Language English
ISSN 1652-8220 Lund University, Faculty of Medicine Doctoral Dissertation Series 2017:22		ISBN 978-91-7619-403-4
Recipient's notes	Number of pages 214	Price
	Security classification	

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature 

Date 2017-01-03

Kissing knees – factors behind the attraction

Knee abduction in individuals with anterior cruciate ligament injury

Anna Cronström



LUND
UNIVERSITY

Coverphoto by Anna Cronström

Copyright Anna Cronström

Faculty of Medicine
Department of Health Sciences
Musculoskeletal function

ISBN 978-91-7619-403-4
ISSN 1652-8220

Printed in Sweden by Media-Tryck, Lund University
Lund 2017



To my family

Content

Content	7
Abstract	9
Populärvetenskaplig sammanfattning	11
List of studies	13
Definitions and Abbreviations	15
Definitions	15
Abbreviations	16
Introduction	17
Anterior cruciate ligament injury	17
Treatment of an ACL injury	18
Risk of re-injury	19
Sensorimotor function and functional knee joint stability	19
Sensory function	19
Muscle strength and muscle activation	21
Postural orientation and knee abduction	22
Overall aim and goal of the thesis	25
Specific aims	25
Methods (I–V)	27
Participants (I–V)	27
Systematic reviews (I and II)	31
Sensory function (III)	32
Muscle strength (IV)	33
Electromyography (V)	35
Functional tasks included in studies III–V	36
Single-leg squat (III–V)	36
Stair descending (III)	36
Forward lunge (III)	37
Drop jump (III)	37
Single-leg hop for distance (IV and IV)	37

Assessment of knee abduction (III–V).....	39
Visual observation and scoring (III).....	39
3D movement analysis (IV and V).....	40
Statistical analysis	40
(I and II).....	40
(III–IV)	41
RESULTS.....	43
Gender differences in knee abduction during weight-bearing activity (I and III)	43
Association between sensory function and medio-lateral knee position (III)	45
Association between joint ROM and knee abduction during weight-bearing activities (II)	46
Association between muscle strength and knee abduction during weight-bearing activities (II and IV)	46
Association between average muscle activation amplitude and knee abduction during weight-bearing activities (II and V)	48
Discussion	51
Gender differences in knee abduction	51
Sensorimotor factors associated with knee abduction.....	53
Sensory function.....	53
Muscle activation.....	54
Muscle strength	55
The role of Knee abduction for ACL injury risk.....	55
Methodological considerations	57
General limitations (I-V)	57
Sensory measures	58
Muscle activation.....	58
3D motion analysis.....	59
Clinical implications	60
Conclusions	63
Acknowledgements	65
References	67
Studies I-V.....	83

Abstract

Anterior cruciate ligament (ACL) injury and patellofemoral pain (PFP) are common sports-related knee injuries. Their consequences include compromised health of the affected individual and substantial financial costs for society. Increased knee abduction or a knee medial to foot position (KMFP), so called “kissing knees”, during weight-bearing activities is reported to be more common in patients with ACL injury or PFP than in non-injured individuals and is also reported to be associated with greater pain and worse function in these patients. Furthermore, increased knee abduction during activity is suggested to increase the risk of sustaining both primary and subsequent ACL injuries. However, the contributing factors for this altered movement pattern are not well understood. Therefore, the primary aim of this thesis was to investigate the association between sensorimotor factors, modifiable by training, and knee abduction in patients with ACL injury.

Two systematic reviews with meta-analyses were conducted (papers I and II). These included patients with ACL injury or PFP and healthy individuals and investigated the association between sex, muscle strength, muscle activation, sensory function and joint range of motion (ROM), respectively, and knee abduction. The association between sensory function (n=51, 23 women), muscle strength and muscle activation (n=29, 11 women), respectively, and a knee abduction/KMFP in patients with ACL injury were investigated in three cross-sectional studies (papers III – V).

This thesis shows that non-injured women and those with PFP perform weight-bearing activities with greater knee abduction compared to their male counterparts. In healthy individuals, lower trunk lateral flexion strength, lower gluteus maximus activation, increased hip external rotation ROM and reduced ankle dorsi-flexion range of motion were associated with increased knee abduction/KMFP. In individuals with ACL injury, reduced sensory function was associated with a KMFP, and lower vastus medialis (VM) activation and iliocostalis activation on the non-injured side but higher semitendinosus (ST) activation and ST/VM ratio were associated with increased 3D knee abduction during weight-bearing activities. Hip and knee strength was not, or only weakly, associated with knee abduction in healthy individuals and in those with knee injury.

The result from this thesis indicates that sensorimotor function, such as proprioception and muscle activation patterns should be considered in the rehabilitation after ACL injury. Given that healthy women perform weight-bearing activities with increased knee abduction compared to men, there may need to be a greater focus on this movement pattern during prevention and rehabilitation training in women, whereas other factors may be more relevant in men. Further investigations are needed to confirm these results and to establish the factors, both structural and neuromuscular, that contribute to knee abduction in men and women with ACL injury.

Populärvetenskaplig sammanfattning

Främre korsbandsskada i knäet och främre knäsmärta är två vanligt förekommande idrottsrelaterade skador som medför stora konsekvenser för den drabbade individen och höga kostnader för samhället. En korsbandsskada resulterar ofta i knäinstabilitet och funktionsnedsättning med försämrad livskvalité som följd. Dessutom innebär en korsbandsskada ofta sjukskrivning och att den drabbade måste undvika fysisk aktivitet helt eller delvis under en längre tid. Omkring hälften går aldrig tillbaka till den aktivitetsnivå de hade innan skadan.

Funktionella test utgör en viktig del i bedömningen av patientens funktion. Det är främst kvantitativa tester som används i kliniken där man mäter t.ex. muskelstyrka eller hoppförmåga. Senare forskning har visat att även kvaliteten på rörelsen har betydelse. När det gäller knäet är det framförallt knäets position i förhållande till foten som utvärderas. Större knäabduktion eller en knäposition där knäet är medialt om foten under aktivitet, så kallade "kissing knees" är vanligare hos patienter med korsbandsskada eller främre knäsmärta än hos individer utan knäskada och samvarierar även med sämre funktion och mer smärta hos dessa patienter. Större knäabduktion samvarierar också med högre risk att drabbas av en korsbandsskada och högre risk för efterföljande korsbandsskada i antingen samma knä eller i andra knäet. Det är inte klarlagt vilka faktorer som ligger bakom detta rörelsemönster. Syftet med denna avhandling var därför att undersöka sambandet mellan olika sensomotoriska faktorer, som kan påverkas med träning, och knäabduktion under funktionella test hos individer med korsbandsskada.

Två systematiska litteraturgranskningar med meta-analys genomfördes (studie I och II), där vi undersökte betydelsen av kön, muskelstyrka, muskelaktivering, sensorisk funktion och rörelseomfång för knäabduktion. Dessa studier inkluderade patienter med främre korsbandsskada eller främre knäsmärta samt friska individer. I tre tvärsnittsstudier (studie III-V) undersöktes sambandet mellan sensorisk funktion (n=51, 23 kvinnor), muskelstyrka och muskelaktivering (n=29, 11 kvinnor) och knäabduktion/medial knäposition för patienter med främre korsbandsskada.

Avhandlingen visar att hos personer utan knäskada och hos de med främre knäsmärta har kvinnor större knäabduktion än män när de utför funktionella test. Hos friska individer samvarierade lägre sidobålstyrka, lägre gluteus maximus aktivering, sämre rörelseomfång i fotleden och ökat rörelseomgång i utåtrotationen

i höften med ökad knäabduktion under funktionella test. Hos individer med främre korsbandsskada var sämre sensorisk funktion relaterat till en medial knäposition mätt visuellt. Lägre vastus medialis (VM) aktivitet och iliocostalis aktivitet för den icke skadade sidan men högre semitendinosus (ST) aktivitet och ST/VM ratio samvarierade med ökad 3D knäabduktion under aktivitet. Det fanns inget samband eller endast ett svagt samband mellan styrka i höft- och knämuskler och knäabduktion för individer utan knäskada och för de med främre korsbandsskada.

Avhandlingens resultat tyder på att sensorisk funktion, som t.ex. proprioception (ledsinne) och muskelaktivering är faktorer att ta hänsyn till i rehabilitering efter korsbandsskada. Eftersom kvinnor verkar vara mer benägna att utföra rörelser med större knäabduktion än män, kan det vara viktigt att lägga mer fokus på detta rörelsemönster under skadeförebyggande träning och rehabilitering efter skada medan andra faktorer kan vara mer relevanta för män. Fortsatta studier behövs för att bekräfta dessa resultat och för att fastställa vilka faktorer, både strukturella och neuromuskulära, som har betydelse för knäabduktion hos män och kvinnor med korsbandsskada.

List of studies

I. Cronström A, Creaby MW, Nae J, Ageberg E. Gender differences in knee abduction during weight-bearing activities: A systematic review and meta-analysis. *Gait Posture*. 2016;49:315-28.

II. Cronström A, Creaby MW, Nae J, Ageberg E. Modifiable factors associated with knee abduction during weight-bearing activities: A systematic review and meta-analysis. *Sports Med*. 2016;46(11):1647-62.

III. Cronström A, Ageberg E. Association between sensory function and medio-lateral knee position during dynamic tasks in patients with anterior cruciate ligament injury. *BMC Musculoskelet Disord*. 2014;15(1):430.

IV. Cronström A, Creaby MW, Blackmore T, Nae J, Ageberg E. Is trunk and lower extremity strength associated with knee abduction angle during weight-bearing activities in patients with anterior cruciate ligament reconstruction? In manuscript.

V. Cronström A, Creaby MW, Franettovich Smith MM, Blackmore T, Nae J, Ageberg E. Alterations in trunk and lower extremity muscle activation are associated with knee abduction during weight-bearing activities in patients with anterior cruciate ligament injury. In manuscript

Definitions and Abbreviations

Definitions

Sensorimotor function

In this thesis, sensorimotor function is defined as the processing of the afferent sensory input, signal integration in the CNS and the activation of efferent motor signals that contribute to control functional joint stability, by modulation of, for example, muscle activation and force production.

Postural orientation

In this thesis defined as the ability to stabilize the body segments in relation to each other, the environment and gravity during static and dynamic tasks.

Neuro muscular control

In this thesis defined as the ability to produce controlled movement through coordinated muscle activity to maintain joint stability.

Proprioception

In this thesis defined as the afferent information arising from mechanoreceptors in the knee joint that contribute to postural control and joint stability.

Abbreviations

ACL	anterior cruciate ligament
ACLD	anterior cruciate ligament deficiency
ACLR	anterior cruciate ligament reconstruction
ADL	activity of daily living
CNS	central nervous system
EMG	electromyography
Gmax	gluteus maximus
Gmed	gluteus medius
IC	iliocostalis
JPS	joint position sense
KMFP	knee medial to foot position
MF	medial femoral condyle
MGc	medial gastrocnemius
MM	medial malleolus
MTP 1	metatarsophalangeal joint 1
MVC	maximum voluntary contraction
N	force in newton
Nm	newton meter
OA	osteoarthritis
PFJ	patellofemoral joint
ROM	range of motion
SDM	standard difference in mean
SLHD	single-leg hop for distance
SLS	single-leg squat
ST	semitendinosus
TDPM	threshold to detection of passive motion

Introduction

Anterior cruciate ligament injury

The anterior cruciate ligament (ACL) is an intraarticular ligament situated in the knee joint with the main purpose of providing mechanical stability to the knee during movements by preventing anterior tibial translation and rotational load [1, 2]. The ACL originates at the posteromedial corner of the lateral femoral condyle and then attaches to the anterior intercondylar fossa on the tibia. The ligament consists of two bundles, the anteromedial and the posterolateral bundles, with different functions on the knee joint. When the knee is in an extended position, it is mainly the posterolateral bundle that contributes to knee stability by passive tension of this part of the ligament, whereas the anteromedial bundle is lax. On the contrary, when the knee is flexed, the anteromedial bundle tautens whereas the posterolateral bundle is lax. Hence, the ACL is able to provide stability to the knee during the whole motion path [1].

Injury to the ACL is a common sports-related injury with major consequences for the individual and with individual life time costs up to 88,000 US dollars for society [3]. In Sweden, the incidence has been estimated to approximately 8.1 in 10,000 individuals [4]. Most injuries occur during sports-related non-contact episodes [5], typically around 50 milliseconds after foot contact, with the foot planted on the surface with a nearly extended knee together with increased trunk lean and increased knee abduction [6, 7]. A non-contact ACL injury has a multifactorial etiology with several potential risk factors. Extrinsic risk factors that have been suggested include dry and warm weather, playing on artificial grass, on a hard surface or indoors as well as the type of shoes used [8, 9]. Intrinsic risk factors may include female sex, increased knee joint laxity, subtalar joint pronation, reduced hip range of motion (ROM), lower extremity asymmetries, joint stiffness, increased knee abduction moment, high ground reaction force, worse core stability, lower hamstring to quadriceps ratio, female hormones, previous knee injury and heredity [7, 10–17]. There is, however, conflicting evidence regarding the role of knee abduction [14, 18–20] and lower extremity strength [12, 17, 21, 22] for ACL injury.

Consequences of an ACL injury

Common consequences after injury are functional limitations such as mechanical instability [23], impaired balance [24], reduced lower extremity strength [25], altered muscle activation patterns [26–28] and impaired proprioception [29]. However, the main symptom after ACL injury is perceived instability during weight-bearing activities, so called “giving way” [30]. Furthermore, these individuals are at an increased risk of developing tibiofemoral and patellofemoral osteoarthritis (OA) when they are only 35 to 40 years old [31, 32]. Reduced self-reported quality of life, both in the short and in the long term is also commonly reported after ACL injury [33, 34]. Also, almost 50% of these patients suffer from depressive symptoms prior to the ACL surgery which has shown to be associated with worse self-reported function one year later [35]. Previous studies report changes in the central nervous system (CNS) after knee injury, in terms of higher frontal theta power [36, 37] and lower activation in sensory cortical areas, but higher activation in motor areas and visual-motor areas [38–40] during activity. This indicates that neuroplasticity allows the brain to compensate for the loss of sensory information from the anterior cruciate ligament when this ligament is ruptured.

Treatment of an ACL injury

An ACL injury is treated with rehabilitation with or without reconstructive surgery. However, previous studies report that surgery does not improve objective or self-reported function [34, 41–43] nor does it prevent the development of early knee osteoarthritis [44] compared to rehabilitation alone. Rehabilitation after ACL injury typically includes trunk and lower extremity strength training, range of motion training and neuromuscular training including coordination, balance and proprioceptive exercises [7, 23], modification of landing techniques and lower extremity alignment during functional activities (the knee in line with the hip and ankle) [5, 7]. Despite thorough rehabilitation, it takes approximately two years for these patients to regain pre-injury function [45]. There are, however, still functional impairments present, and many patients do not return to their pre-injury activity level; this is especially true of women [46]. Only 39% to 55% of the individuals who sustain an ACL injury return to competitive sports after reconstruction and finalized rehabilitation [47, 48]. In addition to physical function such as strength and hop performance, several psychological factors including fear of re-injury, motivation and self-esteem seem to be important for the athletes’ decision whether to return to sport [47, 49–51].

Risk of re-injury

After return to sport there is also a substantial risk of subsequent injuries to the knee. The incidence of either a re-injury to the same knee or a contralateral ACL injury is reported to be approximately 20%, with higher rates in young individuals [52]. Female sex, higher body mass index, early reconstruction, early return to high level sport, decreased postural control, asymmetrical quadriceps strength and walking with lower knee flexion force seem to increase the risk of re-injury [7, 53–55]. In a recent study, Grooms et al. also proposed that neuroplastic changes and re-organization in the CNS after the primary injury may pre-dispose the individual to subsequent injuries [40].

Sensorimotor function and functional knee joint stability

Sensorimotor function involves the processing of the afferent sensory input, signal integration in CNS and the activation of efferent motor signals that contribute to control functional joint stability, by modulation of, for example, muscle activation and force production.

Functional joint stability is traditionally defined as the ability of the joints to remain stable during weight-bearing activities and perturbations, sometimes called neuromuscular control [56]. At the knee, passive joint stability is provided by several structures such as the meniscus, the geometry of the joint, the joint capsule and knee joint ligaments, whereas functional joint stability depends on loads from gravity and muscles acting on the knee joint. Components involved in functional joint stability include sensory function, range of motion, muscle strength, muscle activation and endurance, all factors that can be modified by training [57]. While interaction of all these factors is significant for joint stability, force produced by the surrounding knee muscles is suggested to be one of the most important factors contributing to functional knee joint stability [56, 58].

Sensory function

Proprioceptive information arises from mechanoreceptors situated in the muscles, joints, ligament- and cutaneous tissues. Together with the visual and vestibular system, they all contribute to the ability of defining the position of the body in space and the position of the joints in relation to each other. Proprioceptive acuity is important for the performance of daily tasks, i.e. walking or maintaining balance, and plays an essential role in joint stability [56, 59]. Mechanoreceptors such as the fast adapting Pacinian corpuscles, slow adapting Ruffini endings,

Golgi receptors and free nerve endings are situated in the ACL, collateral ligaments and joint capsule and provide the CNS with important information regarding joint angles, load and movements. Muscle spindles located in the extrafusal muscle fibres provide the nervous system with information about muscle length and contraction velocity, and contribute to the ability to distinguish joint movement (kinesthesia) and joints position sense (JPS) [56].

In animal studies, afferent signals from the receptors situated in the ACL are shown to be important for the gamma-muscle spindle regulation and control of muscle stiffness during movements by providing the muscle spindles with proprioceptive information. The muscle spindles are thereby subconsciously contributing to joint stability by preparatory muscle stiffness regulation via the ACL-reflex hamstring arc [59]. Dyhre-Poulsen et al. and Tsuda et al. verified the existence of this ACL-reflex hamstring arc by electronic stimulation of the ACL in humans [60, 61]. A delayed reflex-hamstring response was also evident in the injured leg compared to the uninjured leg and to non-injured controls in individuals suffering from ACL injury, and this delay was further related to an increased number of give-way symptoms [62]. Furthermore, reflex transmitted co-contraction of the hamstring and quadriceps muscles has been shown to be important for stabilizing the knee joint in response to a perturbation [63].

Individuals with ACL injury have an impaired proprioception sense as assessed with kinesthesia or JPS [64], which may persist even after a return to sport [65]. This deficiency has been reported to be associated with worse hop performance [66], worse balance [67], give-way episodes [68] and worse self-reported knee function [69], indicating that adequate proprioceptive acuity plays an important role for function in these patients.

Another aspect of sensory function that has been proposed to be closely connected to proprioception is vibration sense [70]. Tactile sense, like vibration sense, arises from receptors partly shared with those responsible for proprioception, such as, the Pacinian corpuscles, but also originates from other receptors, i.e. Meissner's corpuscles, Merkel's disks and Ruffini endings, situated in the deeper layer of the skin, between the muscles, and in the periosteum. Afferent information regarding vibration sense then travels along the same pathway as proprioception sense to the cerebral cortex [71]. Vibration sense is known to be impaired in patients with hip or knee OA [72, 73]. Reduced vibration sense was also reported to be related to OA severity [74] and to be associated with a higher knee joint loading during gait, an indicator of OA progression [75]. Furthermore, in a group of individuals with or at high risk of knee OA, poor vibration sense was found to be related to the occurrence of self-reported knee instability episodes [76]. In patients with meniscectomy, poorer vibration sense at the knee was found to be related to worse performance of the number of knee bends in 30 seconds [77]. Vibration sense is

poorly investigated in patients with ACL injury [69, 78]. Thorlund et al. found no difference in vibration sense between a group consisting of patients with ACL injury or meniscectomy and a healthy control group [78]. However, in a recent study from our research group, we reported an association between poorer vibration sense and worse self-reported function as assessed by the Knee Injury and Osteoarthritis Outcome Score [69], indicating some relevance of vibration sense for function in these patients.

Muscle strength and muscle activation

Muscle strength is an important factor in functional ability, and is commonly measured as peak torque. The magnitude of muscle torque is calculated as the product of the force applied and the lever arm length, which can be influenced by several different factors, i.e. muscle activation level, limb weight, movement velocity and the angle between the force vector and the lever arm. Hence, since most of these factors vary when the limb is moved by muscular activation, the muscle may produce different torque values, even though the muscle activation level is the same. Muscle activation on the other hand is a product of both the reflex transmitted signals from mechanoreceptors and the descending alpha motor neuron signals [79]. Co-activation of agonist and antagonist joint muscles is believed to contribute to functional joint stability by increasing muscle stiffness, i.e. the muscles' resistance to lengthening, and thereby the stiffness of the joint [58]. Specifically, co-contraction of the hamstring and quadriceps muscles are reported to be important for frontal plane knee stability [80]. Recent research also indicates that the level (%maximal voluntary contraction (MVC)) to which the muscle is activated is more important for knee stability than the actual force it produces [81].

Individuals suffering from ACL injury have decreased hip and knee torque, compared to a non-injured population [25], that often persists even 20 year after the injury was sustained [82]. In addition, these patients commonly exhibit altered muscle activation pattern, such as decreased voluntary quadriceps activation [26] and delayed muscle activation onset of the quadriceps, hamstrings, gluteus maximus (Gmax), gluteus medius (Gmed) and gastrocnemius muscles [27, 83, 84]. An increased average activation amplitude of the hamstring muscles [28, 85-88], Gmax [86, 88] and gastrocnemius muscles [86] during functional performance compared to non-injured individuals is also reported after ACL injury. Moreover, hamstring stiffness has been found to be greater in the ACL-injured leg compared to the uninjured leg and to healthy controls [89]. Decreased hamstring stiffness was also related to worse perceived functional performance, assessed with the Noyes rating scale, after ACL injury [90].

Given that both the hamstring and gastrocnemius muscles act as ACL agonists and thereby resist anterior tibial translation [91, 92], it has been suggested that these alterations in muscular activity are compensatory mechanisms intended to maintain neuromuscular control after the loss of the ACL [91, 92]. In patients with ACL reconstruction (ACLR), greater knee laxity has been associated with reduced hamstring voluntary activity, [93] and delayed gastrocnemius onset during perturbation [84], respectively. Taken together, this implies that appropriate activation of these muscles is important for functional knee stability in the ACL injured population.

Postural orientation and knee abduction

One aspect of sensorimotor function is postural orientation. Postural orientation is defined as the ability to stabilize the body segments in relation to each other, the environment and gravity; e.g. to keep the trunk, hip, knee and ankle aligned during dynamic activities [94]. Worse postural orientation during weight-bearing activities has been reported in individuals with ACL injury compared to the non-injured leg and to individuals without injury [95, 96]. This altered movement pattern was also associated with worse knee confidence [97], worse self-reported function [96, 98] and reduced hop performance [96] in these patients.

Postural orientation at the knee is commonly evaluated by the position of the knee in relation to the foot. A position where the knee is positioned medial to the foot is considered poor postural orientation, whereas a position where the knee is aligned over the foot is considered good postural orientation. This idea of “good and poor” movement pattern has been founded on anecdotal rather than scientific evidence. However, recent studies show that a knee medial to foot position (KMFP) or increased knee abduction is more common in individuals with ACL injury or patellofemoral pain (PFP) than in uninjured controls [94, 100–102] and is still observed 20 years after injury [81].

Increased knee abduction is also consistently reported to be part of the mechanism of non-contact ACL injury [11, 103–105], and is associated with greater strain on the ACL [106–110]. With respect to PFP, greater knee abduction has been demonstrated to increase contact forces within the lateral patellofemoral joint [111]. Thus, knee abduction has been suggested to be a risk factor in sustaining both PFP and ACL injury [112] as well as subsequent knee injuries [17, 113].

The gold standard to evaluate this movement pattern is by assessing knee abduction in degrees (frontal plane angle between the thigh and shank) with 2D or 3D movement analysis systems [98]. However, a more clinically feasible method is to evaluate medio-lateral knee position by visual observation, which is a reliable method that recently showed good validity against both 2D and 3D analysis [99].

The contributing factors for increased knee abduction are not well established. Several mechanisms have been suggested to be associated with this movement pattern in healthy individuals, such as static alignment, reduced muscle strength, altered muscle activation and decreased foot and ankle mobility [99–104]. Furthermore, women are believed to exhibit greater knee abduction during activity compared to men, which has been suggested to be a contributing factor for the increased knee injury risk observed in women [19]. However, the results of these studies are inconsistent [105–108] and studies on the corresponding factors in patients with knee injury is poorly investigated [109, 110].

Overall aim and goal of the thesis

The overall aim of this thesis was to investigate the association between sensorimotor factors and knee abduction in patients with ACL injury. The goal was to identify factors that are modifiable by training and exercise therapy. This knowledge may be helpful in designing rehabilitation protocols aimed at reducing knee abduction and preventing subsequent knee injuries in this population.

Specific aims

Specific aims were to:

1. Systematically review possible gender differences in knee abduction (I) and the association between modifiable sensorimotor factors and knee abduction (II) during weight-bearing activities in individuals with ACL injury, with PFP and in a healthy population.
2. Investigate the association between sensory function (kinesthesia and vibration sense) and knee abduction assessed by visual observation during weight-bearing activity in individuals with ACL injury and in addition, investigate any gender differences in the presence of knee abduction and/or in the contribution of sensory function for knee abduction (III).
3. Investigate the association between muscle strength (IV) and muscle activation (V), respectively, and knee abduction (3D) during weight-bearing activities in individuals with ACL injury.

Methods (I–V)

Participants (I–V)

The two first studies in this thesis were systematic reviews with meta-analysis with the intention of investigating what was already known about which factors contribute to knee abduction. Thus, in addition to individuals with ACL injury, we also included in these two reviews individuals with PFP and healthy individuals. Study I included 58 articles with in total 114 participants with ACL injury, 116 with PFP and 2,465 healthy individuals. Study II Included 33 articles with in total 67 participants with ACL injury, 35 with PFP and 1,005 healthy individuals (Figures 1 and 2). In study III, the cohort constituted a sample of convenience, including 28 men and 23 women with arthroscopic or MRI-verified ACL deficiency (ACLD), or ACLR, recruited from physiotherapy clinics in Skåne, Sweden. The results in studies IV and V were based on one cohort (18 men and 11 women with ACL reconstruction), where the participants were mainly recruited from the Department of Orthopedics, Skåne University Hospital, Sweden. All patients that had undergone an ACL reconstruction between June 1st and December 8th, 2015 (n=106) were invited to participate (Figure 3). Exclusion criteria common to studies III–V were: age <18 or >39, usage of external devices to assist with weight-bearing, no longer participating in formal rehabilitation and/or suffering from other injuries or diseases that affected function. The participants included had a mean pre-injury activity level [111] of 9 (III) and 8 (IV and V), respectively. We included participants both with and without associated injuries to the knee. In study III, 64% had associated injuries and in studies IV and V, 76% had associated injuries, with meniscus injury being the most commonly associated injury in both cohorts, see Table 1 for characteristics.

The three original studies were approved by the Advisory Committee for Research Ethics in Health Education at the Faculty of Medicine of Lund University (VEN 48–12) (III) or the Regional Ethical Review Board in Lund, Sweden (2015/581) (IV and V).




Fig 1. Flow chart of the inclusion process. KMFP knee medial to foot position (I)




Fig 2. Flow chart of the inclusion process. KMFP knee medial to foot position (II)




Fig 3. Flow chart of the inclusion process (IV and V)

Table 1. Characteristic of the participants included in study III-V

Characteristics	Study III		Study IV and V
	Women (n = 23)	Men (n = 28)	n=29 (11 women, 18 men)
Women, n (%)	NA	NA	11 (38)
Age (years)▫	23 (4.0)	26 (6.1)	27 (5.6)
BMI▫	22.5 (2.21)	24.7 (2.85)	25.9 (3.6)
Tegner activity score pre injury #	9 (8–9)	9 (8–9)	8 (7-9)
Tegner activity score at assessment #	3 (2–6)	4 (3–7)	3 (2-4)
Injured right knee, n (%)	11 (48)	20 (71)	12 (41)
Time since injury, non-reconstruction (weeks)▫	45 (42.1)	66 (57.9)	NA
Reconstruction, n (%)	19 (83)	19 (68)	29 (100)
Time since reconstruction (weeks)▫	36 (40.7)	47 (52.5)	29 (6.4)
Associated injuries, n (%)	13 (57)	20 (71)	22 (76)
Collateral ligament, n (%)	5 (22)	8 (28.6)	8 (28)
Meniscal, n (%)	9 (39)	16 (57)	19 (65)
Cartilage, n (%)	8 (35)	7 (25)	7 (24)

▫ = mean (SD), # = median (quartiles), BMI= body mass index, NA=not applicable

Systematic reviews (I and II)

Studies I and II were both systematic reviews. In the beginning these were meant to be one review, investigating modifiable factors and gender as contributing factors for knee abduction, with the purpose of guiding the content in my subsequent studies. However, after the initial search was performed, we realized that the number of studies and amount of data available were much more substantial than we had initially believed and, therefore, we decided to divide the results into two reviews: (I) gender differences in knee abduction, and (II) modifiable factors associated with knee abduction. These systematic reviews were conducted according to the PRISMA guidelines and the protocol was pre-registered in the International Prospective Register of Systematic reviews (PROSPERO) (2013: CRD42013005415). The initial systematic search conducted in August 2013 included the databases Medline (PubMed), CINAHL and EMBASE. The same search strategy was applied to both reviews, including the following terms;

(healthy[Title/Abstract]) OR non-injured[Title/Abstract]) OR ACL reconstruct*[Title/Abstract]) OR ACL injur*[Title/ Abstract]) OR anterior cruciate ligament injur*[Title/Abstract]) OR anterior cruciate ligament reconstruct*[Title/Abstract]) OR patellofemoral pain[Title/Abstract]) OR “Anterior Cruciate Ligament”[Mesh]) OR “Anterior Cruciate Ligament Reconstruction”[Mesh]) OR “Patellofemoral Pain Syndrome”[Mesh])) OR uninjured[Title/ Abstract])) AND (“joint position sense”[Title/Abstract]) OR propriocept*[Title/Abstract]) OR kinesthesia[Title/Abstract]) OR proprioception[MeSH Terms]) OR Muscle activation[Title/Abstract]) OR Muscle strength[Title/Abstract]) OR muscle strength[MeSH Terms]) OR muscle strength dynamometer [MeSH Terms]) OR range of motion[Title/Abstract]) OR ROM[Title/ Abstract]) OR range of motion, articular[MeSH Terms]) OR gender [Title/Abstract]) OR sex[Title/Abstract])OR females[Title/Abstract]) OR males[Title/Abstract]) OR sex characteristic[MeSH Terms])OR“Sex Factors”[Mesh]))AND (biomechanic*[Title/ Abstract]) OR kinematic*[Title/Abstract]) OR valgus[Title/Abstract]) OR “postural orientation”[Title/Abstract]) OR alignment [Title/Abstract]) OR “movement quality”[Title/Abstract]). This search was then updated in September 2015.

We intended to include all original studies that investigated any of the following factors: gender, muscle strength, muscle activation, proprioception and/or joint range of motion (ROM) as a contributing factor in knee abduction (assessed by visual observation or 2D or 3D motion analysis) during any kind of weight-bearing activity in patients with ACL injury or PFP or in healthy individuals. Two reviewers (AC and JN) independently screened abstracts and full-text articles against the inclusion-exclusion criteria. The same reviewers then assessed all articles that met the inclusion criteria for methodological quality by using a checklist previously developed for quality assessment of observational studies [112]. A quality score of at least 50% was required to be included in the reviews, as the result in low quality studies may be affected by bias due to poor methodology.

Sensory function (III)

The aim of study III was to investigate the importance of adequate sensory function, as measured by kinesthesia and vibration sense, for the ability to control frontal plane knee movement during weight-bearing activities. We chose kinesthesia over JPS, since kinesthesia has been proven to be a more reliable and valid test of proprioception, with lower measurement errors [113]. Kinesthesia was measured by the threshold of detection of passive motion (TDPM) with a device specially built for this purpose. For a full description of the assessment procedure,

see Ageberg et al., [114]. The device consists of a platform mounted on a steel frame with an electrical motor at one end. The subject lies on the platform in a lateral position with their shank in a plaster brace. The brace is connected to the electrical motor with a wire and, by pushing a button on a remote control, the tester can make the brace move the knee in either flexion or extension at an angular speed of 1 degree per second. An analogue scale at the end of the platform registers movements in increments of 0.25°. In order to prevent any false experience of movements derived from the sound when the measuring device was started, all participants wore headphones and listened to a recording of the sound that was produced by the electric motor. The subjects were told to close their eyes and indicate by raising their hand when any movement in the knee was felt. TDPM was measured towards extension and flexion from a 20-degree (knee flexion) starting position. This starting position was chosen because with this device higher reliability and sensitivity have been reported closer to the end point of range of motion compared to the middle part of range of motion [114]. The median value from three measurements of the extension and flexion assessments respectively were determined, and the sum of these measures was then used in the statistical analysis. A higher value indicates poorer TDPM.

To gain a broader understanding of sensory function in patients with ACL injury, we also included vibration sense as another measure of sensory function. Vibration sense is assessed by the vibration perception threshold (VPT) with a biothesiometer, which is a reliable [72] and commonly used measure of sensory function in patients with OA [72, 73] or injury to the lower extremity [78, 115, 116]. During the assessments, the participant was in a supine position on a treatment bench. Before the trials, the biothesiometer was tested on the processus styloideus ulna in order to familiarize the participant with the procedure. The head of the biothesiometer was then held to the participants' most prominent point of the metatarsophalangeal joint 1 (MTP1), the medial malleolus (MM), and the medial femoral condyle (MF) with a standard pressure. These locations were chosen due to the results of previous studies where generalized deficiencies in vibration sense but better vibration sense at more distal body parts compared to more proximal body parts in patients with OA were reported [72, 73]. The participants were told to close their eyes and indicate by raising their hand when any sense of vibration was felt. The amplitude was increased by 1 volt per second, and the voltage when the subject first felt any sensation of vibration was noted as the VPT. A higher value indicated worse VPT.

Muscle strength (IV)

The aim of study IV was to investigate the relation between trunk and lower extremity strength and knee abduction. Therefore, we assessed the isometric peak

force (N) of the hip external rotators, hip abductors, hip extensors, knee flexors, knee extensors, trunk extensors, and trunk lateral flexors (side-bridge test) with a hand-held dynamometer (Commander Echo, JTECH Medical, Salt Lake City, Utah, USA) according to previously described methods [100, 117–121]. During all assessments, a belt was used to fixate the dynamometer, and the participant was encouraged to push their leg/trunk against the dynamometer as forcefully as they could, see Figure 4 a-g and [122] for a full description of testing positions. Prior to the strength assessments, the distance between the dynamometer location and the rotation axis of the joint for each muscle group was assessed with a tape measure and noted as the lever arm. The peak value of three trials was then calculated in Newton metres (Nm) by multiplying the peak force value with the corresponding lever arm for all tests except for the side-bridge test, which was multiplied with the distance between acromion and the lateral malleolus. These values were then normalized to body mass (Nm/kg). The participant held each contraction for 5 seconds with 15 seconds of rest in between. Before the data included in this study were collected, a pilot study including 9 healthy individuals was performed. The reasons for this pilot study were partly for me to learn the method properly, but also to determine test-retest reliability for all strength measures. All strength assessments showed moderate to excellent test-retest reliability; hip external rotators (ICC=0.88), hip abductors (ICC=0.77), hip extensors (ICC=0.80), knee flexors (ICC=0.95), knee extensors (ICC=0.80), trunk extensors (ICC=0.73) and trunk lateral flexors (ICC=0.62). I then collected all the strength measure data in the original study.




Figure 4 a-g. Strength assessment positions (IV)

Electromyography (V)

In study V, I aimed to investigate the possible association between muscle activation of the trunk and lower extremity muscles and knee abduction. Average muscle activation patterns of the Gmax, gluteus medius (Gmed), semitendinosus (ST), vastus medialis (VM) medial gastrocnemius (MGc) and Iliocostalis (IC) muscles for the ACLR leg (both sides for IC) were, therefore, synchronously collected with the kinematic data. For this purpose, surface electromyography (EMG) electrodes with a wireless EMG system (Desktop DTS, Noraxon U.S.A. Inc, Scottsdale, Arizona, USA) with a sampling frequency of 1500Hz and a lowpass filter of 500Hz was used. Prior to the placement of the electrodes, the skin was shaved and lightly abraded with a medical abrasion gel (Nuprep, Weaver and company, Aurora, Colorado, USA). Disposable self-adhesive dual EMG electrodes (Noraxon, USA. Inc, Scottsdale, Arizona, USA) with an inter-electrode distance of 17.5 mm were used. The electrodes were placed parallel to the muscle fibres for all muscles (except for VM which was placed perpendicular to the line between the landmarks) and at locations according to the Surface Electromyography for Non-Invasive Assessment of Muscles (SENIAM) guidelines [123].

MVC data for each muscle was collected synchronously with the muscle strength data used in study IV. The participants were encouraged to push their leg/trunk against a manual resistance as much as they could and hold for 5 seconds. The MVCs were calculated from the maximum value of three repetitions with 15 seconds of rest in between.

All EMG data was collected during the flexion phase for the single-leg squat (SLS) and the single-leg hop for distance (SLHD), respectively. The raw EMG data were then ECG-reduced, high-pass filtered at 20Hz, full-wave rectified and smoothed by a root mean square algorithm over 50ms windows (MyoRESEARCH Biomechanical Analysis, Noraxon, version 3.6). All processed EMG signals for each muscle were then normalized to the maximum value from the respective MVC. Collection of corresponding EMG data was also included in the pilot study on 9 healthy individuals, mentioned in the method section of the muscle strength assessment, in order for me to become familiarized with the method before the original data collection. Moderate to excellent reliability has been reported for surface EMG measurements of the lower extremity and trunk during weight-bearing activities [124].

Functional tasks included in studies III–V

In this thesis, we wanted to investigate contributory factors for increased knee abduction during weight-bearing functional activities with different degrees of difficulty. Thus, in study III we included functional tasks commonly used for assessing postural orientation in the clinical setting, reflecting both activities of daily living (ADL) (SLS, stair descending) and more strenuous sport-related activities (forward lunge, drop jump). However, in the consecutive studies, we only included one task reflecting ADL activity (SLS) and one task reflecting sporting activity (SLHD). In studies IV and V, we chose to replace the drop jump task with the SLHD, since the drop jump has recently been reported to be a poor screening task for predicting future knee injuries [18] and for discriminating between men and women [125]. The participants wore shorts, sports bra (women) and their own personal trainers (when needed) during the tasks. The ADL activities were repeated 5 times, whereas the sporting activities were repeated only 3 times due to the risk of fatigue in these tasks. Up to 3 practice trials were allowed to familiarize participants with the tests.

Single-leg squat (III–V)

The single-leg squat (SLS) was performed according to Ageberg et al. [126], but without allowing fingertip support for maintaining balance. The participant was standing on one leg with the second toe placed on a marked longitudinal line with the contralateral leg lifted from the floor. They were then instructed to flex the injured knee until they could not see their toes (approximately 50°) (study III), or until they touched a treatment bench placed behind them with their bottom (60°) (studies IV and V), and then return to extension. The squat was repeated five times at a speed of 3 seconds per squat. The participant performed the task barefoot (study III) or with shoes (studies IV and V) (Figure 5a).

Stair descending (III)

This task was performed according to Pfeifer et al [127], but modified to use a step board 27 centimetres high. The participant stood with both feet (hip-width apart) on the step board. They were then instructed to step down onto the floor with their non-injured leg and then return to the starting position. The injured leg, which is in contact with the step board throughout the entire movement, was evaluated. The task was repeated 5 times. The participant performed the task barefoot.

Forward lunge (III)


The forward lunge was performed according to Alkjaer et al [128]. The participant was standing with their feet hip-width apart on the floor. They were then instructed to take a long stride forward, about 1 m, with the injured leg and to flex the knee to approximately 90°, and then push back to the starting position by extending the front leg. This task was repeated 3 times. The participant performed the task with shoes.

Drop jump (III)

The drop jump was performed according to Hewett et al., [19] but modified to use the second landing in the analysis instead of the first landing that is more commonly used. This approach was chosen because the second landing may better represent a situation when an ACL injury occurs [129]. The participant stood on a step board (27 cm high) with their feet hip-width apart. They were then instructed to drop from the step board and directly perform a maximal vertical jump. Arm swing was allowed during the jump. The task was repeated 3 times. The participant performed the task with shoes.

Single-leg hop for distance (IV and IV)

Performed according to Zätterström et al.[130] to allow arm swing during the jump to enable a more functional execution of the task. The participant stood on their ACLR leg with the toes behind a marked line, and with the other leg lifted from the floor by flexing the knee. The participant was then told to jump forwards as far as possible, taking off and landing on the same foot, and to maintain balance on landing for 3–5 seconds. The distance in centimetres from toe at take-off to heel in landing was then measured. If there was more than 30 cm between the longest and shortest jump, additional jumps were performed until less than 30 cm was achieved. This approach was chosen so as to reduce the risk of too many repeated jumps on the individual's injured leg. The task was repeated 3 times. The participant performed the task with shoes (Figure 5 b).



Figures 5a-b. The single-leg squat (a) and the single-leg hop for distance (b) (IV and V)

Assessment of knee abduction (III–V)

Visual observation and scoring (III)

In study III, we used visual observation and scoring of the position of the knee in relation to the foot as a measure of knee abduction. Visual observation is an easy of handling and a clinically feasible way of assessing postural orientation of the knee, and has been reported to be valid in both 2D and 3D [131]. Two experienced raters independently scored the position of the knee in relation to the foot on video recordings of each trial. The raters were able to watch the video in slow motion and to review the movie as many times as needed. The knee position relative to the foot during the participant's performance of each task was assessed on an ordinal scale from 0 to 2. If the mid-point of the patella was in line with or lateral to the second toe, a score of 0 = “good” was given for the postural orientation. If the mid-point of the patella was medial to the second toe, a score of 1 = “fair” was given. Finally, if the mid-point of the patella was clearly placed medial to the first toe, a score of 2 = “poor” was given (Figure 6). A consensus score from the two raters was then used in the analysis. These assessments of postural orientation showed good to excellent agreement (ICC = 0.710–0.939).




Figure 6 a-c. The position of the knee in relation to the foot a) A score of 0, b) A score of 1, c) A score of 2 (III)

3D movement analysis (IV and V)

In studies IV and V, we used three-dimensional (3D) motion analysis equipment for assessing knee abduction. A combination of individual reflective markers and marker clusters were attached to the trunk and the participants' injured leg so as to track their kinematics. An eight-camera motion analysis system (Qualisys motion capture system, Gothenburg, Sweden) (150 Hz) was then used for collection of 3D knee abduction data during the flexion phase of the SLS and SLHD. For the SLS, the flexion phase was defined as the time frame from when the knee flexion angle increased by more than 2 degrees from full extension at the commencement of the squat to when the knee flexion angle reached 2 degrees less than the angle achieved at full flexion (the bottom of the squat). An adaptation of the methods of Fellin et al. [132] was used to define foot contact in the SLHD as the time at which lowest height of the distal heel or toe marker occurred (whichever occurred first). The flexion phase for the SLHD was defined as the time frame from foot contact to when the knee flexion angle reached 2 degrees less than the peak knee flexion angle achieved during the landing. From these data, peak knee abduction angle and knee abduction excursion (the change in angle from foot contact to peak knee abduction) during the flexion phase of the two tasks were calculated.

The 3D data was reconstructed and labelled in Qualisys Track Manager (version 2.12). All further processing was carried out in Visual 3-D (version 5.02, C-motion, Germantown, MD, USA). Marker trajectories were filtered with a 12Hz, 4th order, low pass Butterworth filter [133]. Knee kinematics (flexion-extension, abduction-adduction and internal-external rotation) in all movement trials were calculated using a joint coordinate system approach [134]. The mean value of the five squats, and the three jumps, respectively, were then included in the analyses

Statistical analysis

(I and II)

In studies I and II, the results were based on meta-analyses. Comprehensive Meta-Analysis software (version 2.2.064, Biostat, Englewood, USA) was used for this purpose. The effect size with 95% CI was calculated on standardized mean difference (SDM) in knee abduction between men and women (I) and as Pearson's correlation coefficient for the association between the assessed factor and knee abduction (II). If appropriate data was not provided, the relevant effect size was calculated from the p-value and sample size [135]. When only group comparison studies were included in the analysis, the effect size was calculated on SDM in

knee abduction between the two groups also in study II. If studies reported data for more than one leg (I and II) or more than one strength measure (II), right and dominant legs and the strength measure (isometric, eccentric, concentric) represented in most studies in that specific analysis were included. To deal with the problem of including multiple interventions, so called double counting, from one study in meta-analyses, we chose to divide the sample size with the number of tasks to create smaller comparator groups if a study reported data for more than one functional task. Each task was then treated as an independent study. This is one of a few approaches suggested by Cochrane [136] to resolve this problem. A random effect model was used in both reviews (I and II) due to expected heterogeneity between studies, such as the use of different functional tasks and different outcomes, i.e. visual observation, 2D or 3D analysis. Between-studies, heterogeneity in effect size was calculated with the Q-test and expressed as I²-statistics. Where possible, subgroup analyses for specific tasks or gender were performed. Meta-analyses were performed separately for healthy individuals, individuals with ACL injury and individuals with PFP. Since there may be methodological differences in assessing knee abduction with motion analysis or by visual observation, separate analysis was conducted also for all data assessed with motion analysis equipment and visual observation. Publication bias was evaluated using Funnel plots with trim and fill imputations [137].

(III–IV)

The calculations for studies III–V were performed with SPSS (version 20, IBM Corporation, New York, USA) (I) and version 21 (IV and V). All data was checked for normality by visual inspection of histogram, Q-Q plots and the Kolmogorov-Smirnov test (III–V). In studies IV and V, the residuals were also checked for normality. Pearson's correlation coefficient, Spearman's rank correlation coefficient, and the independent T-test were used as appropriate to investigate any possible association between participant demographics and knee abduction in all three studies. Since postural orientation was visually scored on an ordinal scale in study III, Spearman's rank correlation coefficient was used to determine the association between the sensory measures and knee position. In this study, the knee position scores were also dichotomized for each test; one group included subjects who were given a score of "0" (good), and one group included subjects who had been given a score of either "1" or "2" (poor). These values were then used to evaluate any differences in the sensory measures between subjects with good and poor postural orientation using the independent t-test. In studies IV and V, Pearson's correlation coefficient (normally distributed data) or Spearman's rank correlation coefficient (data not normally distributed) were used for evaluating the association between each muscle torque and each muscle activation

variable, respectively, and knee abduction. Each statistically significant factor (IV) or all variables that correlated with knee abduction ($r/rs\text{-value} \geq 0.3$, $p\text{-value} \leq 0.1$) (V) were then included in separate linear regression models (normally distributed data) or in a partial correlation model (data not normally distributed), adjusting for sex and BMI to evaluate the contribution of each measure for the dependent knee abduction variables (peak and excursion). To investigate any possible gender differences in the association between sensory function and knee abduction, men and women were analysed separately in study III. In this study, the Mann–Whitney U-test was also used to detect any differences in knee abduction between men and women.

RESULTS

Gender differences in knee abduction during weight-bearing activity (I and III)

The meta-analyses in study I showed that women with PFP exhibit greater peak knee abduction during weight-bearing activities (SDM; -1.34, 95%CI; -1.83 to -0.84) compared to men with PFP. Non-injured women were found to perform the tasks with increased knee abduction during the entire movement (initial contact, peak abduction, excursion) (SDM; -0.68 to -0.79, 95%CI; -1.04 to -0.37) compared to their male counterparts (Figure 7). Most tasks discriminated between men and women, e.g., walking, running, jump landings and cutting, whereas vertical drop jump did not.

We found no gender difference in knee position score, assessed by visual observation in individuals with ACLD or ACLR during the SLS, stair descending, forward lunge or drop jump ($p>0.116$) (study III).




Figure 7. Gender difference in peak knee abduction during functional tasks in healthy individuals (I).

Association between sensory function and medio-lateral knee position (III)

Sixteen of the 51 subjects were not able to perform all tasks (due to being too close to surgery/injury or fear of performing the jumping tasks), leaving between 13 and 28 individuals in the gender-stratified analysis. We found that worse kinesthesia was associated with a KMFP during the drop jump in both men ($r_s = 0.423$, $p = 0.044$) and women ($r_s = 0.469$, $p = 0.106$). In women, worse vibration sense at MTP1 ($r_s = 0.453$, $p = 0.034$) and MM ($r_s = 0.626$, $p = 0.002$) was associated with a KMFP during stair descending. Worse vibration sense at MM was also associated with a KMFP during the forward lunge ($r_s = 0.544$, $p = 0.016$). Consistent with the correlation analyses, there were significant differences in sensory function between those with good and poor postural orientation, i.e., women with a KMFP during stair descending had worse vibration sense at MTP 1 and MM and women with a KMFP during forward lunge had worse vibration sense at the MM compared to those with a knee over foot position (KOFP) (Figure 8).




Figure 8. Differences in vibration sense between women (ACL/ACL) with good and poor postural orientation during stair descending (a) and the forward lunge (b) (III). VPT=vibration perception threshold, MM=medial malleolus

Association between joint ROM and knee abduction during weight-bearing activities (II)

The meta-analyses on healthy individuals in study II showed that reduced passive ankle dorsi-flexion ROM (knee extended) (SDM: -0.40; 95 % CI -0.85 to 0.05) but increased hip external rotation ROM (SDM: 0.50; 95 % CI 0.09 to 0.91) were associated with a KMFP during bilateral squatting in a group consisting of both men and women. No relation was found between ankle dorsi-flexion with the knee flexed, hip internal rotation or hip abduction ROM and knee abduction.

Association between muscle strength and knee abduction during weight-bearing activities (II and IV)

In study II, the meta-analyses showed no or only weak associations between lower extremity strength and knee abduction in healthy individuals when tasks were pooled ($r \leq 0.21$, $p = 0.013-0.426$) (Figure 9). However, in a sub group analysis of the single-leg squat, lower hip external rotator torque was associated with greater peak knee abduction ($r -0.38$; 95 % CI -0.53 to -0.21). No associations were found between hip abductor, hip extension or hip external rotation strength and knee abduction during the tasks in a subgroups analysis consisting of women only ($r \leq -0.25$; 95% CI -0.48 to 0.16). However, lower trunk lateral flexion torque, assessed with the side-bridge test, was associated with greater peak knee abduction during a SLS ($r -0.39$; 95 % CI -0.54 to -0.22) in a group consisting of both men and women.

In study IV, lower BMI was associated with both greater strength and greater knee abduction and was thus included as a covariate together with sex in the regression models. Consistent with the results in study II, we found mainly no associations between lower extremity peak torque and knee abduction during the SLS or the SLHD in individuals with ACLR after adjusting for sex and BMI. However, higher knee flexion peak torque was associated with increased knee abduction excursion during the SLHD. In this study including individuals with ACLR, we found no relation between the side-bridge test and knee abduction during the two tasks (Table 2).




Figure 9. Association between hip abductor, hip extensor and hip external rotator strength and peak knee abduction during functional tasks in healthy individuals (II).

Association between average muscle activation amplitude and knee abduction during weight-bearing activities (II and V)

In study II, we found that lower Gmax activation was associated with greater 3D peak knee abduction during functional tasks in healthy females (r -0.34; 95 % CI -0.51 to -0.15), whereas there was no relation between Gmed or MGc and knee abduction ($\text{SDM} \leq 0.26$; 95 % CI -0.87 to 1.39).

In individuals with ACLR (V), higher ST activation was associated with greater knee abduction excursion during the SLS after adjusting for sex and BMI in the regression models. Lower VM activation and lower IC activation on the non-injured side were associated with greater peak knee abduction during the SLHD (Table 2). The partial correlations showed that greater ST/VM ratio was associated with greater peak knee abduction during the SLS and greater peak knee abduction and excursion during the SLHD ($r_{\text{sp}} = 0.415 - 0.583$, $p \leq 0.012$).

Table 2. Linear regression (β) of the contribution of each muscle peak torque and each muscle activation variable, adjusted for sex and BMI for the knee abduction variables (peak, excursion) during the single-leg squat (SLS) and single-leg hop for distance (SLHD) (IV and V).

		SLS										SLHD									
		Knee abduction (degrees)										Knee abduction (degrees)									
		Peak					Excursion					Peak					Excursion				
Peak torque (Nm/kg)		β	SE	95% CI	p-value		β	SE	95% CI	p-value		β	SE	95% CI	p-value		β	SE	95% CI	p-value	
Hip abduction		-	-	-	-	-	4.62	3.91	-3.43;12.67	0.249	-	-	-	-	-	-	-	-	-	-	-
Knee flexion		6.78	4.43	-2.34;15.9	0.138	6.26	3.32	6.06	-0.58;13.09	0.071	8.66	6.06	6.06	-3.84;21.14	0.166	7.85	3.59	3.59	0.449;15.25	0.039	
Trunk extension		4.38	3.31	-2.43;11.20	0.197	-	-	4.51	-3.43;15.16	0.206	-	5.86	4.51	-	-	-	-	-	-	-	-
Side-bridge		0.80	0.52	-0.27;1.86	0.137	-	-	1.17	-0.28;2.61	0.109	-	-	-	-	-	-	-	-	-	-	-
Average muscle activation (% MVC)																					
Gmed		-0.11	0.07	-0.25;0.03	0.104	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
ST		0.26	0.13	-0.01;0.52	0.055	0.20	0.10	0.00;0.40	0.048	-	-	-	-	-	-	-	-	-	-	-	-
VM		-	-	-	-	-	-	-	-	-	-	-0.10	0.02	-0.13;-0.03	0.004	-	-	-	-	-	-
IC injured		-0.39	0.22	-0.84;5.06	0.080	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
IC non-injured		-	-	-	-	-	-	-	-	-	-	-0.07	0.03	-0.13;-0.01	0.028	-	-	-	-	-	-

SE=standard error, CI=confidence interval, - indicate variables not included in the regression model, bold characters indicate significant correlation ($p<0.05$).

Discussion

The aim of this thesis was to determine which factors contribute to altered movement patterns in terms of increased knee abduction during weight-bearing activities, with special emphasis to individuals with ACLR. For that purpose, two systematic reviews with meta-analysis and three original studies were conducted. The meta-analyses revealed that female sex, reduced trunk lateral flexion strength, lower Gmax activation, increased hip external rotation ROM and reduced ankle dorsi-flexion ROM were associated with increased knee abduction during activity in healthy individuals, whereas lower extremity strength had, at best, a weak association with knee abduction. In patients with ACL injury the results from this thesis show that impaired sensory function (as assessed by kinesthesia and vibration sense), and alterations in trunk and knee muscle activation may be predictive of knee abduction in these patients. Consistent with reports in healthy individuals, our study found no association between reduced lower extremity strength and knee abduction in individuals with ACLR.

Gender differences in knee abduction

Women are reported to be at a greater risk of sustaining both a primary and subsequent ACL injury compared to men [15, 53]. That women perform weight-bearing activities with greater knee abduction has been proposed to be one explanation for this gender difference [19]. Despite inconsistent evidence, the assumption that women are more prone to knee abduction than men have been widely accepted in research and in clinical sport settings [7]. At the start of the work of this thesis, numerous studies had been conducted aiming at establish this gender-difference but with inconsistent results [105-108]. In our systematic review with meta-analysis (I) including 58 articles, we could conclude that both healthy women and women with PFP performed weight-bearing tasks with greater knee abduction compared to their male counterparts throughout the whole movement cycle, at initial contact, peak abduction and abduction excursion. We did not find a sufficient number of studies in patients with ACL injury to be able to perform meta-analysis to establish if this was the case also after ACL injury. However, in a single study of patients with ACL injury (that we identified as part of the

systematic review), Yamazaki et al., reported that women with ACLD exhibited greater 3D peak knee abduction than men with ACLD during a SLS [139], in line with the result in healthy individuals and in those with PFP.

In study III, we investigated possible gender differences in knee abduction by visual observation of the position of the knee in relation to the foot, during several different functional tasks with increasing difficulty in a group consisting of patients with either ACLD or ACLR. We found no gender difference in frontal plane knee position during any of the tasks, i.e., SLS, stair descending, forward lunge or drop jump. The main differences between our study and the study by Yamazaki et al., [139] are that we included both males and females, both patients with ACLD and ACLR and perhaps most important we assessed knee abduction as a KMFP by visual observation. A KMFP (the frontal plane position of the knee relative to the foot) as assessed visually may not reflect the knee abduction angle (the rotation of the shank in the frontal plane of the thigh) assessed with 3D motion analysis. Previous studies suggest that a visually observed KMFP is in fact a product of adduction and internal rotation of the hip rather than true knee abduction [126, 140]. This discrepancy in how knee abduction was assessed may be one explanation for the lack of gender difference in frontal plane knee position in study III. However, a recent review with meta-analysis from our research group concluded that visual observation of frontal plane knee position is valid in both 2D and 3D, i.e., the individuals that had a KMFP during weight-bearing activity also had greater knee abduction in both 2D and 3D compared to those with a KOPF [131].

Although our research confirms the suggested gender difference in 3D knee abduction in healthy individuals, the underlying reasons for this discrepancy are unclear. Several different factors, both biomechanical and neuromuscular have been proposed to pre-dispose women to this altered movement pattern. These factors include greater pelvic width, femoral anteversion, greater static knee abduction, joint laxity, greater landing forces, lower hamstrings to quadriceps torque ratios and reduced Gmax activation in females compared to males [7]. Also, the mechanism for sustaining an ACL injury may differ between males and females in such ways that knee abduction load seems to be consistently involved in ACL injury in females whereas knee abduction does not appear to be related to the injury mechanism in males [141, 142]. Together with the results from our review, this indicates that different strategies for men and women should be considered in prevention and rehabilitation of ACL injury. Specifically, women may benefit from exercise therapy aiming at improving knee alignment whereas other factors may be considered in men to prevent primary and subsequent ACL injury.

Sensorimotor factors associated with knee abduction

In the second study, the meta-analysis showed that reduced trunk lateral flexion strength as assessed by the side-bridge test, lower Gmax activation, increased hip external rotation ROM and reduced ankle dorsi-flexion ROM were associated with increased knee abduction during activity in healthy individuals, whereas lower extremity muscle strength was not. The factors contributing to knee abduction in individuals with ACL injury were however poorly investigated (II). Thus, our next step was to investigate possible sensorimotor factors that may contribute to increased knee abduction in patients with ACL injury. Patients suffering from ACL injury are known to have decreased sensorimotor function such as impaired proprioception [29], reduced muscle strength [25] and altered muscle activation patterns [26, 83] compared to non-injured individuals. Thus, and because that these factors are all modifiable by training we chose to include sensory function and trunk and lower extremity muscle strength and muscle activation as potential contributors for knee abduction in the subsequent studies.

Sensory function

We found that in patients with ACLD or ACLR, worse kinesthesia was associated with a KMFP during the drop jump in both men and women whereas worse vibration sense was associated with a KMFP during stair descending and forward lunge in women only (III). As mentioned in the introduction part of this thesis, the ACL includes several different mechanoreceptors responsible for the perception of the orientation of the segments in relation to each other and to the environment [143]. When the ACL is ruptured, the afferent input from these proprioceptive receptors is lost and a re-organization of the CNS then likely occurs to compensate for this deterioration to be able to maintain joint stability during activity [38-40]. Thus, it is plausible that impaired proprioception sense after ACL injury may affect the individual's ability to control frontal plane knee motion during activity.

Poor vibration sense has recently been linked to worse self-reported function in patients with ACL injury [69]. We are however not aware of any studies on the relation between vibration sense and objective function in these patients. In this thesis, we found worse vibration sense to be related to a KMFP during activity in women but not in men (III). The underlying cause for this gender difference is not clear. One could speculate that women may rely more on sensory abilities for joint stability whereas muscle function may be more important in men. Further research is however needed to investigate the underlying cause of the gender difference.

Muscle activation

In healthy individuals, we found lower Gmax activation to be associated with increased 3D peak knee abduction during activity (II). We were not able to confirm this result in patients with ACLR (V). However, we did find that lower IC activation on the non-injured side, higher ST activation, lower VM activation as well as greater ST/VM ratio was associated with greater knee abduction during the SLS and SLHD (V). Adequate activation of the trunk and core muscles has been proposed to decrease the risk of knee injury [10, 11]. Specifically, greater ipsilateral trunk motion has been reported to be related to greater peak knee abduction [144], increased knee abduction moment [145, 146] and a higher risk of sustaining an ACL injury [10, 11] in healthy individuals. If IC activation on the contralateral side is reduced, the individual may not be able to prevent lateral trunk lean to the ipsilateral side [147-149] and may thereby also increase knee abduction. This could constitute an explanation for the association between lower IC activation on the non-injured side and increased knee abduction on the injured side observed in study V.

Patients with ACL injury are reported to have higher average activation of the hamstring muscles [85] and lower activation of the quadriceps muscles [26] compared to healthy individuals. Since the hamstring muscles are important ACL agonists and resists anterior tibial translation, whereas the quadriceps is an ACL antagonist, these alterations may be due to central modifications to compensate for the decrease in joint stability that often is present after the loss of the ACL [91, 92]. The ST muscle is involved in adduction of the hip and in medial rotation of the hip and knee [150], and may thus play a role in knee abduction, since hip adduction and internal rotation are known contributors for this movement pattern [151]. Given the above-mentioned function of the hamstring as a joint stabilizer and that most patients are reconstructed using a ST graft, these patients are likely to have a preference of activating this muscle to a relatively high extent during activity. However, this compensatory strategy may in fact contribute to alterations in movement pattern in the form of increased knee abduction that may predispose these individuals to further injuries to the knee. Further research is required to confirm these findings and to assess whether this strategy is a preventive factor or risk factor for further injury.

Muscle strength

Previously, there has been great attention on the role of muscle strength for knee abduction and especially the hip muscles are believed to contribute to knee abduction due to their role of resisting adduction and internal rotation of the hip during activity [152]. But also, the muscles acting at the knee joint has been proposed to influence knee abduction because of the capacity of the hamstring and quadriceps muscles to support knee abduction/adduction moments of the knee [80]. In this thesis, we did not find that lower extremity strength including hip extension, abduction, external rotation or knee flexion and extension peak torque were associated with knee abduction in healthy individuals (II) or in those with ACLR (IV). In support of this finding, several studies report no improvement in knee abduction during activity after participation in a lower extremity strength training program [153-156]. An association between reduced hip muscle strength and increased hip muscle activation has been reported [139, 157-160], suggesting a compensatory increase in muscle activation to maintain joint stability and lower extremity alignment during activity when muscle strength is impaired. This indicates that the level of activation of the lower extremity muscles may be more important for postural orientation of the knee during activity than the absolute strength of the muscle as assessed by peak torque.

Trunk lateral flexion strength as assessed by the side-bridge test was associated with increased 3D peak knee abduction during a SLS in healthy individuals (II). We found however no such association in patients with ACLR (IV). This could indicate that the factors contributing to knee abduction may differ between healthy individuals and those with knee injury. However, study IV only included 29 participants with ACLR compared to 109 participants in the meta-analysis on healthy individuals (II), which could suggest that we were underpowered to detect this relation in study IV. Further studies are therefore needed to confirm the results in patients with ACLR.

The role of Knee abduction for ACL injury risk

Despite the fact that increased knee abduction seems to be involved in the knee injury mechanism in women [11, 161–163], whether knee abduction is a risk factor for sustaining the ACL injury is still a subject of debate. In the meta-analysis in this thesis we found that non-injured women performed weight-bearing activities with significantly greater knee abduction compared to men. This finding may support the hypothesis that knee abduction is a contributing factor for the high ACL injury incidence in women. However, to this date, four studies have

investigated the association between knee abduction at baseline and the risk of sustaining future primary ACL injury using a prospective study design [14, 18-20]. In the study by Hewett et al., 205 high school court sport female athletes were screened at baseline. The nine athletes that sustained an ACL injury during the season had significantly greater knee abduction at initial contact of a vertical drop jump landing compared to those that did not sustain an injury [19]. In contrast, in three other studies (screened athletes n=171-782, sustained ACL injuries n=5-42) there was no association between knee abduction during the drop jump task and the risk of sustaining a primary ACL injury [14, 18, 20]. It should however be noted that two of these latter studies were conducted in Norway in female adult elite team hand ball and soccer players [18, 20]. Norway is known for their rigorous injury prevention programs aiming to improve lower extremity alignment [164]. Consequently, these participants have most likely been participating in neuromuscular prevention training and may, therefore, be well trained to keep their knee over the foot during the assessment, which may have influenced the result of these studies. This is also reflected by the low amount of knee abduction observed during the drop jump in these studies (approximately 2 degrees) compared to the study by Hewett et al., where a mean knee abduction angle of 5 to 9 degrees was observed [18-20]. In a recent study, Fox et al., determined that normal knee abduction values during a vertical drop jump range from 5 to 17 degrees in females [165]. This implies that the participants in these studies were all in the normal range of knee abduction. Also, to date there is no consensus regarding the amount of knee abduction that is considered excessive enough to amplify the risk of injury. Taken together, the effect of knee abduction on knee injury risk in women is equivocal. Furthermore, the association between knee abduction and knee injury risk in men is unclear, since all available studies are conducted in women. That said, knee abduction may be relevant for knee injury in younger individuals, not playing on an elite level. Although the evidence for the role of knee abduction in primary ACL injury is not univocal, there seems to be an association between greater knee abduction during a drop jump at baseline and a second ACL injury in both men and women [18, 151]. However, further studies are still needed to establish the importance of lower extremity alignment during activity for the risk of sustaining both the primary and subsequent ACL injury.

Methodological considerations

General limitations (I-V)

This thesis includes some limitations. First, a knee injury may occur during several different movements, e.g., landings and cutting movements. Since increased knee abduction has been suggested to account for some of the gender-difference in knee injury incidence, we were in study I initially interested in investigating if women in general performed weight-bearing activities with increased knee abduction compared to men. The same applies for study II, where we wanted to elucidate the effect of the investigated factors on knee abduction during weight-bearing activities. Consequently, we included all available tasks in the meta-analyses. We are, however, aware that this approach may introduce some potential bias. It could not be ruled out that the different movement strategies used in each task may influence the magnitude of knee abduction differently or affect the contribution of each sensorimotor factor for knee abduction in a different manner and, thus, have an influence on the results. In an effort to deal with this problem, we performed separate analyses for each task that included two or more articles. Although this was an adequate approach, including separate analysis for most tasks in study I, it was not feasible in study II where only the association between hip strength and knee abduction during the SLS was eligible for separate analysis.

Second, in study III we included patients with ACLD and ACLR in the same analysis to reflect the population that typically presents clinically for rehabilitation post ACL injury. However, studies have shown that there may be differences both in proprioceptive acuity [64] and in the relation between sensory function and self-reported knee function [69] between individuals with ACLD and ACLR. Thus, it is possible that the result would have been different if we had analysed these groups separately. It is also for this reason we chose to only include participants with an ACLR in studies IV and V in an effort to have a more homogeneous population.

Third, no power calculations were performed in studies III-V. The sample in study III constituted a sample of convenience (n=51) and in studies IV and V all individuals that have performed an ACLR at Skåne University Hospital in Sweden between June and December, 2015 were invited to participate in the study (n=106). Out of the 106 that were invited, only 29 participants were finally included in the studies. The data was collected between February 16th and May 30th, 2016. Due to time constraints we were unfortunately not able to include additional participants within the time frame of this thesis. Thus, there is a possibility that the lack of associations between some variables and knee abduction are due to a lack of power. Consequently, the small sample also

prevented us from performing separate analysis for men and women in study IV and V, which was our aim initially. Although we adjusted for sex in the regression models, the result may have been different if we were able to perform gender stratified analyses. Thus, a larger sample size is needed to determine our findings in studies IV and V.

Sensory measures

In study III we investigated the relation between proprioception (kinesthesia) and visually observed frontal plane knee position. Assessment of kinesthesia (and all other measures of proprioception) depends on the individual's conscious perception of the movement and may thus be affected by cognitive abilities. Also, the assessment of "conscious" proprioception may not reflect the unconscious movement strategies that are suggested to play a role in joint stability. Furthermore, although great effort is being put on reducing sensory input from the skin and the equipment by padding and the use of socks, it is not possible to perform such assessment without affecting other sensory mechanoreceptors. For example, receptor in the skin will send information about movement to the CNS when the skin is stretched during flexion/extension of the knee, thus it is hard to distinguish the source/origin of the afferent signals that contribute to the conscious perception of the movement during the assessment [113]. Also, in our study, kinesthesia was assessed during passive movement. It has been proposed that this approach is not ecologically valid [113], since passive movement does not reflect a real-life situation [166]. Thus, more functional measures of proprioception, as used in other studies [167, 168], that includes weight-bearing and active movements may have provided other results.

Muscle activation

Traditionally there have been several methodological limitations associated with surface EMG data collection, such as the difference in EMG signal amplitude due to the size of the electrode, type of electrode fixation, inter-electrode distance and/or location of the reference electrode, which have made it hard to compare results between studies. These limitations are minimized with modern wireless EMG equipment with self-adhesive standard dual electrodes with a fixed inter-electrode distance [123]. However, other limitations that should be considered are the high risk of collecting cross-talks from adjacent muscles [123, 169] and the presence of baseline noise and movement artefacts [170] that may affect the EMG signal. Movement artefacts are a particularly problem during dynamic tasks such as the SLHD used in study V. In fact, we had to exclude 7 out of the 29

participants from the analysis on IC muscle activation on the injured side due to movement artefacts, which may have affected the statistical power of this analysis.

To be able to compare muscle activation amplitude between individuals, you must normalize the data. The most common normalization method is to divide the processed EMG amplitude for each muscle collected during the trial with the amplitude collected for that specific muscle during a MVC [171, 172]. A common problem with this method is however that you cannot be sure that the data collected during the MVCs really reflects the true MVC. This is especially problematic when, as in study V, the data of interest are collected during dynamic movements (SLS and SLHD) but are then normalized to data collected during isometric testing. This was not a general problem in our study (V), but for one muscle (VMO), 11 (38%) participants had a task data/MVC ratio of between 1 and 3. In other words, in these cases the participants activated their VMO up to three times more during the task than during the MVC test. This implies that the isometric MVC did not appear to represent the true MVC in these cases. The only method to ensure that a true MVC is collected is by a twitch interpolation technique [173]. This was not feasible in our study since this technique involves percutaneous electrical impulses that are both uncomfortable for the participants and require advanced equipment. However, in an effort to gain the maximal recruitment of the muscles, we used a standardized protocol, where the participants were given the same instructions on how to perform the task and were verbally encourage to produce maximum force during the task. In addition, a question about perceived pain during the performance of the task was asked to minimize pain related failure during the tasks.

3D motion analysis

Although 3D motion analysis is considered the “gold standard” to assess human movement, this method is not without limitations. Several factors affect the probability of obtaining accurate and reliable movement data from the motion analysis system. Systematic errors, associated with for example laboratory set-up and calibration of the cameras, and random errors originating from electronic noise are common instrument-related problems in 3D assessments [174]. The latter may, however, be minimized through certain filtering and smoothing techniques [133]. Another limitation associated with 3D assessment includes difficulties with accurate identification of the anatomical bone landmarks for marker placement and the centre of the joints which have substantial influence on the reliability of the data obtained [175]. Furthermore, soft tissues artefacts (STA) are an important factor that should be considered in 3D motion analysis. This is a particular problem during high impact movements such as the SLHD. STA is provoked by displacement of the skin where the markers are attached in relation to the

subcutaneous bone structure and may be influenced by muscle contractions, type of activity and the location of the markers [176]. Previous studies have reported that the thigh is associated with the greatest error during functional tasks such as stair descending, cutting and SLHD [176-179]. Also, the use of skin markers showed poor reliability in frontal plane knee rotations with errors up to 70% between the assessed movement and the actual real joint movement [176]. To minimize the errors from STA, the marker model in our studies (IV and V) was designed according to the Visual 3D recommendations based on the work by Cappozzo et al [180]. Despite these limitations, 3D motion analysis is the closest we can get to identify the movement of the joints. 3D knee abduction has shown high inter and intra reliability during several tasks including different jumping tasks and the SLS [181-183] as well as high reliability between different laboratories and systems [184].

Clinical implications

Previous research proposes an association between increased knee abduction during activity and worse self-reported function [97, 98] and worse hop performance [96], respectively, in individuals with ACL injury. In addition, this movement pattern seems to increase the risk of subsequent ACL injury [18, 151]. Thus, exercises with the aim of decreasing knee abduction are today often incorporated in rehabilitation programs after knee injury [5]. Studies report that neuromuscular training including plyometric exercises, strength training, proprioception training and modification of landing techniques may reduce the risk of injury [185, 186] and improve biomechanical factors associated with knee injury such as knee abduction and lateral trunk lean [187]. However, the specific factors that should be targeted to reduce knee abduction have not been established.

In this thesis, we found that altered muscle activation rather than muscle strength seem to play a role in knee abduction in both healthy individuals (II) and in those with ACLR (IV and V). Previous research also found no improvement in frontal plane knee motion after strength training protocols in healthy individuals and in those with PFP [153-156]. This indicates that absolute strength may not be as important for knee abduction as previously believed and that incorporating exercises targeting the activation of the muscles instead of force may be more beneficial for improving this movement pattern. Zebis et al., reported that a 12-week neuromuscular exercise program improved knee muscle activation pattern commonly associated with knee injury risk [188].

Another factor that may be important in the prevention and rehabilitation of ACL injury is ankle dorsiflexion ROM. In study II, we found that those with a KMFP

during activity had significantly reduced ankle-dorsi flexion ROM compared to those with a KOFP. Reduced ankle ROM is proposed to increase foot pronation and prevent knee flexion and may thereby introduce compensatory strategies at the knee such as increased knee abduction [189-191].

Proprioceptive training should also be considered when aiming at reducing knee abduction after ACL injury, since we found that individuals with a KMFP had decreased proprioception during a drop jump compared to those with a KOFP (III). In study I, we reported that women exhibit greater knee abduction during weight-bearing activity than men. This may suggest that exercise therapies aiming at reducing knee abduction would be more beneficial in women than in men. Other factors than knee abduction may therefore be considered in men to improve function and reduce the risk of subsequent knee injury.

The altered knee movement pattern observed after ACL rupture may also be elicited by neuroplastic changes and alterations to the CNS after injury which may be hard to target with conventional rehabilitation strategies. Electromyographic biofeedback, transcutaneous electrical nerve stimulation and neuromuscular electrical stimulation are methods that may be helpful in restoring neuromuscular alteration after knee injury by improving motor unit recruitment, reflex excitability and muscle inhibition as well as exaggerate voluntary muscle contraction [192]. Also, rehabilitation after ACL injury may be enhanced by integrating exercises with higher demands on the brain. Incorporating cognitive tasks that include visual-spatial-motor tasks and mental imagery may ameliorate neuromuscular control and joint stability by increasing muscular activation and improving coordination after ACL injury and thereby potentially decrease frontal plane knee motion [193, 194].

In this thesis, we investigated gender and sensorimotor factors modifiable by training as possible contributors to knee abduction. However, we found only moderate associations between the investigated factors and knee abduction, implying that there are other factors that also contribute to knee abduction. Several non-modifiable factors, such as greater tibial slope, pelvic width and static knee alignment have been proposed to influence frontal plane knee motion during activity [195, 196]. Also, the relative contribution of the different factors for knee abduction is not clear. Thus, future studies that include both structural and sensorimotor factors as potential contributors for knee abduction are warranted. Also, most studies on both knee injury risk and movement pattern alterations are conducted in women. Further studies are thus needed to evaluate which specific factors that is associated with knee abduction and knee injury risk in men.

Conclusions

The result from this thesis indicates that alterations in sensorimotor function play a role in knee abduction during weight-bearing activities. Specifically, lower trunk lateral flexion strength, reduced ankle dorsi-flexion ROM and lower Gmax activation are associated with greater knee abduction in healthy individuals. In patients with ACL injury, reduced sensory function and alterations in trunk and knee muscle activation may increase knee abduction, whereas lower extremity muscle strength does not seem to be important for knee abduction in individuals both with and without knee injury. Given that healthy women perform weight-bearing activities with increased knee abduction compared to men, there may need to be a greater focus on this movement pattern during prevention and rehabilitation training in women, whereas other factors may be more relevant in men. Further investigations are needed to confirm our results and to establish which factors, both structural and neuromuscular, that contributes to knee abduction in men and women with ACL injury.

Acknowledgements

I would like to express my sincere gratitude to all the people that have been with me on this journey. This work would not have been possible without you! A special thanks to:

Eva Ageberg, my main supervisor, for introducing me to research and making me love it, for all creative discussions, for teaching me academic writing, for excellent guidance and support, but most of all thank you for your incredible enthusiasm and your ability to always lift me up when I lost faith in myself.

Mark Creaby, my co-supervisor, for sharing your vast knowledge in biomechanics, for your great attention to details that made this work so much better and for your excellent guidance and support.

Jenny Älmqvist Nae, co-author and fellow PhD student, for your great input and help in collecting data for all my studies.

Tim Blackmore, co-author, for all the hours you have spent with the visual 3D program to sort out and process my movement analysis data.

Melinda Franettovich Smith, co-author, for sharing your knowledge in EMG and for answering all my EMG questions.

All colleges in the research group, for the creative discussions and for all the fun and laughs, it has been a fantastic time and I will really miss you all. A special thanks to Vala Flosadottir for coming up with the title of this thesis.

Vibeke Horstman, *Christel Nielsen* and *Tommy Schyman* for statistical advice and help when I tended to complicate things too much.

Carsten Juhl for sharing your great knowledge, and for answering my questions, on systematic reviews and meta-analyses.

Marianne Gullberg for your generosity in letting us use the Humanities Lab at Lund University to collect our data, and a special thanks to *Stefan Lindgren* and *Henrik Garde* for your time and support when we had trouble with the EMG-synchronization.

My parents, *Anita Ryman* och *Leif Svensson* for your endless support and for making me who I am today and my sister *Annika Heidefors* and brother *Kent Svensson* for making life so much more fun

Anders Cronström, *Rasmus Brandin* and *Olivia Brandin* for your love and support and for putting up with me during these years, I know I have been a real pain sometimes, love you endlessly.

Lena Brandin for your engagement and support during the years especially when it comes to your grandchildren and for just being you

I will also like to thank all men and women who have taken their time to participate in the studies.

I am also grateful for funding from the Swedish Research Council, the Crafoord Foundation, Skåne Regional Council, the Swedish Rheumatism Association, the Swedish National Centre for Research in Sports, and the Faculty of Medicine of Lund University.

References

1. Bicer EK, Lustig S, Servien E, Selmi TA, Neyret P: Current knowledge in the anatomy of the human anterior cruciate ligament. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2010, 18(8):1075-1084.
2. Butler DL, Noyes FR, Grood ES: Ligamentous restraints to anterior-posterior drawer in the human knee. A biomechanical study. *The Journal of bone and joint surgery American volume* 1980, 62(2):259-270.
3. Mather RC, 3rd, Koenig L, Kocher MS, Dall TM, Gallo P, Scott DJ, Bach BR, Jr., Spindler KP: Societal and economic impact of anterior cruciate ligament tears. *The Journal of bone and joint surgery American volume* 2013, 95(19):1751-1759.
4. Frobell RB, Lohmander LS, Roos HP: Acute rotational trauma to the knee: poor agreement between clinical assessment and magnetic resonance imaging findings. *Scandinavian journal of medicine & science in sports* 2007, 17(2):109-114.
5. Renstrom P, Ljungqvist A, Arendt E, Beynnon B, Fukubayashi T, Garrett W, Georgoulis T, Hewett TE, Johnson R, Krosshaug T et al: Non-contact ACL injuries in female athletes: an International Olympic Committee current concepts statement. *British journal of sports medicine* 2008, 42(6):394-412.
6. Stuelcken MC, Mellifont DB, Gorman AD, Sayers MG: Mechanisms of anterior cruciate ligament injuries in elite women's netball: a systematic video analysis. *Journal of sports sciences* 2016, 34(16):1516-1522.
7. Hewett TE, Myer GD, Ford KR, Paterno MV, Quatman CE: Mechanisms, prediction, and prevention of ACL injuries: Cut risk with three sharpened and validated tools. *Journal of orthopaedic research : official publication of the Orthopaedic Research Society* 2016, 34(11):1843-1855.
8. Alentorn-Geli E, Myer GD, Silvers HJ, Samitier G, Romero D, Lazaro-Haro C, Cugat R: Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying risk factors. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2009, 17(7):705-729.
9. Balazs GC, Pavey GJ, Brelin AM, Pickett A, Keblish DJ, Rue JP: Risk of Anterior Cruciate Ligament Injury in Athletes on Synthetic Playing Surfaces: A Systematic Review. *The American journal of sports medicine* 2015, 43(7):1798-1804.
10. Zazulak BT, Hewett TE, Reeves NP, Goldberg B, Cholewicki J: Deficits in neuromuscular control of the trunk predict knee injury risk: a prospective biomechanical-epidemiologic study. *The American journal of sports medicine* 2007, 35(7):1123-1130.

11. Hewett TE, Torg JS, Boden BP: Video analysis of trunk and knee motion during non-contact anterior cruciate ligament injury in female athletes: lateral trunk and knee abduction motion are combined components of the injury mechanism. *British journal of sports medicine* 2009, 43(6):417-422.
12. Rafeeuddin R, Sharir R, Staes F, Dingenen B, George K, Robinson MA, Vanrenterghem J: Mapping current research trends on neuromuscular risk factors of non-contact ACL injury. *Physical therapy in sport : official journal of the Association of Chartered Physiotherapists in Sports Medicine* 2016, 22:101-113.
13. Vacek PM, Slauterbeck JR, Tourville TW, Sturnick DR, Holterman LA, Smith HC, Shultz SJ, Johnson RJ, Tourville KJ, Beynnon BD: Multivariate Analysis of the Risk Factors for First-Time Noncontact ACL Injury in High School and College Athletes: A Prospective Cohort Study With a Nested, Matched Case-Control Analysis. *The American journal of sports medicine* 2016, 44(6):1492-1501.
14. Leppanen M, Pasanen K, Kujala UM, Vasankari T, Kannus P, Ayrano S, Krosshaug T, Bahr R, Avela J, Perttunen J et al: Stiff Landings Are Associated With Increased ACL Injury Risk in Young Female Basketball and Floorball Players. *The American journal of sports medicine* 2016.
15. Walden M, Hagglund M, Werner J, Ekstrand J: The epidemiology of anterior cruciate ligament injury in football (soccer): a review of the literature from a gender-related perspective. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2011, 19(1):3-10.
16. Joseph AM, Collins CL, Henke NM, Yard EE, Fields SK, Comstock RD: A multisport epidemiologic comparison of anterior cruciate ligament injuries in high school athletics. *Journal of athletic training* 2013, 48(6):810-817.
17. VandenBerg C, Crawford EA, Sibilsky Enselman E, Robbins CB, Wojtys EM, Bedi A: Restricted Hip Rotation Is Correlated With an Increased Risk for Anterior Cruciate Ligament Injury. *Arthroscopy : the journal of arthroscopic & related surgery : official publication of the Arthroscopy Association of North America and the International Arthroscopy Association* 2016.
18. Krosshaug T, Steffen K, Kristianslund E, Nilstad A, Mok KM, Myklebust G, Andersen TE, Holme I, Engebretsen L, Bahr R: The Vertical Drop Jump Is a Poor Screening Test for ACL Injuries in Female Elite Soccer and Handball Players: A Prospective Cohort Study of 710 Athletes. *The American journal of sports medicine* 2016.
19. Hewett TE, Myer GD, Ford KR, Heidt RS, Jr., Colosimo AJ, McLean SG, van den Bogert AJ, Paterno MV, Succop P: Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *The American journal of sports medicine* 2005, 33(4):492-501.
20. Nilstad A, Andersen TE, Bahr R, Holme I, Steffen K: Risk factors for lower extremity injuries in elite female soccer players. *The American journal of sports medicine* 2014, 42(4):940-948.
21. Steffen K, Nilstad A, Kristianslund EK, Myklebust G, Bahr R, Krosshaug T: Association between Lower Extremity Muscle Strength and Noncontact ACL Injuries. *Medicine and science in sports and exercise* 2016, 48(11):2082-2089.

22. Khayambashi K, Ghoddosi N, Straub RK, Powers CM: Hip Muscle Strength Predicts Noncontact Anterior Cruciate Ligament Injury in Male and Female Athletes: A Prospective Study. *The American journal of sports medicine* 2016, 44(2):355-361.
23. Ageberg E: Consequences of a ligament injury on neuromuscular function and relevance to rehabilitation - using the anterior cruciate ligament-injured knee as model. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2002, 12(3):205-212.
24. Culvenor AG, Alexander BC, Clark RA, Collins NJ, Ageberg E, Morris HG, Whitehead TS, Crossley KM: Dynamic Single-Leg Postural Control Is Impaired Bilaterally Following Anterior Cruciate Ligament Reconstruction: Implications for Reinjury Risk. *The Journal of orthopaedic and sports physical therapy* 2016, 46(5):357-364.
25. Petersen W, Taheri P, Forkel P, Zantop T: Return to play following ACL reconstruction: a systematic review about strength deficits. *Archives of orthopaedic and traumatic surgery Archiv fur orthopadische und Unfall-Chirurgie* 2014, 134(10):1417-1428.
26. Hart JM, Pietrosimone B, Hertel J, Ingersoll CD: Quadriceps activation following knee injuries: a systematic review. *Journal of athletic training* 2010, 45(1):87-97.
27. Steele JR, Brown JM: Effects of chronic anterior cruciate ligament deficiency on muscle activation patterns during an abrupt deceleration task. *Clinical biomechanics (Bristol, Avon)* 1999, 14(4):247-257.
28. Perraton L, Clark R, Crossley K, Pua YH, Whitehead T, Morris H, Telianidis S, Bryant A: Impaired voluntary quadriceps force control following anterior cruciate ligament reconstruction: relationship with knee function. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2016.
29. Relph N, Herrington L, Tyson S: The effects of ACL injury on knee proprioception: a meta-analysis. *Physiotherapy* 2013.
30. Goldstein J, Bosco JA, 3rd: The ACL-deficient knee: natural history and treatment options. *Bull Hosp Jt Dis* 2001, 60(3-4):173-178.
31. Culvenor AG, Collins NJ, Guermazi A, Cook JL, Vicenzino B, Khan KM, Beck N, van Leeuwen J, Crossley KM: Early knee osteoarthritis is evident one year following anterior cruciate ligament reconstruction: a magnetic resonance imaging evaluation. *Arthritis & rheumatology (Hoboken, NJ)* 2015, 67(4):946-955.
32. Simon D, Mascarenhas R, Saltzman BM, Rollins M, Bach BR, Jr., MacDonald P: The relationship between anterior cruciate ligament injury and osteoarthritis of the knee. *Advances in orthopedics* 2015, 2015:928301.
33. Faltstrom A, Hagglund M, Kvist J: Patient-reported knee function, quality of life, and activity level after bilateral anterior cruciate ligament injuries. *The American journal of sports medicine* 2013, 41(12):2805-2813.
34. Filbay SR, Culvenor AG, Ackerman IN, Russell TG, Crossley KM: Quality of life in anterior cruciate ligament-deficient individuals: a systematic review and meta-analysis. *British journal of sports medicine* 2015, 49(16):1033-1041.
35. Garcia GH, Wu HH, Park MJ, Tjoumakaris FP, Tucker BS, Kelly JD, Sennett BJ: Depression Symptomatology and Anterior Cruciate Ligament Injury: Incidence and

- Effect on Functional Outcome--A Prospective Cohort Study. *The American journal of sports medicine* 2016, 44(3):572-579.
36. Baumeister J, Reinecke K, Weiss M: Changed cortical activity after anterior cruciate ligament reconstruction in a joint position paradigm: an EEG study. *Scandinavian journal of medicine & science in sports* 2008, 18(4):473-484.
 37. Baumeister J, Reinecke K, Schubert M, Weiss M: Altered electrocortical brain activity after ACL reconstruction during force control. *Journal of orthopaedic research : official publication of the Orthopaedic Research Society* 2011, 29(9):1383-1389.
 38. Kapreli E, Athanasopoulos S, Gliatis J, Papathanasiou M, Peeters R, Strimpakos N, Van Hecke P, Gouliamos A, Sunaert S: Anterior cruciate ligament deficiency causes brain plasticity: a functional MRI study. *The American journal of sports medicine* 2009, 37(12):2419-2426.
 39. Grooms DR, Page SJ, Nichols-Larsen DS, Chaudhari AM, White SE, Onate JA: Neuroplasticity Associated With Anterior Cruciate Ligament Reconstruction. *The Journal of orthopaedic and sports physical therapy* 2016:1-27.
 40. Grooms DR, Page SJ, Onate JA: Brain Activation for Knee Movement Measured Days Before Second Anterior Cruciate Ligament Injury: Neuroimaging in Musculoskeletal Medicine. *Journal of athletic training* 2015, 50(10):1005-1010.
 41. Ageberg E, Thomee R, Neeter C, Silbernagel KG, Roos EM: Muscle strength and functional performance in patients with anterior cruciate ligament injury treated with training and surgical reconstruction or training only: a two to five-year followup. *Arthritis and rheumatism* 2008, 59(12):1773-1779.
 42. Ericsson YB, Roos EM, Frobell RB: Lower extremity performance following ACL rehabilitation in the KANON-trial: impact of reconstruction and predictive value at 2 and 5 years. *British journal of sports medicine* 2013, 47(15):980-985.
 43. Frobell RB, Roos EM, Roos HP, Ranstam J, Lohmander LS: A randomized trial of treatment for acute anterior cruciate ligament tears. *The New England journal of medicine* 2010, 363(4):331-342.
 44. Luc B, Gribble PA, Pietrosimone BG: Osteoarthritis prevalence following anterior cruciate ligament reconstruction: a systematic review and numbers-needed-to-treat analysis. *Journal of athletic training* 2014, 49(6):806-819.
 45. Nagelli CV, Hewett TE: Should Return to Sport be Delayed Until 2 Years After Anterior Cruciate Ligament Reconstruction? Biological and Functional Considerations. *Sports medicine (Auckland, NZ)* 2016.
 46. Tan SH, Lau BP, Khin LW, Lingaraj K: The Importance of Patient Sex in the Outcomes of Anterior Cruciate Ligament Reconstructions: A Systematic Review and Meta-analysis. *The American journal of sports medicine* 2016, 44(1):242-254.
 47. Ardern CL, Taylor NF, Feller JA, Webster KE: Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *British journal of sports medicine* 2014, 48(21):1543-1552.

48. Filbay SR, Ackerman IN, Russell TG, Crossley KM: Return to sport matters-longer-term quality of life after ACL reconstruction in people with knee difficulties. *Scandinavian journal of medicine & science in sports* 2016.
49. Christino MA, Fantry AJ, Vopat BG: Psychological Aspects of Recovery Following Anterior Cruciate Ligament Reconstruction. *The Journal of the American Academy of Orthopaedic Surgeons* 2015, 23(8):501-509.
50. Bauer M, Feeley BT, Wawrzyniak JR, Pinkowsky G, Gallo RA: Factors affecting return to play after anterior cruciate ligament reconstruction: a review of the current literature. *The Physician and sportsmedicine* 2014, 42(4):71-79.
51. Sonesson S, Kvist J, Ardern C, Osterberg A, Silbernagel KG: Psychological factors are important to return to pre-injury sport activity after anterior cruciate ligament reconstruction: expect and motivate to satisfy. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2016.
52. Wiggins AJ, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD: Risk of Secondary Injury in Younger Athletes After Anterior Cruciate Ligament Reconstruction: A Systematic Review and Meta-analysis. *The American journal of sports medicine* 2016, 44(7):1861-1876.
53. Shelbourne KD, Gray T, Haro M: Incidence of subsequent injury to either knee within 5 years after anterior cruciate ligament reconstruction with patellar tendon autograft. *The American journal of sports medicine* 2009, 37(2):246-251.
54. Capin JJ, Khandha A, Zarzycki R, Manal K, Buchanan TS, Snyder-Mackler L: Gait mechanics and second ACL rupture: Implications for delaying return-to-sport. *Journal of orthopaedic research : official publication of the Orthopaedic Research Society* 2016.
55. Grindem H, Snyder-Mackler L, Moksnes H, Engebretsen L, Risberg MA: Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: the Delaware-Oslo ACL cohort study. *British journal of sports medicine* 2016, 50(13):804-808.
56. Johansson H, Sjölander P, Sojka P: A sensory role for the cruciate ligaments. *Clinical orthopaedics and related research* 1991(268):161-178.
57. Riemann BL, Lephart SM: The sensorimotor system, part I: the physiologic basis of functional joint stability. *Journal of athletic training* 2002, 37(1):71-79.
58. Solomonow M, Krogsgaard M: Sensorimotor control of knee stability. A review. *Scandinavian journal of medicine & science in sports* 2001, 11(2):64-80.
59. Johansson H: Role of knee ligaments in proprioception and regulation of muscle stiffness. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 1991, 1(3):158-179.
60. Tsuda E, Okamura Y, Otsuka H, Komatsu T, Tokuya S: Direct evidence of the anterior cruciate ligament-hamstring reflex arc in humans. *The American journal of sports medicine* 2001, 29(1):83-87.
61. Dyhre-Poulsen P, Krogsgaard MR: Muscular reflexes elicited by electrical stimulation of the anterior cruciate ligament in humans. *Journal of applied physiology (Bethesda, Md : 1985)* 2000, 89(6):2191-2195.

62. Melnyk M, Faist M, Gothner M, Claes L, Friemert B: Changes in stretch reflex excitability are related to "giving way" symptoms in patients with anterior cruciate ligament rupture. *J Neurophysiol* 2007, 97(1):474-480.
63. Dhaher YY, Tsoumanis AD, Rymer WZ: Reflex muscle contractions can be elicited by valgus positional perturbations of the human knee. *Journal of biomechanics* 2003, 36(2):199-209.
64. Relph N, Herrington L, Tyson S: The effects of ACL injury on knee proprioception: a meta-analysis. *Physiotherapy* 2014, 100(3):187-195.
65. Relph N, Herrington L: Knee joint position sense ability in elite athletes who have returned to international level play following ACL reconstruction: A cross-sectional study. *The Knee* 2016.
66. Roberts D, Ageberg E, Andersson G, Fridén T: Clinical measurements of proprioception, muscle strength and laxity in relation to function in the ACL-injured knee. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2007, 15(1):9-16.
67. Ageberg E, Roberts D, Holmstrom E, Fridén T: Balance in single-limb stance in patients with anterior cruciate ligament injury: relation to knee laxity, proprioception, muscle strength, and subjective function. *The American journal of sports medicine* 2005, 33(10):1527-1535.
68. Roberts D, Fridén T, Zätterström R, Lindstrand A, Moritz U: Proprioception in people with anterior cruciate ligament-deficient knees: comparison of symptomatic and asymptomatic patients. *The Journal of orthopaedic and sports physical therapy* 1999, 29(10):587-594.
69. Cronstrom A, Roos E, Ageberg E: Association between sensory function and hop performance and self-reported outcomes in patients with anterior cruciate ligament injury. *Accepted to Open access journal of sports medicine* 2016.
70. Noback C, Strominger N, Demarest R, Ruggiero D: *The human nervous system: structure and function*, 6th edn. Totowa (NJ): Human Press; 2005.
71. Gilman S: Joint position sense and vibration sense: anatomical organisation and assessment. *J Neurol Neurosurg Psychiatry* 2002, 73(5):473-477.
72. Shakoor N, Agrawal A, Block JA: Reduced lower extremity vibratory perception in osteoarthritis of the knee. *Arthritis and rheumatism* 2008, 59(1):117-121.
73. Shakoor N, Lee KJ, Fogg LF, Block JA: Generalized vibratory deficits in osteoarthritis of the hip. *Arthritis and rheumatism* 2008, 59(9):1237-1240.
74. Shakoor N, Lee KJ, Fogg LF, Wimmer MA, Foucher KC, Mikolaitis RA, Block JA: The relationship of vibratory perception to dynamic joint loading, radiographic severity, and pain in knee osteoarthritis. *Arthritis and rheumatism* 2012, 64(1):181-186.
75. Miyazaki T, Wada M, Kawahara H, Sato M, Baba H, Shimada S: Dynamic load at baseline can predict radiographic disease progression in medial compartment knee osteoarthritis. *Annals of the rheumatic diseases* 2002, 61(7):617-622.
76. Shakoor N, Felson DT, Niu J, Nguyen US, Segal NA, Singh JA, Nevitt MC: The association of vibratory perception and muscle strength with the incidence and

- worsening of knee instability: the Multicenter Osteoarthritis Study. Arthritis & rheumatology (Hoboken, NJ) 2016.
77. Sandal LF: Reliability and construct validity of vibratory perception threshold testing. Master thesis. Copenhagen: University of southern Denmark; 2011.
 78. Thorlund JB, Shakoor N, Ageberg E, Sandal LF, Block JA, Roos EM: Vibratory perception threshold in young and middle-aged patients at high risk of knee osteoarthritis compared to controls. Arthritis care & research 2012, 64(1):144-148.
 79. Riemann BL, Lephart SM: The Sensorimotor System, Part II: The Role of Proprioception in Motor Control and Functional Joint Stability. Journal of athletic training 2002, 37(1):80-84.
 80. Lloyd DG, Buchanan TS: Strategies of muscular support of varus and valgus isometric loads at the human knee. Journal of biomechanics 2001, 34(10):1257-1267.
 81. Blackburn JT, Norcross MF, Padua DA: Influences of hamstring stiffness and strength on anterior knee joint stability. Clinical biomechanics (Bristol, Avon) 2011, 26(3):278-283.
 82. Tengman E, Brax Olofsson L, Stensdotter AK, Nilsson KG, Hager CK: Anterior cruciate ligament injury after more than 20 years. II. Concentric and eccentric knee muscle strength. Scandinavian journal of medicine & science in sports 2014.
 83. Dingenen B, Janssens L, Luyckx T, Claes S, Bellemans J, Staes FF: Lower extremity muscle activation onset times during the transition from double-leg stance to single-leg stance in anterior cruciate ligament injured subjects. Human movement science 2015, 44:234-245.
 84. Oeffinger DJ, Shapiro R, Nyland J, Pienkowski D, Caborn DN: Delayed gastrocnemius muscle response to sudden perturbation in rehabilitated patients with anterior cruciate ligament reconstruction. Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA 2001, 9(1):19-27.
 85. Shanbehzadeh S, Mohseni Bandpei MA, Ehsani F: Knee muscle activity during gait in patients with anterior cruciate ligament injury: a systematic review of electromyographic studies. Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA 2015.
 86. Hall M, Stevermer CA, Gillette JC: Muscle activity amplitudes and co-contraction during stair ambulation following anterior cruciate ligament reconstruction. Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology 2015, 25(2):298-304.
 87. Swanik CB, Lephart SM, Giraldo JL, Demont RG, Fu FH: Reactive muscle firing of anterior cruciate ligament-injured females during functional activities. Journal of athletic training 1999, 34(2):121-129.
 88. Nyland J, Klein S, Caborn DN: Lower extremity compensatory neuromuscular and biomechanical adaptations 2 to 11 years after anterior cruciate ligament reconstruction. Arthroscopy : the journal of arthroscopic & related surgery : official publication of the Arthroscopy Association of North America and the International Arthroscopy Association 2010, 26(9):1212-1225.

89. Jennings AG, Seedhom BB: The measurement of muscle stiffness in anterior cruciate injuries -- an experiment revisited. *Clinical biomechanics* (Bristol, Avon) 1998, 13(2):138-140.
90. McNair PJ, Wood GA, Marshall RN: Stiffness of the hamstring muscles and its relationship to function in anterior cruciate ligament deficient individuals. *Clinical biomechanics* (Bristol, Avon) 1992, 7(3):131-137.
91. Adouni M, Shirazi-Adl A, Marouane H: Role of gastrocnemius activation in knee joint biomechanics: gastrocnemius acts as an ACL antagonist. *Computer methods in biomechanics and biomedical engineering* 2016, 19(4):376-385.
92. Cowling EJ, Steele JR: Is lower limb muscle synchrony during landing affected by gender? Implications for variations in ACL injury rates. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2001, 11(4):263-268.
93. Barcellona MG, Morrissey MC, Milligan P, Amis AA: The effect of thigh muscle activity on anterior knee laxity in the uninjured and anterior cruciate ligament-injured knee. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2014, 22(11):2821-2829.
94. Horak FB: Postural orientation and equilibrium: what do we need to know about neural control of balance to prevent falls? *Age and ageing* 2006, 35 Suppl 2:ii7-ii11.
95. Trulsson A, Garwicz M, Ageberg E: Postural orientation in subjects with anterior cruciate ligament injury: development and first evaluation of a new observational test battery. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2010, 18(6):814-823.
96. Trulsson A, Roos EM, Ageberg E, Garwicz M: Relationships between postural orientation and self reported function, hop performance and muscle power in subjects with anterior cruciate ligament injury. *BMC musculoskeletal disorders* 2010, 11:143.
97. Ageberg E, Roos EM: The Association Between Knee Confidence and Muscle Power, Hop Performance, and Postural Orientation in People With Anterior Cruciate Ligament Injury. *The Journal of orthopaedic and sports physical therapy* 2016, 46(6):477-482.
98. Flosadottir V, Roos EM, Ageberg E: Muscle function is associated with future patient-reported outcomes in young adults with ACL injury. *BMJ Open Sport & Exercise Medicine* 2016, 2(1).
99. Wyndow N, De Jong A, Rial K, Tucker K, Collins N, Vicenzino B, Russell T, Crossley K: The relationship of foot and ankle mobility to the frontal plane projection angle in asymptomatic adults. *Journal of foot and ankle research* 2016, 9:3.
100. Bell DR, Padua DA, Clark MA: Muscle strength and flexibility characteristics of people displaying excessive medial knee displacement. *Archives of physical medicine and rehabilitation* 2008, 89(7):1323-1328.
101. Claiborne TL, Armstrong CW, Gandhi V, Pincivero DM: Relationship between hip and knee strength and knee valgus during a single leg squat. *Journal of applied biomechanics* 2006, 22(1):41-50.

102. Hollman JH, Galardi CM, Lin IH, Voth BC, Whitmarsh CL: Frontal and transverse plane hip kinematics and gluteus maximus recruitment correlate with frontal plane knee kinematics during single-leg squat tests in women. *Clinical biomechanics* (Bristol, Avon) 2014, 29(4):468-474
103. Brown TN, McLean SG, Palmieri-Smith RM: Associations between lower limb muscle activation strategies and resultant multi-planar knee kinetics during single leg landings. *Journal of science and medicine in sport / Sports Medicine Australia* 2013, 17 (4):408-413.
104. Dill KE, Begalle R, Frank B, Zinder S, Padua DA: Altered knee and ankle kinematics during squatting in those with limited weight-bearing lunge ankle-dorsiflexion range of motion. *Journal of athletic training* 2014, 49(6):723-732
105. Baldon Rde M, Lobato DF, Furlan L, Serrao FV: Gender differences in lower limb kinematics during stair descent. *Journal of applied biomechanics* 2013, 29(4):413-420.
106. Tanikawa H, Matsumoto H, Komiyama I, Kiriya Y, Toyama Y, Nagura T: Comparison of Knee Mechanics Among Risky Athletic Motions for Non-Contact Anterior Cruciate Ligament Injury. *Journal of applied biomechanics* 2013.
107. Graci V, Van Dillen LR, Salsich GB: Gender differences in trunk, pelvis and lower limb kinematics during a single leg squat. *Gait & posture* 2012, 36(3):461-466.
108. Sigward SM, Pollard CD, Havens KL, Powers CM: Influence of sex and maturation on knee mechanics during side-step cutting. *Medicine and science in sports and exercise* 2012, 44(8):1497-1503.
109. Boling M, Padua D: Relationship between hip strength and trunk, hip, and knee kinematics during a jump-landing task in individuals with patellofemoral pain. *International journal of sports physical therapy* 2013, 8(5):661-669.
110. Willson JD, Kernozek TW, Arndt RL, Reznichuk DA, Scott Straker J: Gluteal muscle activation during running in females with and without patellofemoral pain syndrome. *Clinical biomechanics* (Bristol, Avon) 2011, 26(7):735-740.
111. Tegner Y, Lysholm J: Rating systems in the evaluation of knee ligament injuries. *Clinical orthopaedics and related research* 1985(198):43-49.
112. Munn J, Sullivan SJ, Schneiders AG: Evidence of sensorimotor deficits in functional ankle instability: a systematic review with meta-analysis. *Journal of science and medicine in sport / Sports Medicine Australia* 2010, 13(1):2-12.
113. Lephart S, Fu F: *Proprioception and Neuromuscular Control in Joint Stability* Champaign: Human Kinetics; 2000.
114. Ageberg E, Flenhagen J, Ljung J: Test-retest reliability of knee kinesthesia in healthy adults. *BMC musculoskeletal disorders* 2007, 8:57.
115. Ageberg E, Björkman A, Rosen B, Roos EM: Principles of brain plasticity in improving sensorimotor function of the knee and leg in patients with anterior cruciate ligament injury: A double-blind randomized exploratory trial. *BMC musculoskeletal disorders* 2012, 13(1):68.

116. Farkas GJ, Shakoor N, Cvetanovich GL, Fogg LF, Orias AA, Nho SJ: Vibratory sense deficits in patients with symptomatic femoroacetabular impingement. *Journal of musculoskeletal & neuronal interactions* 2016, 16(1):40-44.
117. Kemp JL, Schache AG, Makdissi M, Sims KJ, Crossley KM: Greater understanding of normal hip physical function may guide clinicians in providing targeted rehabilitation programmes. *Journal of science and medicine in sport / Sports Medicine Australia* 2013, 16(4):292-296.
118. Beutler A, de la Motte S, Marshall S, Padua D, Boden B: Muscle strength and qualitative jump-landing differences in male and female military cadets: the jump-acl study. *Journal of sports science & medicine* 2009, 8:663-671.
119. Pua YH, Wrigley TV, Cowan SM, Bennell KL: Intrarater test-retest reliability of hip range of motion and hip muscle strength measurements in persons with hip osteoarthritis. *Archives of physical medicine and rehabilitation* 2008, 89(6):1146-1154.
120. Muller R, Strassle K, Wirth B: Isometric back muscle endurance: an EMG study on the criterion validity of the Ito test. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2010, 20(5):845-850.
121. McGill SM, Childs A, Liebenson C: Endurance times for low back stabilization exercises: clinical targets for testing and training from a normal database. *Archives of physical medicine and rehabilitation* 1999, 80(8):941-944.
122. Cronström A, Creaby M, Blackmore T, Nae J, Ageberg E: Is trunk and lower extremity strength associated with knee abduction angle during weight-bearing activities in patients with anterior cruciate ligament reconstruction? In manuscript 2016.
123. Hermens H, Freriks B, Disselhorst-Klug C, Rau G: Development of recommendations for SEMG sensors and sensor placement procedures. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2000, 10(5):361-374.
124. Smoliga JM, Myers JB, Redfern MS, Lephart SM: Reliability and precision of EMG in leg, torso, and arm muscles during running. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2010, 20(1):e1-9.
125. Cronstrom A, Creaby MW, Nae J, Ageberg E: Gender differences in knee abduction during weight-bearing activities: A systematic review and meta-analysis. *Gait & posture* 2016, 49:315-328.
126. Ageberg E, Bennell KL, Hunt MA, Simic M, Roos EM, Creaby MW: Validity and inter-rater reliability of medio-lateral knee motion observed during a single-limb mini squat. *BMC musculoskeletal disorders* 2010, 11:265.
127. Pfeifer K, Banzer W: Motor performance in different dynamic tests in knee rehabilitation. *Scandinavian journal of medicine & science in sports* 1999, 9(1):19-27.

128. Alkjaer T, Henriksen M, Dyhre-Poulsen P, Simonsen EB: Forward lunge as a functional performance test in ACL deficient subjects: test-retest reliability. *The Knee* 2009, 16(3):176-182.
129. Powell JW, Barber-Foss KD: Sex-related injury patterns among selected high school sports. *The American journal of sports medicine* 2000, 28(3):385-391.
130. Zätterström R, Fridén T, Lindstrand A, Moritz U: Rehabilitation following acute anterior cruciate ligament injuries--a 12-month follow-up of a randomized clinical trial. *Scandinavian journal of medicine & science in sports* 2000, 10(3):156-163.
131. Nae J, Creaby MW, Cronstrom A, Ageberg E: Measurement properties of visual rating of Postural Orientation Errors (POEs) of the lower extremity - A systematic review and meta-analysis. Submitted to *Physical Therapy in Sport* November 2016.
132. Fellin RE, Rose WC, Royer TD, Davis IS: Comparison of methods for kinematic identification of footstrike and toe-off during overground and treadmill running. *Journal of science and medicine in sport / Sports Medicine Australia* 2010, 13(6):646-650.
133. Robertson DG, Dowling JJ: Design and responses of Butterworth and critically damped digital filters. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2003, 13(6):569-573.
134. Grood ES, Suntay WJ: A joint coordinate system for the clinical description of three-dimensional motions: application to the knee. *Journal of biomechanical engineering* 1983, 105(2):136-144.
135. Borenstein M, Hedges L, Higgins J, Rothstein H: *Computing Effect Sizes for Meta-analysis*. Chichester: John Wiley & Sons, Ltd; 2009.
136. *Cochrane Handbook for Systematic Reviews of Interventions* [http://handbook.cochrane.org/chapter_16/16_5_4_how_to_include_multiple_groups_from_one_study.htm]
137. Duval S, Tweedy R: A nonparametric "trim and fill method" of accounting for publication bias in meta-analyses *J Am Stat Assoc* 2000, 95:89-98.
138. Cronström A, Ageberg E: Does sensory function influence medio-lateral knee position in patients with anterior cruciate ligament injury? Master. University of Lund; 2013.
139. Yamazaki J, Muneta T, Ju YJ, Sekiya I: Differences in kinematics of single leg squatting between anterior cruciate ligament-injured patients and healthy controls. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2010, 18(1):56-63.
140. Whatman C, Hume P, Hing W: The reliability and validity of physiotherapist visual rating of dynamic pelvis and knee alignment in young athletes. *Physical therapy in sport : official journal of the Association of Chartered Physiotherapists in Sports Medicine* 2013, 14(3):168-174.
141. Myer GD, Ford KR, Di Stasi SL, Foss KD, Micheli LJ, Hewett TE: High knee abduction moments are common risk factors for patellofemoral pain (PFP) and anterior cruciate ligament (ACL) injury in girls: is PFP itself a predictor for subsequent ACL injury? *British journal of sports medicine* 2015, 49(2):118-122.

142. Walden M, Krosshaug T, Bjorneboe J, Andersen TE, Faul O, Hagglund M: Three distinct mechanisms predominate in non-contact anterior cruciate ligament injuries in male professional football players: a systematic video analysis of 39 cases. *British journal of sports medicine* 2015.
143. Lephart SM RB, Fu FH: Introduction to the sensorimotor system. In: *Proprioception and neuromuscular control in joint stability*, Human kinetics. edn.; 2000: xvii-xviii.
144. Nakagawa TH, Maciel CD, Serrao FV: Trunk biomechanics and its association with hip and knee kinematics in patients with and without patellofemoral pain. *Manual therapy* 2015, 20(1):189-193.
145. Mundermann A, Asay JL, Mundermann L, Andriacchi TP: Implications of increased medio-lateral trunk sway for ambulatory mechanics. *Journal of biomechanics* 2008, 41(1):165-170.
146. Weltin E, Mornieux G, Gollhofer A: Influence of gender on trunk and lower limb biomechanics during lateral movements. *Research in sports medicine (Print)* 2015, 23(3):265-277.
147. Beith ID: Reflex control of ipsilateral and contralateral paraspinal muscles. *Experimental brain research* 2012, 218(3):433-440.
148. Zedka M, Kumar S, Narayan Y: Electromyographic response of the trunk muscles to postural perturbation in sitting subjects. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 1998, 8(1):3-10.
149. Peach JP, Sutarno CG, McGill SM: Three-dimensional kinematics and trunk muscle myoelectric activity in the young lumbar spine: a database. *Archives of physical medicine and rehabilitation* 1998, 79(6):663-669.
150. Beltran L, Ghazikhanian V, Padron M, Beltran J: The proximal hamstring muscle-tendon-bone unit: a review of the normal anatomy, biomechanics, and pathophysiology. *European journal of radiology* 2012, 81(12):3772-3779.
151. Paterno MV, Schmitt LC, Ford KR, Rauh MJ, Myer GD, Huang B, Hewett TE: Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *The American journal of sports medicine* 2010, 38(10):1968-1978.
152. Delp SL, Hess WE, Hungerford DS, Jones LC: Variation of rotation moment arms with hip flexion. *Journal of biomechanics* 1999, 32(5):493-501.
153. Snyder KR, Earl JE, O'Connor KM, Ebersole KT: Resistance training is accompanied by increases in hip strength and changes in lower extremity biomechanics during running. *Clinical biomechanics (Bristol, Avon)* 2009, 24(1):26-34.
154. Herman DC, Weinhold PS, Guskiewicz KM, Garrett WE, Yu B, Padua DA: The effects of strength training on the lower extremity biomechanics of female recreational athletes during a stop-jump task. *The American journal of sports medicine* 2008, 36(4):733-740.
155. Ferber R, Kendall KD, Farr L: Changes in knee biomechanics after a hip-abductor strengthening protocol for runners with patellofemoral pain syndrome. *Journal of athletic training* 2011, 46(2):142-149.

156. Palmer K, Hebron C, Williams JM: A randomised trial into the effect of an isolated hip abductor strengthening programme and a functional motor control programme on knee kinematics and hip muscle strength. *BMC musculoskeletal disorders* 2015, 16:105.
157. Nguyen AD, Shultz SJ, Schmitz RJ, Luecht RM, Perrin DH: A preliminary multifactorial approach describing the relationships among lower extremity alignment, hip muscle activation, and lower extremity joint excursion. *Journal of athletic training* 2011, 46(3):246-256.
158. Hollman JH, Hohl JM, Kraft JL, Strauss JD, Traver KJ: Modulation of frontal-plane knee kinematics by hip-extensor strength and gluteus maximus recruitment during a jump-landing task in healthy women. *Journal of sport rehabilitation* 2013, 22(3):184-190.
159. Homan KJ, Norcross MF, Goerger BM, Prentice WE, Blackburn JT: The influence of hip strength on gluteal activity and lower extremity kinematics. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2013, 23(2):411-415.
160. Popovich JM, Jr., Kulig K: Lumbopelvic landing kinematics and EMG in women with contrasting hip strength. *Medicine and science in sports and exercise* 2012, 44(1):146-153.
161. Olsen OE, Myklebust G, Engebretsen L, Bahr R: Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *The American journal of sports medicine* 2004, 32(4):1002-1012.
162. Krosshaug T, Nakamae A, Boden BP, Engebretsen L, Smith G, Slauterbeck JR, Hewett TE, Bahr R: Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *The American journal of sports medicine* 2007, 35(3):359-367.
163. Koga H, Nakamae A, Shima Y, Iwasa J, Myklebust G, Engebretsen L, Bahr R, Krosshaug T: Mechanisms for noncontact anterior cruciate ligament injuries: knee joint kinematics in 10 injury situations from female team handball and basketball. *The American journal of sports medicine* 2010, 38(11):2218-2225.
164. Myklebust G, Engebretsen L, Braekken IH, Skjølberg A, Olsen OE, Bahr R: Prevention of anterior cruciate ligament injuries in female team handball players: a prospective intervention study over three seasons. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine* 2003, 13(2):71-78.
165. Fox AS, Bonacci J, McLean SG, Spittle M, Saunders N: What is normal? Female lower limb kinematic profiles during athletic tasks used to examine anterior cruciate ligament injury risk: a systematic review. *Sports medicine (Auckland, NZ)* 2014, 44(6):815-832.
166. Gibson J: The ecological approach to the visual perception of pictures. *Leonardo* 1978, 11:227-235.
167. Han J, Anson J, Waddington G, Adams R: Proprioceptive performance of bilateral upper and lower limb joints: side-general and site-specific effects. *Experimental brain research* 2013, 226(3):313-323.

168. Han J, Waddington G, Adams R, Anson J: Ability to discriminate movements at multiple joints around the body: global or site-specific. *Perceptual and motor skills* 2013, 116(1):59-68.
169. Riemann BL, Myers JB, Lephart SM: Sensorimotor system measurement techniques. *Journal of athletic training* 2002, 37(1):85-98.
170. De Luca CJ, Gilmore LD, Kuznetsov M, Roy SH: Filtering the surface EMG signal: Movement artifact and baseline noise contamination. *Journal of biomechanics* 2010, 43(8):1573-1579.
171. Bolgla LA, Uhl TL: Reliability of electromyographic normalization methods for evaluating the hip musculature. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2007, 17(1):102-111.
172. Norcross MF, Blackburn JT, Goerger BM: Reliability and interpretation of single leg stance and maximum voluntary isometric contraction methods of electromyography normalization. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2010, 20(3):420-425.
173. Folland JP, Williams AG: Methodological issues with the interpolated twitch technique. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 2007, 17(3):317-327.
174. Chiari L, Della Croce U, Leardini A, Cappozzo A: Human movement analysis using stereophotogrammetry. Part 2: instrumental errors. *Gait & posture* 2005, 21(2):197-211.
175. Della Croce U, Leardini A, Chiari L, Cappozzo A: Human movement analysis using stereophotogrammetry. Part 4: assessment of anatomical landmark misplacement and its effects on joint kinematics. *Gait & posture* 2005, 21(2):226-237.
176. Leardini A, Chiari L, Della Croce U, Cappozzo A: Human movement analysis using stereophotogrammetry. Part 3. Soft tissue artifact assessment and compensation. *Gait & posture* 2005, 21(2):212-225.
177. Tsai TY, Lu TW, Kuo MY, Lin CC: Effects of soft tissue artifacts on the calculated kinematics and kinetics of the knee during stair-ascent. *Journal of biomechanics* 2011, 44(6):1182-1188.
178. Benoit DL, Damsgaard M, Andersen MS: Surface marker cluster translation, rotation, scaling and deformation: Their contribution to soft tissue artefact and impact on knee joint kinematics. *Journal of biomechanics* 2015, 48(10):2124-2129.
179. Barre A, Jolles BM, Theumann N, Aminian K: Soft tissue artifact distribution on lower limbs during treadmill gait: Influence of skin markers' location on cluster design. *Journal of biomechanics* 2015, 48(10):1965-1971.
180. Cappozzo A, Cappello A, Della Croce U, Pensalfini F: Surface-marker cluster design criteria for 3-D bone movement reconstruction. *IEEE transactions on bio-medical engineering* 1997, 44(12):1165-1174.
181. Milner CE, Westlake CG, Tate JJ: Test-retest reliability of knee biomechanics during stop jump landings. *Journal of biomechanics* 2011, 44(9):1814-1816.

182. Ford KR, Myer GD, Hewett TE: Reliability of landing 3D motion analysis: implications for longitudinal analyses. *Medicine and science in sports and exercise* 2007, 39(11):2021-2028.
183. Whatman C, Hume P, Hing W: Kinematics during lower extremity functional screening tests in young athletes - are they reliable and valid? *Physical therapy in sport : official journal of the Association of Chartered Physiotherapists in Sports Medicine* 2013, 14(2):87-93.
184. Miller E, Kaufman K, Kingsbury T, Wolf E, Wilken J, Wyatt M: Mechanical testing for three-dimensional motion analysis reliability. *Gait & posture* 2016, 50:116-119.
185. Donnell-Fink LA, Klara K, Collins JE, Yang HY, Goczalk MG, Katz JN, Losina E: Effectiveness of Knee Injury and Anterior Cruciate Ligament Tear Prevention Programs: A Meta-Analysis. *PloS one* 2015, 10(12):e0144063.
186. Sugimoto D, Myer GD, Foss KD, Hewett TE: Specific exercise effects of preventive neuromuscular training intervention on anterior cruciate ligament injury risk reduction in young females: meta-analysis and subgroup analysis. *British journal of sports medicine* 2015, 49(5):282-289.
187. Pappas E, Nightingale EJ, Simic M, Ford KR, Hewett TE, Myer GD: Do exercises used in injury prevention programmes modify cutting task biomechanics? A systematic review with meta-analysis. *British journal of sports medicine* 2015, 49(10):673-680.
188. Zebis MK, Andersen LL, Brandt M, Myklebust G, Bencke J, Lauridsen HB, Bandholm T, Thorborg K, Holmich P, Aagaard P: Effects of evidence-based prevention training on neuromuscular and biomechanical risk factors for ACL injury in adolescent female athletes: a randomised controlled trial. *British journal of sports medicine* 2016, 50(9):552-557.
189. Macrum E, Bell DR, Boling M, Lewek M, Padua D: Effect of limiting ankle-dorsiflexion range of motion on lower extremity kinematics and muscle-activation patterns during a squat. *Journal of sport rehabilitation* 2012, 21(2):144-150.
190. Tiberio D: Pathomechanics of structural foot deformities. *Physical therapy* 1988, 68(12):1840-1849.
191. Joseph M, Tiberio D, Baird JL, Trojian TH, Anderson JM, Kraemer WJ, Maresh CM: Knee valgus during drop jumps in national collegiate athletic association division i female athletes: the effect of a medial post. *The American journal of sports medicine* 2008, 36(2):285-289.
192. Pietrosimone BG, McLeod MM, Lepley AS: A theoretical framework for understanding neuromuscular response to lower extremity joint injury. *Sports health* 2012, 4(1):31-35.
193. Grooms DR, Myer GD: Upgraded hardware horizontal line What about the software? Brain updates for return to play following ACL reconstruction. *British journal of sports medicine* 2016.
194. Lebon F, Guillot A, Collet C: Increased muscle activation following motor imagery during the rehabilitation of the anterior cruciate ligament. *Applied psychophysiology and biofeedback* 2012, 37(1):45-51.

195. Shultz SJ, Schmitz RJ: Tibial plateau geometry influences lower extremity biomechanics during landing. *The American journal of sports medicine* 2012, 40(9):2029-2036.
196. Nguyen AD, Shultz SJ, Schmitz RJ: Landing Biomechanics in Participants With Different Static Lower Extremity Alignment Profiles. *Journal of athletic training* 2015.