



# LUND UNIVERSITY

Eilifdahlia schwarzii (Caloplacoideae, Teloschistaceae) – a new species from Western Australia

Kondratyuk, Sergey. Y.; Schumm, Felix; Elix, John A.; Kärnefelt, Ingvar; Thell, Arne; Hur, Jae-Seoun

*Published in:*  
Graphis Scripta

2017

*Document Version:*  
Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for published version (APA):*

Kondratyuk, S. Y., Schumm, F., Elix, J. A., Kärnefelt, I., Thell, A., & Hur, J.-S. (2017). Eilifdahlia schwarzii (Caloplacoideae, Teloschistaceae) – a new species from Western Australia. *Graphis Scripta*, 29(1-2), 18-23.

*Total number of authors:*  
6

## General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

## Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00



## ***Eilifdahlia schwarzii* (Caloplacoideae, Teloschistaceae) – a new species from Western Australia**

SERGII Y. KONDRATYUK, FELIX SCHUMM, JOHN A. ELIX, INGVAR KÄRNEFELT, ARNE THELL and JAE-SEOUN HUR

Kondratyuk, S. Y., Schumm, F., Elix, J. A., Kärnefelt, I., Thell, A. & Hur, J.-S. 2017. *Eilifdahlia schwarzii* (Caloplacoideae, Teloschistaceae) – a new species from Western Australia. *Graphis Scripta* **29**(1–2): 18–23. Tartu. ISSN 0901-7593.

The new species *Eilifdahlia schwarzii* F. Schumm & S. Y. Kondr. is described. It differs from the more common southern Australian *E. dahlia* (Elix, S. Y. Kondr. & Kärnefelt) S. Y. Kondr., Kärnefelt, Elix, A. Thell, J. Kim, A. S. Kondr. & J.-S. Hur in having a poorly developed whitish thallus, bright yellow to bright yellow-orange apothecia, with a persistent glossy margin, a proper exciple with scleroplectenchymatous tissue, longer ascospores, and wider ascospore septa. The new species is known from scattered localities in Western Australia. The new combination *E. sergeyana* (Kantvilas) S. Y. Kondr., Elix, Kärnefelt & A. Thell is made, and a key to the species of *Eilifdahlia* is presented.

Key words: New species, new combination, ecology, lichen substance, key

*Sergii Y. Kondratyuk, M. H. Kholodny Institute of Botany, Tereshchenkivska str. 2, 01004 Kiev, Ukraine; E-mail: ksya\_net@ukr.net; Korean Lichen Research Institute, Suncheon National University, 315 Mae-gok dong, Suncheon, 540-742, Korea.*

*Felix Schumm, Mozartstr. 9, 73221, Wangen, Germany. E-mail: fschumm@online.de*

*John A. Elix, Research School of Chemistry, Building 137, Australian National University, Canberra, ACT 2601, Australia. E-mail: john.elix@anu.edu.au*

*Ingvar Kärnefelt, Biological Museum, Lund University, Box 117, SE-221 00 Lund, Sweden. E-mail: ingvar.karnefelt@biol.lu.se*

*Arne Thell, Biological Museum, Lund University, Box 117, SE-221 00 Lund, Sweden. E-mail: arne.thell@biol.lu.se*

*Jae-Seoun Hur, Korean Lichen Research Institute, Suncheon National University, 315 Mae-gok dong, Suncheon, 540-742, Korea. E-mail: jshur1@sunchon.ac.kr*

### **Introduction**

The genus *Eilifdahlia* comprises four species, *E. dahlia*, *E. sergeyana*, *E. schwarzii* and *E. wirthii*. Two of these species, the type of the genus *E. dahlia* as well as *E. wirthii* have been included in phylogenetic analyses which confirmed that *Eilifdahlia* was close to *Franwilsia* (Kondratyuk *et al.* 2014). The other two species will be investigated in the near future (Kondratyuk *et al.* in prep.).

*Eilifdahlia* S. Y. Kondr., Kärnefelt, Elix, A. Thell & J.-S. Hur occurs primarily in the Southern Hemisphere and was first described in 2014. The genus parallels *Blastenia* A. Massal., primarily distributed in the Northern Hemisphere. In the original description of *Eilifdahlia* two species were included, namely *E. dahlia* (Elix, S. Y. Kondr. & Kärnefelt) S. Y. Kondr., Kärnefelt,

Elix, A. Thell, J. Kim, A. S. Kondr. & J.-S. Hur, which is widely distributed in southern Australia, and *E. wirthii* S. Y. Kondr., known from several localities in South Africa (Kondratyuk *et al.* 2014).

The genus *Eilifdahlia* was supported by a three gene phylogeny based on separate and combined analyses of ITS1/ITS2 nrDNA, 28S nrLSU and 12S mtSSU sequences (Kondratyuk *et al.* 2014). The genus is most similar to *Franwilsia* S. Y. Kondr., Kärnefelt, Elix, A. Thell & J.-S. Hur, and in the combined analysis they formed a subclade within the subfamily Caloplacoideae (Kondratyuk *et al.* 2014). In addition to chemical differences, the *Franwilsia* is distinguished in having a hymenium interspersed with oil droplets.

Another species, *Caloplaca sergeyana* Kantvilas, recently described from siliceous rocks in South Australia, is similar to *E. dahlia* (Kantvilas 2016). Here it is combined into *Eilifdahlia*. During a revision of lichen collections from Western Australia by the second author (FS) an additional species was discovered, which forms the subject of this paper. A key to the four species of *Eilifdahlia* is presented.

## Material and Methods

A total of more than 100 specimens collected during last decade and deposited in B and HO, as well as some duplicates in the Korean Lichen Research Institute, Sunchon National University, South Korea (KoLRI), the Hungarian Natural History Museum (BP) and the Lichen Herbarium in the M. H. Kholodny Institute of Botany of National Academy of Sciences of Ukraine (KW-L), have been examined using standard microscopical techniques and hand-sectioned under a dissecting microscope (Nikon SMZ 645). Anatomical characters were observed using a Nikon Eclipse E200 microscope and a Zeiss Scope, complemented with a digital camera AxioCam ERc 5s, as well as in Wanger using Wild M3 and Olympus BX51 DIK microscopes and Canon Eos 600D with Canon MP-E 65 mm Macro Photo and Kryostat Microm HM 560. Sections of apothecia were tested with water, K and IKI (10% potassium iodide).

## The New Species

*Eilifdahlia schwarzi* F. Schumm & S. Y. Kondr. **sp. nov.**

MycoBank 819920

Similar to *E. dahlia* but differs in having a poorly developed, whitish thallus, bright yellow to yellow-orange apothecia, with a persistent, glossy proper margin, a proper exciple of scleroplectenchymatous tissue, longer ascospores and wider ascospore septa.

**Type:** Australia. Western Australia, Yalgorup National Park, Lake Clifton, 32°44'43"S 115°39'20.9"E, 15 m alt., on bark, growing together with *Buellia dissa* (Stirt.) Zahlbr. and species of *Physcia*, *Opegrapha*, *Candelariella* and *Arthonia*, 28.x.2009, F. Schumm & E. Stocker-Wörgötter (PERTH, ex herb. F. Schumm 15956 – holotype); the same locality, growing together with *Physcia* sp., *Buellia dissa* and *Kaerneftia kaernefeltii* (S. Y. Kondr., Elix & A. Thell) S. Y. Kondr. *et al.* (B ex herb. F. Schumm 15937 – isotype); the same locality, growing together with *Buellia* sp. and *Opegrapha* sp. (B ex herb. F. Schumm 15951 – isotype).

Thallus corticolous, up to 3 cm wide or forming aggregations, very thin, continuous to finely granular or warty, white to dirty white or whitish-grey, covered with numerous bright yellow or yellow-orange apothecia with a glossy, well developed, pure yellow proper margin and sparse to numerous, dull orange to bluish-orange or bluish-black conidiomata.

Apothecia 0.5–1.3 mm diam., numerous, dispersed, biatorine (without algae), initially regularly rounded but becoming undulate at the margin. The proper margin distinct, persistent, 0.08–0.14(–0.15) mm wide, raised above the disc, usually somewhat shiny, bright yellow to dull yellow-brown or dull yellow-orange (somewhat paler than the disc). In section 0.2–0.25(–0.35) mm thick, biatorine, proper exciple (40–)50–70 µm thick in the uppermost lateral portion, 80–90 µm thick in the lower lateral and basal portions, palisade plectenchymatous in the outer basal portion, the inner portions scleroplectenchymatous with lumina to 1–1.5 µm diam., the outermost part of the lateral portion to 30 µm thick, brownish-orange, the inner portion hyaline; hymenium 60–70 µm high; epihymenium dark brownish-orange in contrast to the pale yellow-brown outer layer of the proper exciple; paraphyses slightly broader at apices, to 3.5 µm wide in K; subhymenium 50–60(–70) µm thick, dull yellow to dull brownish-yellow in lower portion; ascospores rather long and with long septa, (13–)16–20.5(–22) × 7–11(–12) µm in water and (15–)16–21(–22) × (7–)9–12(–14) µm in K, septa (4–)7–9 µm wide in water and (8–)9–12(–14) µm wide in K. Conidiomata dull orange, bluish-orange or blackish-orange, to 0.05–0.07 mm diam.; conidia narrow bacilliform, 2.5–3.5 × 0.7–0.8 µm.

*Chemistry*: Thallus UV+ yellow, parietin and lichexanthone only present in the apothecia.

*Ecology*: *Eilifdahlia schwarzii* grows on the bark of various trees, together with *Kaernefia kaernefeltii*, *Buellia dissa* and species of *Arthonia*, *Candelariella*, *Opegrapha* and *Physcia*. It sometimes overgrows thalli and even ascomata of other crustose lichens [e.g. specimens 15951 and 15912 where *E. schwarzii* grows over ascomata of *Opegrapha* sp.].

*Etymology*: The new species is named in honour of the German bryologist Uwe Schwarz (17.V.1965–) a friend, colleague and keen field-trip supporter of the second author.

*Distribution*: *Eilifdahlia schwarzii* is only known from several localities in Western Australia.

*Taxonomic notes*: *Eilifdahlia schwarzii* is similar to two other Australian species, the common, corticolous *E. dahlia* and the rare, epilithic *E. sergeyana*; neither species have interspersed hymenia. Both *E. schwarzii* and *E. dahlia* have a brownish-yellow lower subhymenium, but *E. schwarzii* differs in having a membranous thallus in contrast to the thick, well developed, continuous and ± wrinkled thallus of *E. dahlia*. The thallus of *E. schwarzii* varies from white to dirty white or whitish-grey whereas *E. dahlia*, varies from yellow-grey to brownish-grey. Furthermore, *E. schwarzii* has bright yellow to yellow-orange apothecia, whereas the discs are brownish-orange to dark brown in *E. dahlia* (Table 1). The proper margin of the apothecia is persistent and glossy in *E. schwarzii* but becomes excluded in *E. dahlia*. *E. schwarzii* is also characterized by a scleroplectenchymatous proper exciple rather than pseudoprosoplectenchymatous as in *E. dahlia* (Kondratyuk & Kärnefelt 1997). The ascospores are longer with wider septa in *E. schwarzii*, the hymenium is lower and the conidia narrower (Table 1). Finally, the two species differ chemically, since *E. schwarzii* lacks lichexanthone in the thallus, whereas this substance is abundant in *E. dahlia*.

The epilithic *E. sergeyana* lacks lichexanthone in the thallus, but differs in having a very thin, more or less continuous thallus, smaller apothecia and shorter ascospores with narrower septa (Kantvilas 2016).

The epiphytic *E. wirthii* is distinguished from *E. schwarzii* by its thicker thallus, smaller, darker and pruinose discs, 0.2–0.8 mm diam., a bluish black proper exciple, an entirely brownish subhymenium, smaller ascospores, (8–)9–11 × 4–5(–8) µm, with narrower septa, (1–)3–4 µm wide, and slightly broader conidia, 3–3.5 × 0.7–1.2 µm (Kondratyuk *et al.* 2014).

In some respects *E. schwarzii* resembles *Franwilsia kilcundaensis* (S. Y. Kondr. & Kärnefelt) S. Y. Kondr. *et al.*, which is characterized by its brownish-grey thallus containing lichexanthone, but this species differs from *E. schwarzii* by having rusty orange-brown discs with a pale grey thalline margin, and a densely oil-inspersed hymenium and subhymenium (Kondratyuk *et al.* 2009b, Kantvilas 2016).

*Additional specimen examined:* Australia: Western Australia, Two Peoples Bay east of Albany, 34°58'10.8"S 118°10'44.4"E, 6 m alt., on bark, growing together with species of *Opegrapha* and *Megalospora*, 27.x.2009 F. Schumm & E. Stocker-Wörgötter (B, ex herb. F. Schumm 15912 – paratype).



**Figure 1.** *Eilifdahlia schwarzii* (HOLOTYPE), general habit. Photo: Felix Schumm.

### **New combination**

*Eilifdahlia sergeyana* (Kantvilas) S. Y. Kondr., Elix, Kärnefelt & A. Thell, **comb. nov.**

MycoBank 819921

Basionym: *Caloplaca sergeyana* Kantvilas, Journal of the Adelaide Botanic Gardens 29: 57 (2016).

**Table 1.** A comparison of *E. schwarzii* and related species.

Character	<i>Eilifdahlia schwarzii</i>	<i>Eilifdahlia dahliei</i>	<i>Eilifdahlia sergeyana</i>
<b>Thallus development</b>	Poorly developed	Distinct	Thin but distinct
<b>Thallus colour</b>	Whitish	Yellow-grey to brownish grey or grey-green	Greenish grey to brownish grey
<b>Apothecial colour, disc</b>	Yellow-orange	Dull brownish orange to dark brown	Orange to yellow-orange
<b>Apothecial width, mm</b>	0.5–1.3	0.4–0.9(–2.5)	0.5–0.8(–1)
<b>Apothecial proper margin</b>	Persistent and glossy	Indistinct and excluded, dull	Initially inrolled, with age slightly flexuose, persistent
<b>Tissue of true exciple</b>	Scleroplectenchymatous	Pseudoprosoplectenchymatous	Of radiating, parallel, anastomosing hyphae
<b>Ascospore form and size, µm</b>	Oblong, (13–)16–20.5(–22) × 7–11(–12)	Subspherical, (12–)13–16(–19) × (6–)7–9(–10) after Kondratyuk <i>et al.</i> (2009a; 12–17 × 5–9 according to Kantvilas (2016)	Oblong, (11–)11.5–16(–17) × (5–)6–8(–8.5)
<b>Ascospore septum µm</b>	(4–)7–9	(4–)6–7(–9)	3–7(–9)
<b>Hymenium height, µm</b>	60–70	80–90	70–80
<b>Conidia width, µm</b>	0.8–0.9	0.9–1.0	unknown
<b>Lichexanthone</b>	Present in apothecia; absent in thallus	Abundant in both apothecia and thallus	Absent in both apothecia and thallus
<b>Distribution</b>	Scattered in Western Australia	Widespread in southern Australia	Known only from type collection in South Australia

### Key to the species of *Eilifdahlia*

- 1 On rock ..... *E. sergeyana*
- On bark of trees or on lignum ..... 2
- 2. Disc dark yellow-brown to blackish brown with yellowish pruina; proper margin dark brown or reddish brown; ascospores 8–12 × 4–5 µm ..... *E. wirthii*
- Disc dull orange or brownish orange, epruinose; proper margin bright yellow to dull yellow-orange; ascospores 13–21 × 5–11 µm ..... 3
- 3 Ascospores 12–16 µm long; septum to 6 µm wide; proper margin initially present, then excluded ..... *E. dahliei*
- Ascospores 16–21 µm long; septum 7–12 µm wide; proper margin persistent ..... *E. schwarzii*

## Acknowledgements

Sergii Kondratyuk is grateful to Dr Jae-Seoun Hur (Sunchon, South Korea) for his kind assistance during his stay at the KoLRI in 2015 and 2016 and during the field work. SK was supported financially by The *Ministry of Education and Science of Ukraine* (M/90-2015 and M/34-2016) and by the Korean Brain Pool Program (161S-4-3-1659). The Almborn Foundation is thanked for financial support.

## References

- Kantvilas, G. 2016. A synopsis and key for the lichen genus *Caloplaca* (Teloschistaceae) on Kangaroo Island, with the description of two new species. *Journal of the Adelaide Botanic Gardens* **29**: 53–69.
- Kondratyuk, S. Y. & Kärnefelt, I. 1997. *Josefpoeltia* and *Xanthomendoza* two new genera in the family Teloschistaceae (Ascomycotina). *Bibliotheca Lichenologica* **68**: 19–44.
- Kondratyuk, S. Y., Kärnefelt, I., Elix, J. A. & Thell, A. 2007. Contributions to the Teloschistaceae of Australia. *Bibliotheca Lichenologica* **96**: 157–174.
- Kondratyuk, S. Y., Kärnefelt, I., Elix, J. A. & Thell, A. 2009a. New *Caloplaca* species from Australia. *Bibliotheca Lichenologica* **99**: 259–278.
- Kondratyuk, S. Y., Kärnefelt, I., Elix, J. A. & Thell, A. 2009b. Contributions to the Teloschistaceae, with particular reference to the Southern Hemisphere. In: Thell, A., Seaward, M. R. D. & Feurer, T. (eds), *Diversity of Lichenology – Anniversary Volume*. *Bibliotheca Lichenologica* **100**: 207–282.
- Kondratyuk, S. Y., Jeong, M.-H., Yu, N.-N., Kärnefelt, I., Thell, A., Elix, J. A., Kim, J., Kondratiuk, A. S. & Hur, J.-S. 2014. A revised taxonomy for the subfamily Caloplacoideae (Teloschistaceae, Ascomycota) based on molecular phylogeny. *Acta Botanica Hungarica* **56**: 93–123.