



# LUND UNIVERSITY

## Evolutionary genomics of host-microbe interactions

Videvall, Elin

2018

[Link to publication](#)

*Citation for published version (APA):*

Videvall, E. (2018). *Evolutionary genomics of host-microbe interactions*. Lund University, Faculty of Science, Department of Biology.

*Total number of authors:*

1

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

The background is a stylized illustration of a savanna landscape. In the foreground, there are dark blue silhouettes of bushes and a large acacia tree on the right. In the middle ground, an ostrich and two smaller birds are visible. The birds are dark brown with three yellow, circular, patterned spots on their chests, representing microbes. The background features rolling yellow hills and blue mountains under a solid orange sky. White branches with small white berries or flowers are scattered across the left side.

# Evolutionary genomics of host-microbe interactions

ELIN VIDEVALL

DEPARTMENT OF BIOLOGY | FACULTY OF SCIENCE | LUND UNIVERSITY



# Evolutionary genomics of host-microbe interactions

Elin Videvall





# Evolutionary genomics of host-microbe interactions

Elin Videvall



**LUND**  
UNIVERSITY

DOCTORAL DISSERTATION

by due permission of the Faculty of Science, Lund University, Sweden.  
To be defended in the Blue Hall, Ecology Building, Sölvegatan 37, Lund, Sweden  
on Friday 6 April 2018, 13.00.

*Faculty opponent*

Dr. Susan Perkins

Curator and Professor at the Sackler Institute for Comparative Genomics,  
American Museum of Natural History, New York, NY, USA.

<b>Organization</b> LUND UNIVERSITY  Department of Biology Sölvegatan 37, 223 62 Lund, Sweden		<b>Document name</b> DOCTORAL DISSERTATION	
		<b>Date of issue</b>	
<b>Author(s)</b> Elin Videvall		<b>Sponsoring organization</b>	
<b>Title</b> Evolutionary genomics of host-microbe interactions			
<b>Abstract</b> <p>The microbes living inside hosts have highly important consequences for host health and fitness. From the host's perspective, some microbes exhibit mutualistic tendencies, others parasitic, and some commensal, but this is context-dependent and opportunistic lifestyles are widespread in nature. Our knowledge of how hosts interact molecularly with different microbes is, however, poor, and little research has been done on non-model organisms from a genomic and community-wide perspective. In this PhD thesis, I investigate host-microbe interactions from multiple angles, and utilize high-throughput sequencing techniques to paint a broad, overarching picture of the relationship between hosts and microbes.</p> <p>My PhD comprised two related projects, 1) host-microbiome interactions and 2) host-parasite interactions. In the former, I have evaluated how to best sample and measure the gut microbiomes of avian hosts (Paper I and II). Different sections of the ostrich gastrointestinal tract were characterized and shown to harbour divergent microbial communities (Paper I, II, and IV). I have further demonstrated that the gut microbiome of juvenile ostriches is colonized in a successional manner and gradually develops over time (Paper III), and is strongly linked to growth and mortality (Paper III and IV). In the second project I described the avian transcriptome response to malaria infection over time and to parasites with different virulence (Paper V and VI). Birds with malaria infection experience a range of transcriptional changes that involves for example the immune system, stress response, cell death regulation, and regulatory genes. To evaluate the molecular response of the malaria parasite, I assembled the blood transcriptome of <i>Plasmodium ashfordi</i> and showed that parasite gene expression is host-specific (Paper VII). This transcriptome was subsequently used, together with a genome assembly of <i>Haemoproteus tartakovskyi</i>, to construct a phylogeny of haemosporidian parasites which showed strong support for a monophyletic clade of mammalian malaria parasites (Paper VIII). Finally, the assembled transcriptome and genome were utilized to identify thiamine biosynthesis enzymes in avian <i>Plasmodium</i> (Paper IX), and to demonstrate that the avian <i>Plasmodium</i> parasites exhibit the most AT-rich genes of eukaryotes (Paper X).</p> <p>In summary, this work offers new insights into host-microbiome and host-parasite interactions, and enables a greater understanding of the multifaceted relationship between hosts and their microbes.</p>			
<b>Key words</b> host, microbiota, avian malaria, ostrich, parasite, dual RNA-seq			
<b>Classification system and/or index terms (if any)</b>			
<b>Supplementary bibliographical information</b>		<b>Language</b> English	
<b>ISSN and key title</b>		<b>ISBN</b> 978-91-7753-577-5 (print) 978-91-7753-578-2 (pdf)	
<b>Recipient's notes</b>		<b>Number of pages</b> 297	<b>Price</b>
		<b>Security classification</b>	

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature 

Date 27 February 2018

# Evolutionary genomics of host-microbe interactions

Elin Videvall



**LUND**  
UNIVERSITY

Copyright Elin Videvall

Cover illustration by Maj Persson and Anna Persson

Faculty of Science  
Department of Biology

ISBN (print) 978-91-7753-577-5

ISBN (pdf) 978-91-7753-578-2

Printed by Media-Tryck, Lund University  
Lund, Sweden 2018



MADE IN SWEDEN 

Media-Tryck is an environmentally  
certified and ISO 14001 certified  
provider of printed material.  
Read more about our environmental  
work at [www.mediatryck.lu.se](http://www.mediatryck.lu.se)

*We have never been individuals – We are all lichens*

Scott F. Gilbert

# Table of contents

List of papers .....	10
Author contributions .....	12
Additional papers published during PhD .....	14
Abstract .....	15
Svensk sammanfattning .....	16
Glossary .....	18
Introduction .....	19
Background .....	21
Host-microbiome interactions .....	21
A world without microbes .....	21
Microbiota in health and disease .....	22
Factors affecting host microbiota .....	24
The importance of microbiota in evolutionary processes .....	26
Unresolved questions in host-microbiome research .....	27
Host-parasite interactions .....	29
Host coevolution with parasites .....	29
Host defence .....	30
Parasite virulence .....	32
Immune evasion by parasites .....	32
Controlled host-parasite experiments .....	34
Unresolved questions in host-parasite research .....	34
Genomics .....	36
High-throughput sequencing of microbiomes .....	36
High-throughput sequencing of host responses .....	37
High-throughput sequencing of parasites .....	38
Aim of thesis .....	39
General methodology .....	40
Ostrich gut microbiomes .....	40
Siskin malaria .....	42
Results and discussion .....	45

Measuring the gut microbiome in birds .....	45
Development of gut microbiota in juvenile ostriches .....	47
Strong link between gut microbiota and mortality in ostriches .....	49
Avian transcriptome responses to malaria parasites .....	51
Malaria parasite transcriptome responses to hosts .....	53
Evolution of malaria parasite genomes .....	55
References .....	57
Acknowledgements .....	66
Thank you .....	67
Paper I.....	75
Paper II .....	99
Paper III.....	121
Paper IV .....	149
Paper V .....	179
Paper VI.....	199
Paper VII .....	223
Paper VIII .....	251
Paper IX.....	273
Paper X .....	283

# List of papers

- I. **Elin Videvall**, Maria Strandh, Anel Engelbrecht, Schalk Cloete, Charlie K Cornwallis. (2017) Measuring the gut microbiome in birds: Comparison of faecal and cloacal sampling. *Molecular Ecology Resources*. doi: 10.1111/1755-0998.12744.
- II. **Elin Videvall**, Maria Strandh, Anel Engelbrecht, Schalk Cloete, Charlie K Cornwallis. (2017) Direct PCR offers a fast and reliable alternative to conventional DNA isolation methods for gut microbiomes. *mSystems*, 2: e00132-17.
- III. **Elin Videvall**, Se Jin Song, Hanna M Bensch, Maria Strandh, Anel Engelbrecht, Naomi Serfontein, Olof Hellgren, Adriaan Olivier, Schalk Cloete, Rob Knight, Charlie K Cornwallis. (2018) The development of gut microbiota in ostriches and its association with juvenile growth. *bioRxiv*. 270017. doi: 10.1101/270017.
- IV. **Elin Videvall**, Se Jin Song, Hanna M Bensch, Maria Strandh, Anel Engelbrecht, Naomi Serfontein, Olof Hellgren, Adriaan Olivier, Schalk Cloete, Rob Knight, Charlie K Cornwallis. (2018) Early-life mortality linked to gut dysbiosis in ostriches. Manuscript.
- V. **Elin Videvall**, Charlie K Cornwallis, Vaidas Palinauskas, Gediminas Valkiūnas, Olof Hellgren. (2015) The avian transcriptome response to malaria infection. *Molecular Biology and Evolution*, 32 (5): 1255–1267.
- VI. **Elin Videvall**, Vaidas Palinauskas, Gediminas Valkiūnas, Olof Hellgren. (2018) Avian transcriptome responses to high- and low-virulent malaria strains over the course of infection. Manuscript.
- VII. **Elin Videvall**, Charlie K Cornwallis, Dag Ahrén, Vaidas Palinauskas, Gediminas Valkiūnas, Olof Hellgren. (2017) The transcriptome of the avian malaria parasite *Plasmodium ashfordi* displays host-specific gene expression. *Molecular Ecology*, 26 (11): 2939–2958.



- VIII. Staffan Bensch, Björn Canbäck, Jeremy D DeBarry, Tomas Johansson, Olof Hellgren, Jessica C Kissinger, Vaidas Palinauskas, **Elin Videvall**, Gediminas Valkiūnas. (2016) The genome of *Haemoproteus tartakovskyi* and its relationship with human malaria parasites. *Genome Biology and Evolution*, 8 (5): 1361-1373.
- IX. Olof Hellgren, Staffan Bensch, **Elin Videvall**. (2017) *De novo* synthesis of thiamine (vitamin B<sub>1</sub>) is the ancestral state in *Plasmodium* parasites – evidence from avian haemosporidians. *Parasitology*. doi: 10.1017/S0031182017002219.
- X. **Elin Videvall**. (2018) *Plasmodium* parasites of birds have the most AT-rich genes of eukaryotes. *Microbial Genomics*, 4 (2): 1–9. doi: 10.1099/mgen.0.000150.

Paper I, VII, and IX have been reprinted with permission from the publishers.

# Author contributions

- I. E.V. and C.K.C. planned and designed the study. S.C. provided animal facilities and A.E. supervised the experiment. A.E., C.K.C., and E.V. collected the samples. M.S. planned and performed the laboratory work. E.V. performed the bioinformatic and data analyses. E.V. wrote the paper with assistance of C.K.C. All authors reviewed and approved of the final manuscript.
- II. M.S., E.V., and C.K.C. planned the study. S.C. provided animal facilities, and A.E. supervised the experimental part of the study. M.S. planned and performed the laboratory work. E.V. performed the bioinformatic and data analyses. E.V., M.S., and C.K.C. wrote the paper with input from S.C. and A.E.
- III. E.V. and C.K.C. planned and designed the study. S.C. provided animal facilities. A.E. supervised the experimental part of the study. N.S., A.E., C.K.C., and E.V. performed the sampling and cared for the animals. A.O. advised on sampling procedure. M.S. supervised the laboratorial part of the study, and together with H.M.B. prepared the samples for sequencing. E.V. performed the bioinformatic and data analyses. C.K.C., S.J.S., R.K., and O.H. provided advice on analyses and the interpretation of results. E.V. and C.K.C. wrote the paper with input from all authors.
- IV. E.V. and C.K.C. planned and designed the study. S.C. provided animal facilities. A.E. supervised the experimental part of the study. N.S., A.E., C.K.C., and E.V. performed the sampling and cared for the animals. A.O. performed the euthanization procedure and advised on sampling. M.S. supervised the laboratorial part of the study, and together with H.M.B. prepared the samples for sequencing. E.V. performed the bioinformatic and data analyses. C.K.C., S.J.S., R.K., and O.H. provided advice on analyses and the interpretation of results. E.V. and C.K.C. wrote the paper with input from all authors.
- V. O.H. and G.V. planned and designed the study. V.P. performed the experiment. O.H. performed the RNA extraction. E.V. performed the bioinformatic and data analyses. C.K.C. and O.H. provided advice on data analyses and the inter-

pretation of results. E.V., O.H., and C.K.C. wrote the paper with input from G.V. and V.P.

- VI. O.H. planned and designed the study. G.V. and V.P. performed the experiment. E.V. performed the bioinformatic and data analyses. O.H. provided advice on analyses. E.V. wrote the paper with input from all authors.
- VII. O.H. initially conceived the study design. O.H., V.P., and G.V. further developed the study. G.V. and V.P. planned the experiment. V.P. performed the experiment. O.H. performed the RNA extraction. E.V., O.H., and C.K.C. planned the analyses and interpreted the results. E.V. performed the assembly and the bioinformatic and data analyses. D.A. provided advice on the bioinformatic analyses. E.V. wrote the paper with extensive input from all authors.
- VIII. S.B., O.H., V.P., and G.V. conceived the study. V.P. and G.V. collected the samples. T.J. sequenced the *H. tartakovskyi* genome. B.C. assembled the genome and performed the gene prediction. O.H. organized sequencing of *P. ashfordi* and E.V. performed the transcriptome assembly. E.V., B.C., and S.B. performed genome analyses. J.D.B. and J.C.K. performed the ortholog clustering, the multiple sequence alignments, and the syntenic analyses. B.C. led the phylogenetic analyses with help from J.D.B. and S.B. The paper was written by S.B. with input from the other authors who approved the manuscript before submission.
- IX. O.H. conceived the study, performed the analysis and wrote the first draft of the manuscript. E.V. performed the bioinformatic work on *P. ashfordi* and *P. relictum*. S.B. performed the bioinformatic work on *H. tartakovskyi*. All authors participated in interpreting the data, contributed to the writing and approved of the final manuscript.
- X. E.V. planned the study, analysed the data, and wrote the paper.

# Additional papers published during PhD

Additional papers to which I have devoted significant parts of my time during the PhD. They are not included in this thesis, but are available as open access at the corresponding journal web site.

- XI. Martin N Andersson, **Elin Videvall**, Kimberly KO Walden, Marion O Harris, Hugh M Robertson, Christer Löfstedt. (2014) Sex- and tissue-specific profiles of chemosensory gene expression in a herbivorous gall-inducing fly (Diptera: Cecidomyiidae). *BMC Genomics*, 15 (1): 501.
- XII. **Elin Videvall**, Nina Sletvold, Jenny Hagenblad, Jon Ågren, Bengt Hansson. (2016) Strong maternal effects on gene expression in *Arabidopsis lyrata* hybrids. *Molecular Biology and Evolution*, 33 (4): 984–994.
- XIII. **Elin Videvall**, Erik Öckinger, Lars Pettersson. (2016) Butterfly monitoring using systematically placed transects in contrasting climatic regions – exploring an established spatial design for sampling. *Nature Conservation*, 14: 41–62.
- XIV. Hannah Watson, **Elin Videvall**, Martin N Andersson, Caroline Isaksson. (2017) Transcriptome analysis of a wild bird reveals physiological responses to the urban environment. *Scientific Reports*, 7: 44180.

# Abstract

The microbes living inside hosts have highly important consequences for host health and fitness. From the host's perspective, some microbes exhibit mutualistic tendencies, others parasitic, and some commensal, but this is context-dependent and opportunistic lifestyles are widespread in nature. Our knowledge of how hosts interact molecularly with different microbes is, however, poor, and little research has been done on non-model organisms from a genomic and community-wide perspective. In this PhD thesis, I investigate host-microbe interactions from multiple angles, and utilize high-throughput sequencing techniques to paint a broad, overarching picture of the relationship between hosts and microbes.

My PhD comprised two related projects, 1) host-microbiome interactions and 2) host-parasite interactions. In the former, I have evaluated how to best sample and measure the gut microbiomes of avian hosts (**Paper I and II**). Different sections of the ostrich gastrointestinal tract were characterized and shown to harbour divergent microbial communities (**Paper I, II, and IV**). I have further demonstrated that the gut microbiome of juvenile ostriches is colonized in a successional manner and gradually develops over time (**Paper III**), and is strongly linked to growth and mortality (**Paper III and IV**). In the second project I described the avian transcriptome response to malaria infection over time and to parasites with different virulence (**Paper V and VI**). Birds with malaria infection experience a range of transcriptional changes that involves for example the immune system, stress response, cell death regulation, and regulatory genes. To evaluate the molecular response of the malaria parasite, I assembled the blood transcriptome of *Plasmodium ashfordi* and showed that parasite gene expression is host-specific (**Paper VII**). This transcriptome was subsequently used, together with a genome assembly of *Haemoproteus tartakovskyi*, to construct a phylogeny of haemosporidian parasites which showed strong support for a monophyletic clade of mammalian malaria parasites (**Paper VIII**). Finally, the assembled transcriptome and genome were utilized to identify thiamine biosynthesis enzymes in avian *Plasmodium* (**Paper IX**), and to demonstrate that the avian *Plasmodium* parasites exhibit the most AT-rich genes of eukaryotes (**Paper X**).

In summary, this work offers new insights into host-microbiome and host-parasite interactions, and enables a greater understanding of the multifaceted relationship between hosts and their microbes.

# Svensk sammanfattning

Mikrober finns överallt, runt omkring oss i vår miljö och inuti våra kroppar. De som lever tillsammans med djur har mycket stor påverkan på världens hälsa och evolutionära fitness. Från ett djurs perspektiv är vissa mikrober goda, andra onda, och vissa har ingen större påverkan på hälsan, men egentligen beror dessa egenskaper på sammanhanget, eftersom det är vanligt att mikrober byter strategi till det som passar stunden bäst. Vi har tyvärr väldigt lite kunskap om hur värdar interagerar molekylärt med olika mikrober, och få studier har gjorts ur ett genomiskt perspektiv på djur som inte tillhör de klassiska studieorganismerna. I denna doktorsavhandling undersöker jag interaktioner mellan mikrober och värdar från flera vinklar, och jag använder mig av nya DNA-sekvenseringstekniker för att illustrera en övergripande bild av förhållandet mellan djur och deras mikrober.

Mitt avhandlingsarbete omfattar två stora projekt, 1) värd-mikrobiom interaktioner och 2) värd-parasit interaktioner. I det första projektet har jag utvärderat hur man bäst kan provta och mäta mikrobiomet i magen hos fåglar (**Kapitel I och II**). Olika delar av mag-tarmkanalen analyserades på DNA och visade sig innehålla samhällen av olika mikrober (**Kapitel I, II och IV**). Jag har sedan kunnat visa att strutsungars mikrobiom koloniserar på ett successivt sätt och gradvis mognar över tid (**Kapitel III**), samt att mikrobiomet är starkt kopplat till både tillväxt och dödlighet (**Kapitel III och IV**). I det andra projektet beskrev jag fåglars transkriptom (totala genuttryck) i respons mot en malaria-infektion över tid och i respons mot parasiter med olika virulens (**Kapitel V och VI**). Fåglar med malariainfektion sätter igång en rad förändringar i sitt genuttryck som involverar till exempel immunsystemet, stressresponsen, regleringen av celledöd och regulatoriska gener. För att utvärdera den molekylära responsen hos malariaparasiter byggde jag ihop transkriptomet av *Plasmodium ashfordi* och kunde visa att genuttryck hos parasiten är specifik beroende på vilken värd den befinner sig i (**Kapitel VII**). Detta transkriptom användes därefter, tillsammans med ett genom av *Haemoproteus tartakovskyi*, för att konstruera ett fylogenetiskt träd av blodparasiter som resulterade i starka bevis för att malariaparasiter som infekterar människor är närmare släkt däggdjursparasiter än fågelparasiter (**Kapitel VIII**). Slutligen använde jag både parasit-transkriptomet samt genomet för att identifiera vitamin B<sub>1</sub>-gener i fågelparasiter (**Kapitel IX**) och för att visa att malariaparasiterna som infekterar fåglar är de eukaryoter som uppvisar de mest AT-rika generna (**Kapitel X**).

Sammanfattningsvis innebär detta arbete nya insikter inom de molekylära interaktioner som äger rum mellan värdar och mikrobiom samt mellan värdar och parasiter. Denna nyvunna kunskap möjliggör en större förståelse för det mångfacetterade förhållandet mellan värdar och deras mikrober.

# Glossary

<b>Microbiota</b>	The community of microorganisms within a confined host environment.
<b>Microbiome</b>	The community of microorganisms <i>and</i> their confined host environment.
<b>Dysbiosis</b>	A state of microbial imbalance resulting from a disturbed microbiota.
<b>Commensal</b>	An organism that lives in/on a host organism and benefits from the partnership, while the host is unaffected.
<b>Mutualist</b>	An organism that lives in/on a host organism and where both partners benefit from the association.
<b>Parasite</b>	An organism that lives in/on a host organism and benefits from the partnership, but at a fitness expense of the host.
<b>Pathogen</b>	An organism that causes disease in a host organism.
<b>Pathobiont</b>	A commensal or mutualist with opportunistic pathogenic potential.
<b>Coevolution</b>	Reciprocal evolution of interacting species.
<b>Holobiont</b>	A unit of biological organization composed of a host and all its associated microorganisms.
<b>Symbiosis</b>	A relationship between two organisms of dissimilar species that is constant and intimate. Can be mutualistic, commensalistic, or parasitic.
<b>Phylosymbiosis</b>	A strong coevolutionary trajectory between several host species and their microbiomes, resulting in a similar topology between the host phylogeny and the microbiome distance dendrogram.
<b>Virulence</b>	The degree of fitness reduction in a host by a given parasite.
<b>Resistance</b>	The capacity of a host to defend itself against parasites.
<b>Tolerance</b>	The capacity of a host to withstand parasites.
<b>Parasitemia</b>	The number of parasites in a host (e.g. proportion infected cells).
<b><i>in vivo</i> research</b>	Experiments using live whole organisms (as opposed to e.g. cell culture).
<b><i>Plasmodium</i></b>	Genus of protist parasites causing the disease malaria in vertebrates.
<b><i>Haemoproteus</i></b>	Genus of protist parasites, closely related to <i>Plasmodium</i> .
<b>Haemosporidian</b>	Protist parasites in the order Haemosporida, of which the majority belong to the genera <i>Plasmodium</i> , <i>Haemoproteus</i> , and <i>Leucocytozoon</i> .
<b>Genome</b>	The entire collection of DNA present in the cell nucleus of an organism.
<b>Transcriptome</b>	The collective transcripts from all expressed genes and their relative expression levels in a given tissue of an organism at a given time point.



# Introduction

We live in a world dominated by microorganisms. All around us are miniscule microbes of various kinds, and inside our bodies we harbour entire ecosystems where diverse microbes coexist (**microbiomes**). The human body contains more microbes than human cells (Sender *et al.* 2016), and it has been estimated that as many as 37 million bacteria and 7 million fungal microbes are added to the indoor air when a person enters a room (Qian *et al.* 2012). Although we have, in modern times, gone to great lengths to exterminate as many microbes as possible from both our bodies and our environment, it has been discovered that we, together with other animals, cannot in fact survive without the many crucial functions performed by microbes.

Some microbes that are associated with hosts have been labelled as harmful (**parasites**), others beneficial (**mutualists**), and many do not affect hosts substantially (**commensalists**). However, as with most living things, it can quickly become difficult to categorize microbial species simply as good or bad, as their effects on hosts are largely contextual and environmentally-dependent. Certain microbes can be beneficial, but when opportunity strikes, they might change strategy to exploit their host (**pathobionts**). Then there are microbes which have been considered purely parasitic for a long time, but new research has illuminated some of their hidden mutualistic properties. A small amount of infectious microbes might even be advantageous, and in some cases necessary, to properly develop into a functional healthy adult (Gensollen *et al.* 2016; Ramanan *et al.* 2016). The famous ‘hygiene hypothesis’ (Strachan 1989) is based on this idea, and states that a lack of exposure to microbes and parasites during early childhood contributes to allergies such as asthma, and chronic inflammatory diseases.

The microbes hosts live with and are exposed to can have enormous consequences on their health and fitness. Several animal species have gone extinct because of the presence of parasitic microbes (van Riper *et al.* 1986; Atkinson & LaPointe 2009), while others may starve or become unable to mature into adulthood if they fail to acquire the correct microbial composition (Hirakawa 2002; Wahl *et al.* 2012). Microorganisms with parasitic tendencies often play major roles in their hosts’ evolutionary trajectory. For example, it has been estimated that the greatest selection pressure exerted on humans in recent times is that from the parasitic microbe which cause malaria (Kwiatkowski 2005; McManus *et al.* 2017). In fact, the evolutionary influence of parasitic microorganism on hosts is so large, that one of the main hypotheses to the long-standing question of why

there are two sexes, implicates the presence of parasites as a causal factor by which hosts select for genetic heterogeneity (Hamilton *et al.* 1990). This **coevolutionary** arms race between hosts and parasites has been popularized as 'The Red Queen hypothesis' (van Valen 1973), derived from the Red Queen's comment to Alice in Lewis Carroll's book 'Through the Looking-Glass': "It takes all the running you can do, to keep in the same place". In a similar fashion, hosts need to constantly evolve to stay in the same place (survive), in response to parasitic microbes.

The genetic basis of the interactions that underlie the coevolution between hosts and their microorganisms was previously studied by examining single candidate genes and using targeted model organisms. For example, the host molecular response could be evaluated by characterizing parts of a specific immune gene, and a particular microbe could be identified by growing a culture. With the advent of high-throughput sequencing, we now have the opportunity to take an unbiased approach to examine the intricate host-microbe relationship. We now know that diverse communities of unculturable microorganisms reside within hosts and that interactions taking place between hosts and microbes are determined by a large and complex network of genes.

In this thesis, I use high-throughput sequencing techniques and genomic tools, to generate a broad but detailed view of the interactions taking place between hosts and their microbes.



**Figure 1.** The author, Elin Videvall (E.V.), at field work in South Africa. Photo: Charlie Cornwallis.

# Background

## Host-microbiome interactions

### **A world without microbes**

Microbes are essential organisms in our world. Already in the 1800's, Louis Pasteur speculated that microbes are necessary for multicellular life (Pasteur 1885).

“Life would not long remain possible in the absence of microbes.”  
— Louis Pasteur

Although this statement is true, to some extent, all life would not suddenly cease to exist in the absence of microbes. Gilbert and Neufeld (2014) speculated that if all bacteria and archaea would suddenly disappear, humans would likely survive, initially, if we can artificially synthesize the essential vitamins and amino acids normally supplied by our gut microbiota. However, many other animals like ruminants and termites would starve to death, plants would rapidly deplete nitrogen levels and die, and the ocean would become virtually dead without bacteria supplying the essential nutrients to support phytoplankton. Most global biogeochemical cycling of nutrients would stop in a world without microbes and the authors predicted that complete human societal collapse would occur within a year (Gilbert & Neufeld 2014). The field of gnotobiology is the study of animals reared in germfree environments, and is built upon the concept of absent microbes. Gnotobiology altered the face of medical research, with a focus that lied within generating “pure units” of biology for sterile experimental research, however some misconceptions about the field has also contributed to a widespread cultural phenomenon revolving around an obsession with eliminating all microorganisms in our environment.

We now know that the microbiota of the gut is absolutely crucial for the development of the vertebrate immune system. Germ-free animals suffer from enlarged caecum, smaller lymph nodes, a poorly developed immune system, and reduced organ sizes, including heart, lungs, and liver (Gordon & Pesti 1971; Macpherson & Harris 2004).


Furthermore, animals without a developed gut microbiota are more susceptible to infection by **pathogenic** bacteria, viruses, and eukaryotes (Sprinz *et al.* 1961; Inagaki *et al.* 1996; Round & Mazmanian 2009), and eliminating bacteria in mice before an influenza infection initiates a reduced immune response, resulting in a higher viral load (Ichinohe *et al.* 2011). Besides modulating the host immune response, microbes in the gut perform a variety of important metabolic and biochemical functions, like metabolism of cholesterol to coprostanol (Sadzikowski *et al.* 1977), and the production of vitamin K (Hill 1997) and essential amino acids (Nicholson *et al.* 2012). Microbes are, as previously mentioned, not only important to vertebrates, but to many other animal hosts. They create biofilms which are essential for a large number of oceanic animals to complete their life cycles (Wahl *et al.* 2012), and a fifth of all insects are so dependent on having bacteria around that they have evolved unique cells, called bacteriocytes, or organs – bacteriomes, where they nurture tens of thousands of their important bacterial symbionts.

## Microbiota in health and disease

Despite the many important roles gut microbiota have for human health, it has been largely understudied until recently, and is therefore sometimes nicknamed “the forgotten organ”. Studies are now finding strong associations between the gut microbiome and multiple diseases such as autoimmune diseases (Macpherson & Harris 2004; Round & Mazmanian 2009), obesity (Turnbaugh *et al.* 2009), depression (Cryan & Dinan 2012; Maes *et al.* 2012), gut-inflammatory diseases, and cancer (Hope *et al.* 2005). Higher taxonomical diversity of the microbial composition in the gut has been associated with health both in humans and other animals (Spor *et al.* 2011). Low diversity of gut bacteria has been linked to disease-related imbalances of the microbial community, termed **dysbiosis** (Turnbaugh *et al.* 2009; Qin *et al.* 2010). However, that does not necessarily mean that a diverse microbiome is healthy and a less diverse is sick. The idea to directly associate a particular gut microbial composition with health has turned out to be very complicated task. The sheer variety of gut microbes, their interactions amongst each other, their interactions with the host, their interactions with macrobiota such as helminths, and the large inter-individual differences in host microbiomes complicate matter tremendously.

A healthy immune system is one that is able to differentiate between microbes, and allow commensal and mutualist microbes to flourish while keeping pathogens and pathobionts in check. To avoid infections by pathogenic strains via food sources, the intestinal immune system constantly monitors the gut microbiota. As many as 70% of all immune cells in the body are located in the gut-associated lymphoid tissue because of this reason (van der Heijden *et al.* 1987; Vighi *et al.* 2008). When pathogenic microbes are detected, specific antibodies called immunoglobulin A (IgA) are released and bind to the intruder, allowing B cells to target them (**Figure 2**). A highly developed

and complex communication system between the gut microbiota and the vertebrate mucosal immune system has evolved in response to the significance microbes have on host health. When parts of this communication fail, autoimmune or autoinflammatory diseases such as inflammatory bowel disease (IBD) may occur (Wu & Wu 2012).



**Figure 2.** Healthy commensal gut microbiota prevents colonization by exogenous pathogens and pathobionts. Reprinted from Kamada *et al.* (2013) with permission from Springer Nature.


Recently, studies have increasingly been able to demonstrate some of the benefits hosts receive from their microbes. ‘Defensive microbes’ are those that provide their host with beneficial mechanisms against pathogens, and they can be divided into two main groups: those with direct effect on pathogens, and those that mediate host responses. Microbes that exhibit direct defensive mechanisms towards pathogens have been shown to produce toxic compounds that kill or reduce pathogen fitness, act as hyperparasites on the pathogens (Tollenaere *et al.* 2014), or directly compete with pathogens for host resources (Mideo 2009; Gerardo & Parker 2014). Microbes that have host-mediated

effects can influence either host resistance or host tolerance (Mideo 2009; Gerardo & Parker 2014; Yilmaz *et al.* 2014). For example, in *Aedes* mosquitoes, the **symbiotic** bacteria *Wolbachia* negatively impact infection by parasitic protozoans and viruses through resource competition and by positively influencing host immune responses (Moreira *et al.* 2009). Another example is the American chestnut tree (*Castanea dentata*), which was under the risk of extinction because of the parasitic fungi *Cryphonectria parasitica*, but was saved with the help of defensive fungal viruses in the family Hypoviridae (MacDonald 1991; Milgroom & Cortesi 2004). In humans, phage therapy using hyperparasitic phage viruses to kill targeted pathogenic bacteria, and faecal transplantation of healthy gut microbiota to treat *Clostridium difficile* infections, have shown great success and are two very promising methods of using defensive microbes to kill pathogens (Gough *et al.* 2011; Ford & King 2016). Mutualistic and defensive microbes will most likely become an important tool for future disease control in both humans and non-human animals, especially in the light of the increase in prevalence of drug-resistant pathogens. Probiotics-based vaccines containing defensive gut microbes are already being considered for certain human diseases, such as malaria (Ngwa & Pradel 2015).

## Factors affecting host microbiota

The composition of the host microbiota can potentially be influenced by both genetic and environmental factors. When it comes to linking host genotype with gut microbiome composition, there has been some conflicting evidence. One of the largest microbiome studies to date investigated the gut microbiomes of 416 twin pairs and showed differentiated microbiota even in homozygotic twins, suggesting that genotype has a minor effect compared to the environment (Goodrich *et al.* 2014). However, the authors found that the effect of host genotype varied across different bacterial families with Christensenellaceae being the most heritable taxon in the human gut. Many other studies investigating the effects of host genotype on gut microbiota have failed, however, to find major contributions (Davenport 2016; Goodrich *et al.* 2016). The general view has therefore resulted in that individual gut microbiome composition in humans is mostly a product of environmental factors. Nevertheless, a few candidate immune genes, for example the major histocompatibility complex (MHC) (Toivanen *et al.* 2001; Bolnick *et al.* 2014; Kubinak *et al.* 2015), toll-like receptor 5 (TLR5) (Vijay-Kumar *et al.* 2010), myeloid differentiation primary response 88 (MYD88) (Turnbaugh *et al.* 2008; Wu & Wu 2012), and nucleotide-binding oligomerization domain containing 2 (NOD2) (Petnicki-Ocwieja *et al.* 2009; Frank *et al.* 2011), have been associated with effects on the gut microbial community. Vijay-Kumar *et al.* (2010) could show that, not only did mice with knocked-out TLR5 become obese and developed metabolic disease, but transplanting their gut microbiota to germ-free mice instilled the same physiological and metabolic changes as the donor mice, suggesting a strong causal effect of the gut

microbiota on host health. Although microbiome studies of laboratory mice and humans contribute much to our understanding, research on the effects of host genotype on microbiota in other animals is crucial for our understanding of this relationship. A handful of studies have evaluated the effects of host phylogenies on gut microbiota, called **phylosymbiosis** (Hird *et al.* 2015; Brooks *et al.* 2016; Groussin *et al.* 2017; Kohl *et al.* 2017). These studies have usually been able to show a minor to moderate effect of coevolution on the gut microbial composition (**Figure 3**), though they suffer from small and biased sampling efforts, with confounding effects such as physiological, ecological, and environmental differences between species. With an increasing number of animal gut microbiomes becoming sequenced, however, future phylosymbiosis research shows a lot of promise, and an increasing amount of data will allow for meta-analyses and more rigorous investigations across species. The outcome of these studies will be highly useful in further evaluating the effect of host genotype on the microbiota.



**Figure 3.** Bird phylogeny (left) compared to a dendrogram of gut microbiota distances (right). Individuals are tracked across the two topologies with lines. Reprinted from Hird *et al.* (2015) under a CC BY 4.0 license.

One of the most prominent drivers behind variation in gut microbiome of hosts is diet. Dramatic changes in the microbial composition due to a switch in diet may be seen as

early as within a day (David *et al.* 2014; Sonnenburg *et al.* 2016). An animal-based diet leads to an increase in the abundance of bile-tolerant microbes (Alistipes, Bilophila, Bacteroides), while Firmicutes bacteria primarily metabolize plant polysaccharides and are thus abundant in plant-based diets (David *et al.* 2014). Other factors that have been strongly associated with gut microbiota composition are age, antibiotics, drug usage, and climatic conditions. Chevalier *et al.* (2015) found that environmental temperature had a major influence on the gut microbiome and gut physiology. Cold exposure of mice markedly shifted the gut microbiota composition which caused energy homeostasis, leading to increases in gut size, insulin sensitivity, energy expenditure and absorptive gut surface (Chevalier *et al.* 2015). The development and maturation of gut microbiota in young animals are believed to be largely influenced by age. Human babies are born largely sterile and acquire their first gut microbes from their mother's vaginal microbiome. In babies born with caesarean section, however, the vaginal seeding of microbes is lost, and the gut microbiomes of these babies instead show similarities to the mother's skin microbiome (Dominguez-Bello *et al.* 2010). The mode of birth, together with early-life antibiotic exposure, is now believed to be an important factor explaining the higher levels of asthma and allergies in children born with caesarean section (Renz-Polster *et al.* 2005; Russell *et al.* 2012).

## **The importance of microbiota in evolutionary processes**

In recent years, some people studying host microbiomes have argued that the field of evolutionary biology do not properly account for how organisms evolve together with their symbionts. The traditional view, to view species as separate entities evolving due to processes such as genetic drift, assortative mate selection, or local adaptation, has been challenged by some studies. For example, Brucker and Bordenstein (2013) found that when two species of *Nasonia* wasps mated with a more distant relative, the hybrid offspring would normally die. Treating the hybrids with antibiotics, however, made them viable and thriving. By simply eliminating the gut microbiota in the hosts, the authors could essentially tear down the ecological barrier separating the different species.

The importance of studying host microbiomes in light of evolution has recently been shown in the context of mate selection and its effect on speciation and trait evolution. Because many animals communicate by producing chemical signals, odour-releasing microbes can have significant influence on their interactions. In humans, skin bacteria produce enzymes which interact with sterile apocrine sweat to produce axillary odour (Froese *et al.* 1990). Without the bacteria, no pheromone signal is released. Symbiotic fermentative bacteria in the scent glands of striped and spotted hyenas produce odours that are species-specific, and further signal both sex and reproductive state (Theis *et al.* 2013). Similarly, the gut microbiota of termites produce colony-specific chemical cues which enable nestmate recognition. Manipulating the composition of gut bacteria with




antibiotics changes the termites' recognition behaviour toward nestmates (Matsuura 2001). Dodd (1989) discovered that rearing *Drosophila* fruit flies on two different diets drastically altered their mate choice. Flies reared on starch-based media preferred to mate with other flies that had been eating starch, and similarly, flies reared on maltose-based diet preferred maltose-eating flies, resulting in strong positive assortative mating. What was causing the changes in mate selection remained unknown until Sharon *et al.* (2013) successfully repeated the experiment and found assortative mating again after only one generation. This time, however, symbiotic microbes were the primary suspect, and Sharon *et al.* (2013) demonstrated that treating the flies with antibiotics suddenly removed their assortative mate preferences.

The idea that animals and plants are not separate entities that evolve discretely; but instead that individuals are a fusion of both host and microbe cells that evolve in symbiosis, called **holobionts**, was proposed already in 1994 by Jefferson. 'The hologenome theory of evolution' states that the fitness of a multicellular organism is derived from the entire collection of genomes (mostly microbial) representing the organism in an environment where selection occurs. Because microbes can be inherited both vertically and horizontally, the theory incorporates Lamarckian as well as Darwinian concepts. Holobionts and the hologenome theory of evolution have been widely discussed in the context of coral bleaching by infection of the bacteria *Vibrio shiloi* (Reshef *et al.* 2006; Rosenberg *et al.* 2007; Zilber-Rosenberg & Rosenberg 2008). After years of repeated bleaching events of the *Oculina patagonica* corals in the Mediterranean sea, the eastern population suddenly became resistant to *V. shiloi* infection, despite their decade-long lifespan and lack of adaptive immune system (Rosenberg *et al.* 2007). The phenomenon was termed 'experience-mediated tolerance' and attributed to the corals' symbiotic microbial communities, which can evolve and adapt much faster than the corals themselves in response to changing environmental conditions. However, the hologenome theory of evolution has encountered criticism, mainly because several microbes are not viewed as having high host fidelity, but rather environmentally acquired and context-dependent (Douglas & Werren 2016). Nevertheless, the currently rapid emerging importance of host microbiota expands our current view on how symbiotic microbes shape the evolution of hosts (Shapira 2016).

## Unresolved questions in host-microbiome research

The research surrounding host microbiomes is relatively new, since it was only recently it became possible to evaluate entire communities of microbes simultaneously. In the last couple of years, a large part of the effort in microbiome research has been directed towards developing new methods and characterizing various microbial communities. The microbiome field is therefore still largely in its development phase with a large focus on evaluating methods that give reliable and repeatable results, and to describe the present variation in microbial communities within hosts. It is crucial to first

characterize and describe what constitutes a specific host microbiome, what makes it stable, and how it changes over time within and between individual hosts, in order to understand the processes and factors governing it. For microbiome research in model organisms, such as mice and humans, a lot of effort has been put into developing reliable and repeatable extraction and sequencing techniques, bioinformatic software, as well as statistical methods to analyse microbial communities (Lozupone & Knight 2005; Caporaso *et al.* 2010; Song *et al.* 2016; Callahan *et al.* 2016; Debelius *et al.* 2016; Morton *et al.* 2017). Microbiome studies using these model host organisms have recently shifted focus, from broad-scale characterisations and comparisons, to attempts at developing theoretical frameworks to understand the observed patterns.



**Figure 4.** Evaluation of two different DNA isolation techniques (conventional DNA extraction versus direct PCR) for high-throughput amplicon sequencing of animal gut microbiomes. (A) PCoA, (B) distance network, and (C) taxonomic proportion in different sample types. Reprinted from **Paper II** (Videvall *et al.* 2017b).

However, the microbiome research of other animals are, as expected, lagging behind and is still in the development phase. As a result, there are both important research questions and methodological questions that are currently unresolved in the field of non-

model microbiomes. Almost no studies are available on how to best sample and measure the gut microbiota of animals (**Figure 4**). We also know extremely little of how the gut microbiota of different species look like and how it is colonized in offspring and subsequently matures over time. The adult microbiome seems to be heavily influenced by its earlier maturation process during juvenile stages, and this developmental window is therefore highly important to evaluate. One of the burning microbiome questions currently on several people's mind is how does the gut microbiome affect host fitness? Does the gut microbiota play a role in the growth and survival of hosts? And does the microbial community shift during disease? These are some of the questions I will address in this thesis. For an overview of host-microbiome related questions, please see 'Aim of thesis'.

## Host-parasite interactions

### Host coevolution with parasites

Though recent work on microbiomes has highlighted some of the beneficial effects microbes can have on hosts, several microbes ultimately pose a threat to host health and fitness. Being parasitic offers so many advantages, it is the most common way of life (Price 1977), and has evolved independently numerous times. The never-ending arms race between hosts and parasites has profound effects on their respective evolutionary path. Parasites continuously invent new invasion and avoidance strategies to circumvent the barriers raised by hosts. They usually have an evolutionary advantage relative their host because they tend to have short generation times, produce numerous offspring, and contain small genomes with high mutation rates. Particularly pathogenic parasites therefore place strong selective pressures on host genomes to evolve counter-adaptive measurements and defences to suppress the infection.

The extent to which parasitic organisms can influence the evolution of hosts was also highlighted by Dawkins (1999), who used parasites as one of his metaphors in 'The Extended Phenotype'. Dawkins expanded the gene-centric view, and reasons that the behaviour and physiology of hosts can be greatly modified by parasite genes, which ultimately determines the extended host phenotype. This means that the resulting host phenotype will depend, not only on the expression of genes in the host genome, but also on the expression of parasite genes, and the parasite's interactions with the host. A classic example illustrating this idea is trematode parasites which infect snails. Infected snails have thicker shells compared to uninfected snails, but are castrated by the parasite (Cheng 1973), presumably because the parasite has nothing to gain from the reproductive success of the snail, but are instead under strong selection pressure to keep


the snail alive. Hence, the parasite genes end up with a large influence on both host reproduction and host survival.

Although trematodes are not microbes, the same concept applies. There are many examples where parasitic microbes are known to significantly alter the physiology or behaviour of their host to their advantage. *Toxoplasma gondii* makes rats attracted to cat urine in order for the parasite to transfer to a feline host, the rabies virus induces e.g. hyperactivity, confusion, and hydrophobia in its host, and the malaria parasite makes vertebrates more attractive to blood-feeding mosquitoes. Although the concept of the extended phenotype is now commonly used to emphasize the importance of viewing both host and parasite genomes in light of their interactions, most studies still focus solely on the molecular aspect of one of the parties (Lambrechts *et al.* 2006).

## Host defence

Vertebrate coevolution with parasites over hundreds of millions of years has created an intricate and complex defence system, known as the immune system. It consists of the innate immunity, acting as the first layer of defence, and the adaptive immunity, which adjusts to the presence of specific pathogens. The adaptive immune system can be further classified into humoral immunity and cell-mediated immunity, which primarily defend against extracellular and intracellular parasites, respectively, and utilize different cells and molecules. The memory lymphocytes protects the host against reinfection by pathogens with the same antigens, but not against similar pathogens that have different antigenic variants. Each host carries a specific memory profile based on its history of infections, which is why the immunological memory of older individuals are usually broader relative to that of younger ones. (Frank 2002; Abbas *et al.* 2014). Together, the strength and efficiency of different aspects of the host immune system can have major implications for the resulting disease progression and severity (**Figure 5**).

Host genomes continuously evolve **resistance** and **tolerance** mechanisms to minimize the negative effects of pathogenic microbes. For example, the resistance genes of the major histocompatibility complex (MHC) constitute the most polymorphic genes in the human genome, due to their role in recognizing and binding parasite antigens. Tolerance mechanisms allow hosts to cope with pathogens by expressing host genes involved in e.g. DNA repair. Where resistance is usually measured as the inverse of infection intensity (number of parasites per unit host tissue), tolerance can be defined as the rate of change in fitness as parasite burden increases, or more informally as ‘the ability to limit the damage of a given parasite burden’ (Simms & Triplett 1994; Råberg *et al.* 2009). In order to quantify tolerance, one needs to measure it across individuals of a given host species, since variation among hosts can be caused by factors other than tolerance, such as host condition or environmental factors.



**Figure 5.** Different components of the immune system have implications for the course of infection. As an example are different mouse strains infected with the rodent malaria parasite *Plasmodium chabaudi*. Reprinted from Stevenson & Riley (2004) with permission from Springer Nature.

One of the best known examples of how hosts have evolved mechanisms to resist parasites is the human sickle cell trait, which confers resistance to malaria parasites (*Plasmodium* species). Malaria is a dangerous disease which constitutes one of the leading causes of death worldwide in humans from an infectious agent. The haemoglobin allele that gives rise to the sickle red blood cell, HbS, has evolved independently several times in different areas, and is maintained at approximately 10% frequency in malaria-endemic regions (Flint *et al.* 1998). An individual homozygous for HbS suffers from sickle cell disease, but heterozygotes have a ten-fold reduction in the risk of acquiring severe malaria (Hill *et al.* 1991; Ackerman *et al.* 2005). Remarkably (as noted already by Haldane 1949), the selection pressure from *Plasmodium* parasites on the human genome has been so strong that, besides sickle cell disease, several other red blood cell disorders that increase resistance or tolerance to malaria have independently evolved in humans. For example, alpha and beta thalassemia, ovalocytosis, glucose-6-phosphate dehydrogenase deficiency, and the Duffy-negative blood group (Flint *et al.* 1998; Kwiatkowski 2005).

## Parasite virulence

Pathogens may cause disease in hosts because they possess structural, biochemical, or genetic traits that make them **virulent**. In an ecological context, virulence is usually defined as the reduction in fitness a host experiences by a particular parasite. Before the 1980's, conventional wisdom regarded virulence as a fixed artefact based on recent associations between a parasite and its host (Levin 1996), and assumed that selection always led parasites on a directional path to becoming commensals. Evolutionary biology was able to show, however, that selection processes could favour either an increase or a decrease in virulence. Work by Anderson and May (1979; 1981) improved our understanding of the natural selection pressures that act on parasite dynamics. A higher reproductive rate may allow a parasite to increase its likelihood of transmission to new hosts, thereby increasing its fitness, but this can result in higher virulence. If virulence increases too much, the parasite may risk killing the host before successful transmission has taken place. As such, the parasite faces important economic trade-offs between transmission and virulence.

Studying the evolution of virulence and its consequences on hosts is very important from both a medical and veterinary viewpoint, because a better understanding of parasite fitness optima may allow us to reduce the negative effects parasites have on hosts. Yet, very little is known about the distribution and effects of virulence in natural populations. Theory suggests that parasite virulence should increase with coinfection of multiple strains due to competition within hosts (Nowak & May 1994; Ebert 1998). It has also been hypothesized that the existence of multiple parasite strains of the same species are the outcome of host genotype-by-parasite genotype interactions. MHC has figured as one of the main genetic host components driving this view that some alleles function better toward certain parasite lineages (Westerdahl *et al.* 2005), but fare worse against other lineages due to binding properties. Few studies have been able to demonstrate specific genotype-by-genotype interactions, but some can clearly point to the fact that disease severity results from an interaction of both host and parasite genotype together (Lambrechts *et al.* 2005).

## Immune evasion by parasites

A major part in the host-parasite relationship consists of parasites evolving variation in antigenic loci to escape host immunity. Having variable antigenic molecules can extend the time a parasite can persist in a particular host individual, and helps avoid the immunological memory of hosts. The large evolutionary pressure to generate antigenic variation in parasite genomes often cause antigenic loci to show signs of positive selection (Endo *et al.* 1996; Yang & Bielawski 2000). To cope with this strong selection pressure, some parasites have evolved genome-wide, or even localized, hypermutation mechanisms that allow them to generate more nucleotide variants via increased mutation

rates (Moxon *et al.* 1994; Caporale 1999; Ripley 1999). For example, mutator lines of the pathogenic bacterium *Pseudomonas aeruginosa*, has displayed non-random distribution of mutations along its chromosomes (Dettman *et al.* 2016), and pathogenic strains of *Escherichia coli* and *Salmonella enterica* have increased overall mutation rates (LeClerc *et al.* 1996; de Visser *et al.* 1999).

Other parasites store several genetic variants for an antigenic locus within their genome, effectively creating a vast library of antigenic variation to choose from. Parasites with this strategy express only one genetic variant at a time, called monoallelic expression, and is able to switch gene expression between the different variants. This is a strategy employed by some malaria parasites (Frank 2002; Ferreira *et al.* 2004; Recker *et al.* 2011; Bachmann *et al.* 2011; Guizetti & Scherf 2013). *Plasmodium falciparum* has a huge archive in its genome of up to 60 variants of the *var* genes coding for the surface-exposed PfEMP-1 protein. Switching gene expression between the different *var* genes can lead to some interesting population dynamics within a single host. When the host's immune system starts recognizing a particular type of antigen in a parasite, massive amounts of immune cells are proliferated that target this specific antigen. To escape this immune storm, the malaria parasite then switches expression from one *var* locus to another. If the parasite switches expression between variants too quickly, the host will develop immunity to all types early on and the infection will not be sustained. However, if the parasite is too slow switching between expression variants, it risks being cleared by the immune system before the switch has been made. This means that the timing of regulating expression between antigenic variants is crucial and finely tuned to the host's immune response. Which malaria parasites that utilize this immune evasion strategy, the rate of antigenic switching, and how the coordinated regulation of the switch works, are unanswered questions that future studies need to investigate.

More recent research has revealed the intricate immune evasion manoeuvres that parasites use to directly and effectively manipulate host gene expression to their advantage. This strategy have been detected in several organisms, but seems to be especially common in apicomplexan parasites. *Toxoplasma gondii* is well-known for inhibiting host gene expression of MHC (Leroux *et al.* 2015), *Plasmodium* parasites prevent host cell death expression pathways in liver cells (Hakimi & Cannella 2011), and *Theileria* parasites have a unique ability to completely transform host leukocytes and hijack their cellular machinery (Plattner & Soldati-Favre 2008). The challenge for the studies investigating these behaviours lies within proving that the altered host expression is indeed a direct manipulation by the parasite, and not a defence response induced by the host cell itself. It is clear, however, that parasites have evolved a wide range of sophisticated strategies to evade the immune defences of hosts; yet the molecular mechanisms underlying these activities are still largely unknown.

## Controlled host-parasite experiments

Natural systems provide great opportunities to study the ongoing evolution and distribution of hosts and parasites. The drawback of using completely wild systems, however, is the inherent difficulties to disentangle the effects due to the large environmental variation and controlling previous host exposure to microbes. Vertebrates in nature are commonly infested with a plethora of various parasites; including (but not limited to) viruses, bacteria, fungi, intestinal nematodes, blood parasites, ticks, and mites (see e.g. Biard *et al.* 2015). The microbes inside natural hosts compete with and influence each other in various ways, and as such complicate any effort to evaluate their interactions with hosts. Taking control of the confounding environmental variables in the laboratory lets us investigate the dynamics and consequences of coevolution with fewer factors. This approach may allow for the distinction between cause and effect, and can reveal detailed molecular interactions between hosts and their microbes.


Though inbred laboratory mouse strains have been used as host models in studies of infectious disease for decades, they tend to give a highly specific picture of host-parasite interactions. The immune system of lab mice do not develop properly (Beura *et al.* 2016; Abolins *et al.* 2017), and because they have been heavily artificially selected, they are not very accurate at representing natural host responses in other vertebrate taxa (White *et al.* 2010). Investigating the molecular mechanisms of host-parasite interactions using natural hosts in controlled environments offers several potential advantages. The way natural selection shapes the evolution of host responses can be studied without interference by artificial selection, vaccinations, medicines, vector control, and other anthropogenic preventive measures. It is also possible to perform controlled infection experiments in wild hosts in order to follow host responses over the course of infections. Several non-model host organisms have now been successfully studied *in vivo* using endoparasites within controlled laboratory environments. These include for example *Daphnia magna* with microsporidia parasites (Ebert 1994; Ebert *et al.* 2000), the bumblebee *Bombus terrestris* with trypanosome gut parasites (Baer and Schmid-Hempel 1999; Barribeau *et al.* 2014), the freshwater snail *Potamopyrgus antipodarum* with trematode parasites (Koskella & Lively 2007, 2009), and passerine birds with malaria parasites (Atkinson *et al.* 2000; Zehindjiev *et al.* 2008; Palinauskas *et al.* 2008, 2011; Cellier-Holzem *et al.* 2010; Ellis *et al.* 2015; Dimitrov *et al.* 2015).

## Unresolved questions in host-parasite research

Understanding the mechanisms behind host-parasite interactions poses great challenges as the outcome depends both on the host, the pathogen, and the interaction between the two. Why do some host individuals get sick while others not? What are the mechanisms



causing these inter-individual differences in host resistance and tolerance? To start investigating this, it is highly important that we examine molecular responses to infection across different taxa (**Figure 6**). To date, there has been little integration between different host systems and between wild and model organisms. Despite extensive research conducted on a subset of parasitic microbes causing human disease, we know very little about the effects of parasitic infection on different hosts. Furthermore, we need to start investigating host molecular responses to parasites over the course of infection. Most studies investigating disease systems in wildlife are dependent on first catching the host in the wild in order to sample them. This approach can be useful to measure the prevalence of low, chronic infections in hosts that survived the acute infection. However, it will not enable an improved understanding of the full infection event, including initial disease stage with parasite proliferation, peak **parasitemia**, decreasing parasitemia, and the recovery period, as well as potential relapses of the infection.



**Figure 6.** Similarities and differences in host processes related to stress response and regulation of cell death among genes differentially expressed in the avian transcriptome during peak parasitemia stage (left column) and decreasing parasitemia stage (right column). Black bars indicate significantly overrepresented processes and a higher  $-\log_{10}$  q-value means higher statistical significance. Adapted from **Paper V** (Videvall *et al.* 2015) under a CC BY 4.0 license.

Finally, we have extremely little knowledge about the specific genes implicated in disease and infection, both from a host perspective and a parasite point of view. A


handful of candidate immune genes have been intensely studied in hosts, but there are thousands of genes with potentially critical roles in the host response to infection. Likewise, the complex network of interacting genes in the parasite genome has large implications for successful transmission and infection in hosts. If we want to better understand the molecular warfare between different parasites and their hosts, it will be important to look at their interactions from a **genome**-wide point of view and over the course of infection. These are some of the aspects I intend to take into account in this thesis. For an overview of research questions related to host-parasite interactions, please see ‘Aim of thesis’.

## Genomics

### High-throughput sequencing of microbiomes

In 1977, Carl Woese and George Fox revolutionized the microbiology field by showing that the bacterial 16S ribosomal gene contains both highly conserved and hypervariable regions, and could be sequenced with universal primers to use in phylogenies and taxonomic identifications. Sanger sequencing, with its single-sequence approach, proved nonetheless highly impractical and time-consuming when it came to characterizing samples containing thousands of microbial species. Before the advent of high-throughput sequencing, most studies investigating microbial compositions had to first culture the organisms in the samples before sequencing them. Characterizing microbial communities with culture-based methods greatly underestimates the diversity, sometimes as much as 80% (Eckburg *et al.* 2005), because standard culturing medium is unsuitable for most microbes, of which many are anaerobes. This means that studies based on cultures become drastically biased towards aerobic microbes that are culturable and severely underestimate the true taxonomic diversity in a sample. However, even studies avoiding cultures drastically underestimated the microbial diversity previously. The reason was because PCR reactions with universal primers usually only result in single sequences derived from one or a few species that are common in the sample and/or bind well to the probes. As a result, when species composition was diverse, researchers had previously a lot of difficulties in recovering the full extent of the community.

With the new high-throughput sequencing techniques, the microbiome field has expanded dramatically (**Figure 7**). For the first time, we are now able to view the species making up entire bacterial communities, and we can accurately estimate their abundance and diversity. Excitingly, most of the species initially found have been completely novel (Venter *et al.* 2004), even inside the human body (Eckburg *et al.* 2005). The modern DNA sequencing techniques allow us to take a more holistic view of all the microbes in



**Figure 7.** Number of published papers containing the term "microbiome" indexed by PubMed each year.

a particular environment, and has led to a shift in the field of microbiology. Where previous studies mostly targeted single human disease-causing bacteria, such as *Escherichia coli* and *Salmonella* spp., today researchers have the possibility to evaluate the whole community of microbes in various parts of the body, and how they interact and change over time. This has led to the appreciation that the vast majority of microbes residing within our bodies are not pathogenic, but instead mutualistic or commensal, yet still have great implications for human and animal health.

## High-throughput sequencing of host responses

Understanding the role of a specific gene, molecule, or pathway in the immune system of vertebrates is not an easy task. Specialized immune cells have been found to express at least 67% of all the genes in the genome, and less than 1% of genes in the genome are only expressed in one cell type (Hyatt *et al.* 2006). Given the size and complexity of the vertebrate immune defence, we need to consider all its components in the framework as a whole, in order to discern it. Moreover, any given immune molecule can have widely contrasting outcomes depending on the epistatic effects it has with other interacting gene products (Heng *et al.* 2008). The study of single classical immune-related genes is highly valuable, and contributes much to our understanding; however, such a narrow focus risks missing other genes with highly important functions, and does not advance our knowledge of the complex interacting network of responses that constitute the immune defence. Therefore, it is crucial that we decipher expression patterns of immune cells in the context of a network, and not limit our view to one type of cell, or molecule. By using high-throughput sequencing techniques, we can start investigating the roles of the various components of the immunity, their interactions

amongst each other, and how they vary in response to different diseases. Genomics provides us with the ability to view the immune system as a whole, and characterize it as the dynamic model it constitutes.

## High-throughput sequencing of parasites

Before modern sequencing techniques, most research on parasite genetics targeted single candidate genes. It was not possible to study genome-wide expression of parasite genes. Later, microarray applications were developed and became useful at measuring gene expression, although they were inherently biased and required sequence information from a genome. With genomic tools, however, pathogens of humans quickly came into focus because of their importance in clinical studies, and their smaller, gene-dense genomes enabled easier assembly. The very first organism to have its genome sequenced, excluding viral phages, was indeed a pathogen, the bacterium *Haemophilus influenzae* (Fleischmann *et al.* 1995). After this pivotal moment, more pathogen and parasite species have had their genomes sequenced and characterized. Still lacking, however, are the many diverse parasite taxa infecting non-model organisms. Sequencing the genome of a new parasite does not only allow for the description of that particular species, but greatly enhances our understanding of parasite evolution through phylogenomic and comparative genomic studies. As a result, obtaining genomic sequences of non-model parasites have multiple benefits in the advancement of science.

In addition, with the development of RNA-sequencing methods, we now have the possibility to sequence the entire repertoire of expressed transcripts from a parasite. Some of the first studies using RNA-seq achieved this in eukaryotic pathogens such as *Schistosoma mansoni* (Almeida *et al.* 2012), *Giardia intestinalis* (Franzén *et al.* 2013), *Candida albicans* (Bruno *et al.* 2010), *Trypanosoma brucei* (Kolev *et al.* 2010), and *Plasmodium falciparum* (Otto *et al.* 2010). These **transcriptome** studies have been highly valuable as tools for improving parasite genome annotation; including information on splice sites, transcription start sites, UTR locations, novel ORFs, transcripts, and non-coding RNA. It is only recently, however, that studies have begun to evaluate parasite expression in several samples, over time, and compare different strains or genotypes (see e.g. Wurtzel *et al.* 2012). This approach allows for analyses of differential gene expression, coregulatory genes, alternative splicing events, and expression variation during infections. Furthermore, the new RNA-sequencing methods open up fantastic possibilities of evaluating simultaneous genome-wide gene expression of both host and parasite together, so-called dual RNA-seq. In conclusion, high-throughput sequencing techniques allow us for the first time to paint complete pictures of the complex molecular interactions taking place between hosts and microbes.

# Aim of thesis

With this PhD thesis, I aim to provide a deeper understanding to the following broad research questions:

1. How do we reliably and accurately measure the gut microbiome of non-model animals? (**Paper I and II**)
2. How does the gut microbiome colonize juvenile hosts and develop over time? (**Paper III and IV**)
3. How does the gut microbiota affect host fitness? (**Paper III and IV**)
4. What constitutes a healthy and a diseased microbiome, and which microbes are specifically associated with health and disease? (**Paper IV**)
5. How does the transcriptome of a host respond to a parasitic infection over time and in relation to parasites with different virulence? (**Paper V and VI**)
6. How does the transcriptome of a parasite respond to different host individuals and which genes are being utilized at different stages of the infection? (**Paper VII**)
7. How do genomes of parasites evolve and what are their evolutionary relationships? (**Paper VIII, IX, and X**)

# General methodology

## Ostrich gut microbiomes

To study the gut microbiomes of hosts in relation to health and fitness (thesis aims 1–4), I have utilized a system of ostriches (*Struthio camelus*). The ostrich system consists of a research farm located in Oudtshoorn, Western Cape, South Africa (**Figure 8**). Ostriches are the world's largest bird species, and they are a valuable economic resource being farmed for feathers, eggs, meat, and leather, yet have only been kept in captivity for a short period of time relative to other agricultural animals (Cloete *et al.* 2012). Their chicks (**Figure 9**) are highly precocial, allowing them to be raised independently from parents. Studying ostriches at a research farm brings many benefits, including highly repeatable sampling, identification of individuals and their biological parents, simultaneous hatching, exclusion of predation, and reduced influence from environmental factors such as variation in diet. Both the genetic and growth variation among the ostrich chicks in the population is high, and I have had the privilege of working with a very large sample size of 234 individuals.


In the microbiome project, we collected faecal samples and weight measurements from all ostrich chicks every second week during the first three months of their lives. We also dissected all chicks that died naturally of disease ( $n = 68$ ), as well as healthy controls



**Figure 8.** Study site in Oudtshoorn, South Africa with a group of ostrich chicks to the left and an adult male ostrich in his enclosure to the right. Photo: E.V.

(n = 60), in order to collect samples from three different sections of the gastrointestinal tract and the cloaca. All gut samples were collected in between rigorous cleaning routines, which included rinsing scalpels with hot water, soap, and 70% ethanol, followed by sterilization with a bunsen burner. Other samples we collected as part of this project were environmental samples (soil, food, water) and faecal samples from adults.


To analyse gut microbiota we have been using 16S rRNA gene sequencing. DNA from the samples was extracted using two different methods, standard DNA isolation according to the Earth Microbiome Project and Direct PCR where you circumvent the extraction steps. The results from these two DNA extraction methods were evaluated in **Paper II**. After 16S-amplicon library preparations, a total of 1152 samples were sequenced to produce 300 bp-long reads over three full sequencing runs on a Illumina MiSeq machine at the Department of Biology, Lund University. I analysed the sequence reads with bioinformatics and performed the statistical analyses in R. Further details of the study system, sample collection, laboratory procedures, and software used can be found within the methods section of each paper (**Paper I, II, III, and IV**).



**Figure 9.** Ostrich chick, two weeks old. Photo: E.V.

## Siskin malaria

To study host responses to parasites and parasite responses to hosts (thesis aims 5–7), I have been using a system of avian malaria. The eukaryotic protist parasites that cause the disease malaria are transmitted via dipteran vectors to a diverse range of vertebrate hosts, including primates, bats, rodents, ungulates, reptiles, and birds. Most of the research on host responses to malaria have been performed using primates and mice, with the effects of malaria on other vertebrate hosts remaining largely unknown. The definition of the term ‘malaria parasite’ varies greatly, but it is most commonly ascribed to all the species within the genus of *Plasmodium*. Species of *Plasmodium* should contain haemozoin pigment and use mosquitoes as their vector, however, exceptions to both of these rules exist. The term ‘malaria parasite’ is therefore sometimes also used for any haemosporidian parasites, including the genera *Haemoproteus* and *Leucocytozoon*. Malaria in birds is a highly suitable system for studying host-parasite interactions because the parasite replicates asexually in red blood cells of the host, which means we can perform repeated blood sampling of an individual to follow both the host response and the parasite over the course of an infection. The same parasite genotype



**Figure 10.** (A) Parasitemia levels over time in siskins infected with a low-virulent malaria parasite (GRW4) and a high-virulent parasite (SGS1). Reprinted from **Paper VI**. (B) Photograph of a microscope view of avian malaria parasites (in pink) on a blood smear. Photo: E.V. (C) Drawing of a malaria parasite merozoite highlighting major organelles. Reprinted from Cowman & Crabb (2006) with permission from Elsevier.



can reach different infection intensities (**parasitemia**) in different individuals (**Figure 10**), creating a great opportunity to investigate gene expression differences in hosts.

In collaboration with colleagues at the Nature Research Centre, in Vilnius, Lithuania who have the facilities to perform controlled infection experiments in birds, we have studied the molecular effects of avian malaria infection. Wild-caught juvenile Eurasian siskins (*Carduelis spinus*; **Figure 11**) were caught with mist nests and housed in aviaries at the Biological Station Rybachy of the Zoological Institute of the Russian Academy of Sciences on the Curonian Spit in the Baltic Sea. Siskins have previously proven to be excellent study organisms for avian malaria experiments (Palinauskas *et al.* 2008, 2011). They are susceptible to several *Plasmodium* lineages, yet juvenile siskins caught early in the summer are uninfected because the vectors have not yet emerged. Siskins are also abundant at the study site and suitable to house in captivity as opposed to several other small wild birds, which may not provide enough blood for high-throughput sequencing.



**Figure 11.** Eurasian siskin resting. Photo: Eva Mårtensson.

Three naturally infected siskins with high parasitemia intensities of *Haemoproteus tartakovskyi* lineage SISKIN1 were used to sequence the genome of this parasite (**Paper VIII**). Siskins used in the infection experiments were inoculated with blood from a single infected donor bird, and control birds were injected with blood from an uninfected


bird. The birds were then observed and blood samples for parasitemia calculations and for RNA-sequencing were collected at specific time points during the infection. We have used three different lineages of *Plasmodium* in three different infection experiments: *P. ashfordi* GRW2 (**Paper V** and **VII**), *P. relictum* GRW4 (**Paper VI**), and *P. relictum* SGS1 (**Paper VI**).

Collected blood samples were frozen in liquid nitrogen and subsequently extracted to retrieve RNA for transcriptome sequencing and DNA for genome sequencing. After library preparations, samples in the *H. tartakovskyi* genome project were sequenced on a 454 Life Sciences Genome Sequencer FLX+ machine at the Department of Biology, Lund University. The blood samples from the *Plasmodium* infection experiments were sequenced using paired-end Illumina HiSeq 2000 RNA-sequencing to generate transcriptome-wide gene expression data. The sequences were subsequently analysed with bioinformatic methods and software. For further details regarding the specific methods used in the avian malaria project, please see the individual papers (**Paper V**, **VI**, **VII**, **VIII**, **IX**, and **X**).

# Results and discussion

## Measuring the gut microbiome in birds


In order to investigate a new trait in vertebrates, it is crucial to first develop a methodological procedure that has been verified to give accurate and repeatable results. As an example, hormone levels in birds are often found to be non-repeatable, even



**Figure 12.** Microbiota differences between two sampling techniques (cloacal swabs and faeces) and three parts of the gastrointestinal tract (ileum, caecum, colon). (a) Network, (b) NMDS, (c) microbial diversity, (d) taxonomic composition. Reprinted from **Paper I** (Videvall *et al.* 2017c) with permission from John Wiley and Sons.

within the same study, as they can yield very high variation within individuals (Ouyang *et al.* 2011). Researchers must therefore be aware of what exactly they are measuring and how repeatable they can expect the results to be. Failure to do so may result in spurious relationships and false positives. Before we started large-scale investigations of ostrich gut microbiomes, we therefore set out to test the accuracy of two commonly used sampling techniques for bird microbiomes (**Paper I**), as well as the repeatability of two DNA isolation techniques (**Paper II**).

In gut microbiome research of birds and reptiles, a large amount of studies have sampled the animals by swabbing the cloacae with cotton swabs. By using ostrich juveniles, we tested in **Paper I** the ability of faecal and cloacal sampling at recovering the gut microbiota. We found that cloacal swabs yield a microbial community that is different from that of the ileum, caecum, and colon (**Figure 12**). Faecal samples were also poor representatives of the microbiota of the ileum and the caecum. However, faecal samples were significantly better than cloacal swabs at measuring the microbiota of the colon. We can therefore be confident that sampling faeces yields a microbial community that is not identical, but largely similar to that of the colon, and we recommend gut microbiome researchers to sample faeces, whenever possible, when studying birds.



**Figure 13.** Repeatability of extraction replicates (purple) and PCR replicates (green) when using a direct PCR method (light colour) and a conventional DNA isolation method (dark colour). Adapted from **Paper II** (Videvall *et al.* 2017b) under a CC BY 4.0 license.


When preparing microbiome samples for 16 rRNA gene sequencing, the standard approach includes the isolation of DNA during a series of 32 cleaning and purification steps. In human microbiome research, a recent paper evaluated the possibility of using a potentially faster method with only 4 steps called ‘Direct PCR’, to circumvent the demanding laboratory procedure of DNA extraction (Flores *et al.* 2012). This method seemed promising, however, the study was performed on human samples only, and with

a small sample size and outdated sequencing technology. It was therefore unclear whether this potentially time-saving preparation technique would yield accurate results using different sample types from a bird species.

In **Paper II**, we set out to evaluate these two methods using five gut-related sample types from ostrich juveniles: gut content from the ileum, caecum, and colon, faecal samples, and cloacal swabs. In terms of costs, we found that the direct PCR method was faster and cheaper compared to the standard DNA extraction procedure. The direct PCR method produced highly repeatable and comparable microbial communities to the DNA extraction method in the caecal, colon, and faecal samples. However, the repeatability and accuracy of both library preparation techniques were much lower in the cloacal and ileal samples (**Figure 13**). In conclusion, both the direct PCR method and the DNA extraction method performed well and had high repeatability when using samples with high DNA concentration, but were equally poor at measuring the microbial community of low biomass samples. We can therefore recommend that researchers use whichever one of these methods they prefer.


## Development of gut microbiota in juvenile ostriches

To evaluate how the gut microbiome develops and matures over time in juveniles of a non-model species, we studied in **Paper III** the faecal microbiota of ostrich chicks during their first three months of life. We found that age has a very strong effect on the microbial community (**Figure 14**). For example, the microbial diversity and richness progressively increased with age and the dissimilarities between the microbiota in individuals initially increased and later decreased with age. One week old individuals were highly dissimilar to all other ages, with a community largely dominated by *Akkermansia muciniphila*, likely a consequence of the internal yolk sac they still were



**Figure 14.** NMDS plot (left) and microbial richness (right) of ostrich gut microbiomes, with colours indicating age of individuals. Adapted from **Paper III** (Videvall *et al.* 2018).

absorbing during this early stage. There were striking differences in colonization and extinction patterns of different bacterial taxa. Some classes increased in relative abundances with age, for example Bacilli, Clostridia, and Planctomycetia, while others rapidly decreased with age, such as Verrucomicrobiae, Erysipelotrichi, and Gamma-proteobacteria (**Figure 15**).




**Figure 15.** Relative abundances of microbial classes in the ostrich gut during development, with age in weeks on the x-axes and A = Adults. Reprinted from **Paper III** (Videvall *et al.* 2018).

The growth rates of the ostrich chicks were extremely variable, with some individuals weighing five times as much as their contemporaries at 12 weeks of age. We evaluated the effect of the microbiome on juvenile growth and found that individuals one week of age showed a strong positive correlation of microbial diversity with growth, while an overall negative association was found when evaluating all ages. Detailed analyses of specific taxa associated with growth showed that the family Bacteroidaceae was positively correlated with juvenile weight at week 1 of age, while Enterobacteriaceae, Enterococcaceae, and Lactobacillaceae were negatively correlated with weight at either week 2 or week 6.

## Strong link between gut microbiota and mortality in ostriches


A large number of ostrich chicks died of suspected disease ( $n = 68$ ) during the first three months of age (**Figure 16**). Despite attentive daily care and provisioning, high mortality rates of juveniles is common in ostrich rearing facilities, with some years up to 80% of the population dying within the first three months (Cloete *et al.* 2001). This high mortality is believed to be associated with some sort of gut-related disease, as many individuals that die first display characteristic disease behaviour such as lethargy and poor appetite, together with symptoms of diarrhea and an inflamed gastrointestinal tract. Several candidate bacterial pathogens have been appointed as potential culprits, but they differ between studies, and many of them are also present in a normal vertebrate gut microbiota as commensals or pathobionts.

In **Paper IV**, we found that weight was a good predictor of the ostrich chicks' probability of survival, but primarily during the first weeks, where low weight was associated with a low probability of surviving to the next week. Most of the chicks that died lost weight rapidly shortly before dying (**Figure 16**), likely an effect of disease symptoms and not eating. The gut microbiota showed major dissimilarities between diseased and control individuals, and this pattern was so strong, it explained a much larger part of the variation than that of the age of individuals, despite age being the most prominent explanatory variable of the faecal microbiota in healthy individuals (**Paper III**). These large differences between diseased and control individuals were highly significant in all three regions of the gastrointestinal tract. Interestingly, however, the beta diversity between individuals differed depending on gut region, with the diseased individuals being more similar to each other in the upper gastrointestinal tract (ileum) than the control individuals were to each other, but with a reversed pattern present in the lower part of the gut (caecum and colon).



**Figure 16.** Log-transformed weight of control individuals (blue) and individuals that died from suspected disease (red) during the first three months of age. Grey lines represent all other individuals. Adapted from **Paper IV**.

The microbial diversity of diseased individuals was much lower than that of control individuals, in all three gut regions. Surprisingly, the ileum showed no sign of increasing in alpha diversity with age after controlling for disease, but the effects of age increased progressively along the gastrointestinal tract. We found several taxa associated with disease, i.e. enriched in the ileum, caecum, and colon of diseased individuals (for example Enterobacteriaceae, Peptostreptococcaceae, Porphyromonadaceae, *Clostridium*, *Paeniclostridium*), and several health-associated taxa depleted in diseased individuals (e.g. S24-7, Lachnospiraceae including *Roseburia*, *Coprococcus*, and *Blautia*, Ruminococcaceae, Erysipelotrichaceae, and *Turicibacter*) (**Figure 17**).




**Figure 17.** Taxonomic composition of Clostridia in the ileum (top) and of Bacteroidia in the caecum and colon (bottom). Left column = control individuals and right column = diseased individuals. Adapted from **Paper IV**.



We could also show that the food, water, and soil did not contribute significantly to the gut microbiota of diseased individuals, suggesting a small likelihood of bacterial contamination from these environmental sources. Taken together, all our results in **Paper IV** points to a pattern of extreme gut dysbiosis in the ostrich chicks that died from suspected disease.

## Avian transcriptome responses to malaria parasites

In **Paper V** and **VI** we evaluated transcriptome responses of siskins to malaria parasites over the course of infection. A large number of genes was found differentially expressed in birds during infection with the high-virulent lineages *P. ashfordi* GRW2 (**Paper V**) and *P. relictum* SGS1 (**Paper VI**), but not during infection with the low-virulent malaria lineage *P. relictum* GRW4 (**Paper VI**). The differentially expressed genes in the birds infected with GRW2 were largely similar during the peak and decreasing parasitemia stages (**Figure 18**). However, a high number of genes involved in the immune system was upregulated during peak parasitemia compared to decreasing parasitemia. The decreasing parasitemia stage had instead genes overrepresented within functions related to 'mature B cell differentiation'. The most highly expressed genes in the blood were those related to haemoglobin and betaglobin (**Paper V**).




**Figure 18.** Transcriptome expression in birds with malaria infection during peak parasitemia (left) and decreasing parasitemia (right) compared to control birds. Each point illustrates the mean normalized log-transformed expression levels for one gene ( $n = 18,618$  genes) and significant differentially expressed genes are coloured in red. Adapted from **Paper V** (Videvall *et al.* 2015) under a CC BY 4.0 license.

In the transcriptome responses to the high-virulent parasites GRW2 and SGS1, we found that up- and down-regulated genes in infected birds were overrepresented within processes related to oxidative stress, negative regulation of cell death, metabolic and

catabolic processes, and regulation of gene expression. In fact, the regulatory miRNA genes belonged to the most significantly upregulated genes during both malaria infection experiments, and some of them were also highly correlated with parasitemia intensity. In **Paper VI** we further evaluated the expression of protein-coding genes that have essential roles in producing mature miRNA molecules and in the miRNA gene silencing pathway, and could demonstrate that these miRNA-related genes were also significantly upregulated during infection.

In **Paper VI**, we had the possibility of sequencing a much larger number of individuals and more time points, resulting in a total of 76 bird transcriptomes. Interestingly, the host molecular response to the high-virulent parasite SGS1 resulted in a clear circular trajectory of individuals over time in the principal component analysis (**Figure 19**), with the size of the trajectory being highly correlated with the number of parasites in the blood. That the magnitude of the host response was highly associated with the quantity




**Figure 19.** (A) PCA of all 76 host transcriptomes. (B) Correlation between host individual movement on the PC1 dimension and parasitemia levels. (C) The high-virulent parasite SGS1 causes a circular trajectory of host transcriptomes in the PCA, whereas (D) the low-virulent parasite GRW4 does not. Reprinted from **Paper VI**.

of parasites during infection was also demonstrated for ~800 genes which showed significant positive or negative associations with parasitemia intensity.

## Malaria parasite transcriptome responses to hosts


Because of the largely unbiased nature of high-throughput sequencing, as compared to e.g. targeted microarrays, we retrieve sequence data from all the material in our sample, regardless of origin. This means that dual RNA-sequencing can be utilized to collect reads in samples that contain transcripts from more than one organism, for example in the case of a host and a parasite. At the time I started my PhD, the field of dual RNA-seq was in its infancy and the only relevant study that had used this method was one evaluating the fungal model pathogen *Candida albicans* in mice (Tierney *et al.* 2012).



**Figure 20.** Density graphs of transcript GC content in (A) the initial unfiltered transcriptome assembly. After rigorous bioinformatic filtering steps, I constructed (B) the filtered *Plasmodium ashfordi* assembly, (C) transcripts matching bird sequences, and (D) unknown contigs containing both bird and parasite transcripts which were subsequently utilized. Reprinted from **Paper VII** (Videvall *et al.* 2017a) with permission from John Wiley and Sons.

In several aspects, my journey into this field of dual RNA-seq data was novel. Not only was I working with a relatively understudied wildlife disease system in terms of molecular studies, the dual RNA-seq data was derived from an infection experiment involving a non-model host species without any genome sequence *and* a non-model parasite species without any genome sequence. The aim of building a clean parasite transcriptome assembly from scratch without any genomic references was therefore monumental.

In **Paper VII**, I show that this endeavour was possible through multiple bioinformatic filtering steps, because we present the first transcriptome assembly of a wildlife bird malaria parasite, *Plasmodium ashfordi* lineage GRW2 (**Figure 20**). Expression of *P. ashfordi* transcripts was analysed in three different host individuals and during two time points of the infection, peak parasitemia and decreasing parasitemia. We could demonstrate that *P. ashfordi* exhibits host-specific gene expression as there were several differentially expressed parasite transcripts between individual hosts, but no differences between time points. We further identified single nucleotide polymorphisms in the *P. ashfordi* transcriptome, that could potentially be a result of real variants in the parasite population, or rare gene duplication events where the transcribed transcripts remain near-identical to each other. When evaluating sequence identity with other apicomplexans, *P. ashfordi* was most similar to the human parasite *P. falciparum*, and we directly compared annotated gene functions of these two parasites (**Figure 21**). Finally, we identified a long list of *P. ashfordi* transcripts derived from genes documented to be involved in the *Plasmodium* invasion of red blood cells.




**Figure 21.** Differences between the transcriptomes of the avian malaria parasite *Plasmodium ashfordi* (light blue) and the human malaria parasite *Plasmodium falciparum* (dark blue) in terms of transcripts annotated with specific gene ontology terms. *P. ashfordi* has fewer transcripts annotated as processes involved in host interactions because these sequences are likely evolving faster compared to other genes, and thus have become too differentiated in avian malaria parasites for sequence similarity searches. Reprinted from **Paper VII** (Videvall *et al.* 2017a) with permission from John Wiley and Sons.

## Evolution of malaria parasite genomes

Because the recently assembled *P. ashfordi* transcriptome represented the only available genomic resource of a non-mammalian malaria parasite (**Paper VII**), it became incredibly valuable in our phylogenomic analyses of **haemosporidians**. Previous studies have found sequence similarities between human malaria parasites and avian malaria parasites. It has therefore been repeatedly suggested that mammalian *Plasmodium* are not monophyletic, and that instead host switches of parasites from birds to primates are responsible for these similarities (Waters *et al.* 1991; Pick *et al.* 2011).

We assembled the genome of *Haemoproteus tartakovskiyi*, a bird blood parasite in the sister genus to *Plasmodium*, to use as an outgroup in a new phylogeny based on genomic data. In **Paper VIII**, we described the genome characteristics of *H. tartakovskiyi*, and use it together with *P. ashfordi* to create a well-supported phylogeny of *Plasmodium* parasites. The *H. tartakovskiyi* genome is small and AT-rich (25.4%), similar to many *Plasmodium* parasites. It also contains expansion clusters of gene families that are slightly higher in GC%, a possible sign of species-specific invasion-related genes similar to *rif*, *var*, and *stevor* in *P. falciparum*.


Phylogenetic analyses were conducted in two steps. First, an initial phylum-wide phylogeny of 17 apicomplexan species was constructed, which confirmed the placement of *H. tartakovskiyi* as an outgroup to *Plasmodium*. Second, we constructed two phylogenetic trees with only Haemosporidian sequence data, which resulted in a supported monophyletic clade of mammalian-infecting *Plasmodium* parasites, with the only available non-mammalian *Plasmodium* (*P. ashfordi*) as a sister taxon (**Figure 22**).



**Figure 22.** Two *Plasmodium* phylogenetic trees based on genomic sequence data with *H. tartakovskiyi* as outgroup, supporting a monophyletic clade of mammalian malaria parasites (green) with avian haemosporidians (blue) as sister taxa. Reprinted from **Paper VIII** (Bensch *et al.* 2016) under a CC BY-NC 4.0 license.

In **Paper IX**, we utilized both of these newly assembled genomic resources of *P. ashfordi* and *H. tartakovskyi* to search for three key genes involved in the thiamine (vitamin B<sub>1</sub>) biosynthesis pathway. Interestingly, the genes coding for this essential vitamin are present in the genomes of primate malaria parasites, but absent in the genomes of rodent malaria parasites and other closely related apicomplexans. It has been suggested that these genes in primate malaria parasites are a result of horizontal gene transfer from bacteria (Frech & Chen 2011). However, our analyses show that these genes are not only present in the genomes of all three avian *Plasmodium* parasites evaluated, but are also actively expressed as transcripts. Furthermore, we located these genes in the genome of a species in the sister genus, *Haemoproteus*, which suggests that the thiamine genes have been present in the common ancestor but subsequently lost in the rodent malaria species (Hellgren *et al.* 2017).

Finally, in **Paper X**, I again utilized our assembled *H. tartakovskyi* genome and *P. ashfordi* transcriptome, together with the genomes of two newly sequenced avian malaria parasites, *P. relictum* and *P. gallinaceum* (Böhme *et al.* 2016). I conducted comparative genomic analyses of a large number of eukaryotes with AT-rich genomes to identify the species with the most extreme AT-bias. The human malaria parasite *P. falciparum*, has repeatedly been described as the most extreme eukaryotic organism when it comes to nucleotide composition. However, my results show that the sequenced avian malaria parasites (all three of them) have a much lower GC content, by a large margin (**Figure 23**).



**Figure 23.** Density of transcript GC content in three avian malaria parasites (green–blue) compared to five AT-rich mammalian parasites (columns). Adapted from **Paper X** (Videvall 2018) under a CC BY-NC 4.0 license.

# References

- Abbas AK, Lichtman AH, Pillai S (2014) *Basic immunology: functions and disorders of the immune system*. Elsevier Health Sciences.
- Abolins S, King EC, Lazarou L *et al.* (2017) The comparative immunology of wild and laboratory mice, *Mus musculus domesticus*. *Nature Communications*, **8**, 14811.
- Ackerman H, Usen S, Jallow M *et al.* (2005) A Comparison of Case-Control and Family-Based Association Methods: The Example of Sickle-Cell and Malaria. *Annals of Human Genetics*, **69**, 559–565.
- Almeida GT, Amaral MS, Beckedorff FCF *et al.* (2012) Exploring the *Schistosoma mansoni* adult male transcriptome using RNA-seq. *Experimental Parasitology*, **132**, 22–31.
- Anderson RM, May RM (1979) Population biology of infectious diseases: Part I. *Nature*, **280**, 361–367.
- Anderson RM, May RM (1981) The Population Dynamics of Microparasites and Their Invertebrate Hosts. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **291**, 451–524.
- Atkinson CT, Dusek RJ, Woods KL, Iko WM (2000) Pathogenicity of Avian Malaria in Experimentally- Infected Hawaii Amakihi. *Journal of Wildlife Diseases*, **36**, 197–204.
- Atkinson C, LaPointe D (2009) Introduced avian diseases, climate change, and the future of Hawaiian honeycreepers. *Journal of Avian Medicine and Surgery*, **23**, 53–63.
- Bachmann A, Predehl S, May J *et al.* (2011) Highly co-ordinated var gene expression and switching in clinical *Plasmodium falciparum* isolates from non-immune malaria patients. *Cellular Microbiology*, **13**, 1397–1409.
- Baer B, Schmid-Hempel P (1999) Experimental variation in polyandry affects parasite loads and fitness in a bumble-bee. *Nature*, **397**, 151–154.
- Barribeau SM, Sadd BM, du Plessis L, Schmid-Hempel P (2014) Gene expression differences underlying genotype-by-genotype specificity in a host–parasite system. *Proceedings of the National Academy of Sciences*, **111**, 3496–3501.
- Bensch S, Canbäck B, DeBarry JD *et al.* (2016) The Genome of *Haemoproteus tartakovskyi* and Its Relationship to Human Malaria Parasites. *Genome Biology and Evolution*, **8**, 1361–1373.
- Beura LK, Hamilton SE, Bi K *et al.* (2016) Normalizing the environment recapitulates adult human immune traits in laboratory mice. *Nature*, **532**, 512–516.
- Biard C, Monceau K, Motreuil S, Moreau J (2015) Interpreting immunological indices: The importance of taking parasite community into account. An example in blackbirds *Turdus merula*. *Methods in Ecology and Evolution*, **6**, 960–972.
- Böhme U, Otto TD, Cotton J *et al.* (2016) Complete avian malaria parasite genomes reveal host-specific parasite evolution in birds and mammals. *bioRxiv*, 1–32. doi: 10.1101/086504.

- Bolnick DI, Snowberg LK, Caporaso JG *et al.* (2014) Major Histocompatibility Complex class IIb polymorphism influences gut microbiota composition and diversity. *Molecular Ecology*, **23**, 4831–4845.
- Brooks AW, Kohl KD, Brucker RM, van Opstal EJ, Bordenstein SR (2016) Phylosymbiosis: Relationships and Functional Effects of Microbial Communities across Host Evolutionary History. *PLoS Biology*, **14**, e2000225.
- Brucker RM, Bordenstein SR (2013) The Hologenomic Basis of Speciation: Gut Bacteria Cause Hybrid Lethality in the Genus *Nasonia*. *Science*, **341**, 667–669.
- Bruno VM, Wang Z, Marjani SL *et al.* (2010) Comprehensive annotation of the transcriptome of the human fungal pathogen *Candida albicans* using RNA-seq. *Genome Research*, **20**, 1451–1458.
- Callahan BJ, McMurdie PJ, Rosen MJ *et al.* (2016) DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, **13**, 581–583.
- Caporale LH (1999) Chance favors the prepared genome. In: *Annals of the New York Academy of Sciences*, pp. 1–21.
- Caporaso JG, Kuczynski J, Stombaugh J *et al.* (2010) QIIME allows analysis of high-throughput community sequencing data. *Nature Methods*, **7**, 335–336.
- Cellier-Holzem E, Esparza-Salas R, Garnier S, Sorci G (2010) Effect of repeated exposure to *Plasmodium relictum* (lineage SGS1) on infection dynamics in domestic canaries. *International Journal for Parasitology*, **40**, 1447–1453.
- Cheng TC (1973) *General Parasitology*. Academic Press, New York.
- Chevalier C, Stojanović O, Colin DJ *et al.* (2015) Gut Microbiota Orchestrates Energy Homeostasis during Cold. *Cell*, **163**, 1360–1374.
- Cloete SWP, Brand TS, Hoffman L *et al.* (2012) The development of ratite production through continued research. *World's Poultry Science Journal*, **68**, 323–334.
- Cloete SWP, Lambrechts H, Punt K, Brand Z (2001) Factors related to high levels of ostrich chick mortality from hatching to 90 days of age in an intensive rearing system. *Journal of the South African Veterinary Association*, **72**, 197–202.
- Cowman AF, Crabb BS (2006) Invasion of Red Blood Cells by Malaria Parasites. *Cell*, **124**, 755–766.
- Cryan JF, Dinan TG (2012) Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour. *Nature Reviews Neuroscience*, **13**, 701–712.
- Davenport ER (2016) Elucidating the role of the host genome in shaping microbiome composition. *Gut Microbes*, **7**, 178–184.
- David LA, Maurice CF, Carmody RN *et al.* (2014) Diet rapidly and reproducibly alters the human gut microbiome. *Nature*, **505**, 559–563.
- Dawkins R (1999) *The extended phenotype: the long reach of the gene*. Oxford University Press.
- Debelius J, Song SJ, Vazquez-Baeza Y *et al.* (2016) Tiny microbes, enormous impacts: what matters in gut microbiome studies? *Genome Biology*, **17**, 217.
- Dettman JR, Szepeanacz JL, Kassen R (2016) The properties of spontaneous mutations in the opportunistic pathogen *Pseudomonas aeruginosa*. *BMC Genomics*, **17**, 27.
- Dimitrov D, Palinauskas V, Iezhova TA *et al.* (2015) *Plasmodium* spp.: An experimental study on vertebrate host susceptibility to avian malaria. *Experimental Parasitology*, **148**, 1–16.



- Dodd DMB (1989) Reproductive Isolation as a Consequence of Adaptive Divergence in *Drosophila pseudoobscura*. *Evolution*, **43**, 1308–1311.
- Dominguez-Bello MG, Costello EK, Contreras M *et al.* (2010) Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proceedings of the National Academy of Sciences*, **107**, 11971–11975.
- Douglas AE, Werren JH (2016) Holes in the Hologenome: Why Host-Microbe Symbioses Are Not Holobionts. *mBio*, **7**, e02099-15.
- Ebert D (1994) Virulence and Local Adaptation of a Horizontally Transmitted Parasite. *Science*, **265**, 1084–1086.
- Ebert D (1998) Experimental Evolution of Parasites. *Science*, **282**, 1432–1436.
- Ebert D, Lipsitch M, Mangin KL (2000) The Effect of Parasites on Host Population Density and Extinction: Experimental Epidemiology with *Daphnia* and Six Microparasites. *The American Naturalist*, **156**, 459–477.
- Eckburg PB, Bik EM, Bernstein CN *et al.* (2005) Diversity of the Human Intestinal Microbial Flora. *Science*, 1635–1638.
- Ellis VA, Cornet S, Merrill L *et al.* (2015) Host immune responses to experimental infection of *Plasmodium relictum* (lineage SGS1) in domestic canaries (*Serinus canaria*). *Parasitology Research*, **114**, 3627–3636.
- Endo T, Ikeo K, Gojobori T (1996) Large-scale search for genes on which positive selection may operate. *Molecular Biology and Evolution*, **13**, 685–690.
- Ferreira MU, Nunes S, Wunderlich G (2004) Antigenic diversity and immune evasion by malaria parasites. *Clinical and Diagnostic Laboratory Immunology*, **11**, 987–995.
- Fleischmann R, Adams M, White O *et al.* (1995) Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science*, **269**, 496–512.
- Flint J, Harding RM, Boyce AJ, Clegg JB (1998) The population genetics of the haemoglobinopathies. *Baillière's Clinical Haematology*, **11**, 1–51.
- Flores GE, Henley JB, Fierer N (2012) A Direct PCR Approach to Accelerate Analyses of Human-Associated Microbial Communities. *PLoS ONE*, **7**, e44563.
- Ford SA, King KC (2016) Harnessing the Power of Defensive Microbes: Evolutionary Implications in Nature and Disease Control. *PLoS Pathogens*, **12**, e1005465.
- Frank SA (2002) *Immunology and Evolution of Infectious Disease*. Princeton University Press.
- Frank DN, Robertson CE, Hamm CM *et al.* (2011) Disease phenotype and genotype are associated with shifts in intestinal-associated microbiota in inflammatory bowel diseases. *Inflammatory Bowel Diseases*, **17**, 179–184.
- Franzén O, Jernström-Hultqvist J, Einarsson E *et al.* (2013) Transcriptome Profiling of *Giardia intestinalis* Using Strand-specific RNA-Seq. *PLoS Computational Biology*, **9**, e1003000.
- Frech C, Chen N (2011) Genome Comparison of Human and Non-Human Malaria Parasites Reveals Species Subset-Specific Genes Potentially Linked to Human Disease. *PLoS Computational Biology*, **7**, e1002320.
- Froebe C, Simone A, Charig A, Eigen E (1990) Axillary Malodor Production - a New Mechanism. *Journal of the Society of Cosmetic Chemists*, **41**, 173–185.
- Gensollen T, Iyer SS, Kasper DL, Blumberg RS (2016) How colonization by microbiota in early life shapes the immune system. *Science*, **352**, 539–544.

- Gerardo NM, Parker BJ (2014) Mechanisms of symbiont-conferred protection against natural enemies: an ecological and evolutionary framework. *Current Opinion in Insect Science*, **4**, 8–14.
- Gilbert JA, Neufeld JD (2014) Life in a World without Microbes. *PLoS Biology*, **12**, e1002020.
- Goodrich JK, Davenport ER, Waters JL, Clark AG, Ley RE (2016) Cross-species comparisons of host genetic associations with the microbiome. *Science*, **352**, 532–535.
- Goodrich JK, Waters JL, Poole AC *et al.* (2014) Human Genetics Shape the Gut Microbiome. *Cell*, **159**, 789–799.
- Gordon HA, Pesti L (1971) The gnotobiotic animal as a tool in the study of host microbial relationships. *Bacterial Reviews*, **35**, 390–429.
- Gough E, Shaikh H, Manges AR (2011) Systematic Review of Intestinal Microbiota Transplantation (Fecal Bacteriotherapy) for Recurrent *Clostridium difficile* Infection. *Clinical Infectious Diseases*, **53**, 994–1002.
- Groussin M, Mazel F, Sanders JG *et al.* (2017) Unraveling the processes shaping mammalian gut microbiomes over evolutionary time. *Nature Communications*, **8**, 14319.
- Guizetti J, Scherf A (2013) Silence, activate, poise and switch! Mechanisms of antigenic variation in *Plasmodium falciparum*. *Cellular Microbiology*, **15**, 718–726.
- Hakimi M, Cannella D (2011) Apicomplexan parasites and subversion of the host cell microRNA pathway. *Trends in Parasitology*, **27**, 481–486.
- Haldane JBS (1949) The rate of mutation of human genes. *Hereditas*, **35**, 267–273.
- Hamilton WD, Axelrod R, Tanese R (1990) Sexual reproduction as an adaptation to resist parasites (a review). *Proceedings of the National Academy of Sciences*, **87**, 3566–3573.
- van der Heijden PJ, Stok W, Bianchi AT (1987) Contribution of immunoglobulin-secreting cells in the murine small intestine to the total “background” immunoglobulin production. *Immunology*, **62**, 551–555.
- Hellgren O, Bensch S, Videvall E (2017) *De novo* synthesis of thiamine (vitamin B<sub>1</sub>) is the ancestral state in *Plasmodium* parasites – evidence from avian haemosporidians. *Parasitology*, 1–6. doi: 10.1017/S0031182017002219.
- Heng TSP, Painter MW, Elpek K *et al.* (2008) The Immunological Genome Project: networks of gene expression in immune cells. *Nature Immunology*, **9**, 1091–1094.
- Hill MJ (1997) Intestinal flora and endogenous vitamin synthesis. In: *European journal of cancer prevention: the official journal of the European Cancer Prevention Organisation (ECP)*, pp. S43–S45.
- Hill AVS, Allsopp CEM, Kwiatkowski D *et al.* (1991) Common West African HLA antigens are associated with protection from severe malaria. *Nature*, **352**, 595–600.
- Hirakawa H (2002) Coprophagy in leporids and other mammalian herbivores. *Mammal Review*, **32**, 150–152.
- Hird SM, Sánchez C, Carstens BC, Brumfield RT (2015) Comparative Gut Microbiota of 59 Neotropical Bird Species. *Frontiers in Microbiology*, **6**, 1403.
- Hope ME, Hold GL, Kain R, El-Omar EM (2005) Sporadic colorectal cancer – role of the commensal microbiota. *FEMS Microbiology Letters*, **244**, 1–7.
- Hyatt G, Melamed R, Park R *et al.* (2006) Gene expression microarrays: glimpses of the immunological genome. *Nature Immunology*, **7**, 686–691.

- Ichinohe T, Pang IK, Kumamoto Y *et al.* (2011) Microbiota regulates immune defense against respiratory tract influenza A virus infection. *Proceedings of the National Academy of Sciences*, **108**, 5354–5359.
- Inagaki H, Suzuki T, Nomoto K, Yoshikai Y (1996) Increased susceptibility to primary infection with *Listeria monocytogenes* in germfree mice may be due to lack of accumulation of L-selectin+ CD44+ T cells in sites of inflammation. *Infection and Immunity*, **64**, 3280–3287.
- Jefferson R (1994) The Hologenome in “A Decade of PCR: Celebrating 10 Years of Amplification, Proceedings of a Symposium” Video released by Cold Spring Harbor Laboratory Press, 1994.
- Kamada N, Chen GY, Inohara N, Núñez G (2013) Control of pathogens and pathobionts by the gut microbiota. *Nature Immunology*, **14**, 685–690.
- Kohl KD, Dearing MD, Bordenstein SR (2017) Microbial communities exhibit host species distinguishability and phyllosymbiosis along the length of the gastrointestinal tract. *Molecular Ecology*. doi: 10.1111/mec.14460.
- Kolev NG, Franklin JB, Carmi S *et al.* (2010) The Transcriptome of the Human Pathogen *Trypanosoma brucei* at Single-Nucleotide Resolution. *PLoS Pathogens*, **6**, e1001090.
- Koskella B, Lively CM (2007) Advice of the rose: Experimental coevolution of a trematode parasite and its snail host. *Evolution*, **61**, 152–159.
- Koskella B, Lively CM (2009) Evidence for negative frequency-dependent selection during experimental coevolution of a freshwater snail and a sterilizing trematode. *Evolution*, **63**, 2213–2221.
- Kubinak JL, Stephens WZ, Soto R *et al.* (2015) MHC variation sculpts individualized microbial communities that control susceptibility to enteric infection. *Nature Communications*, **6**, 8642.
- Kwiatkowski DP (2005) How Malaria Has Affected the Human Genome and What Human Genetics Can Teach Us about Malaria. *The American Journal of Human Genetics*, **77**, 171–192.
- Lambrechts L, Fellous S, Koella JC (2006) Coevolutionary interactions between host and parasite genotypes. *Trends in Parasitology*, **22**, 12–16.
- Lambrechts L, Halbert J, Durand P, Gouagna LC, Koella JC (2005) Host genotype by parasite genotype interactions underlying the resistance of anopheline mosquitoes to *Plasmodium falciparum*. *Malaria Journal*, **4**, 3.
- LeClerc JE, Li B, Payne WL, Cebula TA (1996) High Mutation Frequencies Among *Escherichia coli* and *Salmonella* Pathogens. *Science*, **274**, 1208–1211.
- Leroux LP, Dasanayake D, Rommereim LM *et al.* (2015) Secreted *Toxoplasma gondii* molecules interfere with expression of MHC-II in interferon gamma-activated macrophages. *International Journal for Parasitology*, **45**, 319–322.
- Levin BR (1996) The Evolution and Maintenance of Virulence in Microparasites. *Emerging Infectious Diseases*, **2**, 93–102.
- Lozupone C, Knight R (2005) UniFrac: a New Phylogenetic Method for Comparing Microbial Communities. *Applied and Environmental Microbiology*, **71**, 8228–8235.
- MacDonald WL (1991) Biological Control of Chestnut Blight: Use and Limitations of Transmissible Hypovirulence. *Plant Disease*, **75**, 653.

- Macpherson AJ, Harris NL (2004) Interactions between commensal intestinal bacteria and the immune system. *Nature Reviews Immunology*, **4**, 478–485.
- Maes M, Kubera M, Leunis J-C, Berk M (2012) Increased IgA and IgM responses against gut commensals in chronic depression: Further evidence for increased bacterial translocation or leaky gut. *Journal of Affective Disorders*, **141**, 55–62.
- Matsuura K (2001) Nestmate recognition mediated by intestinal bacteria in a termite, *Reticulitermes speratus*. *Oikos*, **92**, 20–26.
- McManus KF, Taravella AM, Henn BM *et al.* (2017) Population genetic analysis of the DARC locus (Duffy) reveals adaptation from standing variation associated with malaria resistance in humans. *PLoS Genetics*, **13**, 48–65.
- Mideo N (2009) Parasite adaptations to within-host competition. *Trends in Parasitology*, **25**, 261–268.
- Milgroom MG, Cortesi P (2004) Biological Control of Chestnut Blight with Hypovirulence: A Critical Analysis. *Annual Review of Phytopathology*, **42**, 311–338.
- Moreira LA, Iturbe-Ormaetxe I, Jeffery JA *et al.* (2009) A *Wolbachia* Symbiont in *Aedes aegypti* Limits Infection with Dengue, Chikungunya, and *Plasmodium*. *Cell*, **139**, 1268–1278.
- Moret Y, Schmid-Hempel P (2000) Survival for Immunity: The Price of Immune System Activation for Bumblebee Workers. *Science*, **290**, 1166–1168.
- Morton JT, Sanders J, Quinn RA *et al.* (2017) Balance Trees Reveal Microbial Niche Differentiation. *mSystems*, **2**, e00162-16.
- Moxon ER, Rainey PB, Nowak MA, Lenski RE (1994) Adaptive evolution of highly mutable loci in pathogenic bacteria. *Current Biology*, **4**, 24–33.
- Ngwa CJ, Pradel G (2015) Coming soon: probiotics-based malaria vaccines. *Trends in Parasitology*, **31**, 2–4.
- Nicholson JK, Holmes E, Kinross J *et al.* (2012) Host-Gut Microbiota Metabolic Interactions. *Science*, **336**, 1262–1267.
- Nowak MA, May RM (1994) Superinfection and the Evolution of Parasite Virulence. *Proceedings of the Royal Society B: Biological Sciences*, **255**, 81–89.
- Otto TD, Wilinski D, Assefa S *et al.* (2010) New insights into the blood-stage transcriptome of *Plasmodium falciparum* using RNA-Seq. *Molecular Microbiology*, **76**, 12–24.
- Ouyang JQ, Hau M, Bonier F (2011) Within seasons and among years: When are corticosterone levels repeatable? *Hormones and Behavior*, **60**, 559–564.
- Palinauskas V, Valkiūnas G, Bolshakov CV, Bensch S (2008) *Plasmodium relictum* (lineage P-SGS1): Effects on experimentally infected passerine birds. *Experimental Parasitology*, **120**, 372–380.
- Palinauskas V, Valkiūnas G, Bolshakov CV, Bensch S (2011) *Plasmodium relictum* (lineage SGS1) and *Plasmodium ashfordi* (lineage GRW2): The effects of the co-infection on experimentally infected passerine birds. *Experimental Parasitology*, **127**, 527–533.
- Pasteur L (1885) Observations relatives à la note précédente de M. Duclaux. *CR Acad Sci (Paris)*, **100**, 68.
- Petnicki-Ocwieja T, Hrcir T, Liu Y-J *et al.* (2009) Nod2 is required for the regulation of commensal microbiota in the intestine. *Proceedings of the National Academy of Sciences*, **106**, 15813–15818.

- Pick C, Ebersberger I, Spielmann T, Bruchhaus I, Burmester T (2011) Phylogenomic analyses of malaria parasites and evolution of their exported proteins. *BMC Evolutionary Biology*, **11**, 167.
- Plattner F, Soldati-Favre D (2008) Hijacking of Host Cellular Functions by the Apicomplexa. *Annual Review of Microbiology*, **62**, 471–487.
- Price PW (1977) General Concepts on the Evolutionary Biology of Parasites. *Evolution*, **31**, 405.
- Qian J, Hospodsky D, Yamamoto N, Nazaroff WW, Peccia J (2012) Size-resolved emission rates of airborne bacteria and fungi in an occupied classroom. *Indoor Air*, **22**, 339–351.
- Qin J, Li R, Raes J *et al.* (2010) A human gut microbial gene catalogue established by metagenomic sequencing. *Nature*, **464**, 59–65.
- Råberg L, Graham AL, Read AF (2009) Decomposing health: tolerance and resistance to parasites in animals. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **364**, 37–49.
- Ramanan D, Bowcutt R, Lee SC *et al.* (2016) Helminth infection promotes colonization resistance via type 2 immunity. *Science*, **352**, 608–612.
- Recker M, Buckee CO, Serazin A *et al.* (2011) Antigenic Variation in *Plasmodium falciparum* Malaria Involves a Highly Structured Switching Pattern. *PLoS Pathogens*, **7**, e1001306.
- Renz-Polster H, David MR, Buist AS *et al.* (2005) Caesarean section delivery and the risk of allergic disorders in childhood. *Clinical and Experimental Allergy*, **35**, 1466–1472.
- Reshef L, Koren O, Loya Y, Zilber-Rosenberg I, Rosenberg E (2006) The Coral Probiotic Hypothesis. *Environmental Microbiology*, **8**, 2068–2073.
- van Riper C, van Riper SG, Goff ML, Laird M (1986) The Epizootiology and Ecological Significance of Malaria in Hawaiian Land Birds. *Ecological Monographs*, **56**, 327–344.
- Ripley LS (1999) Predictability of mutant sequences. Relationships between mutational mechanisms and mutant specificity. In: *Annals of the New York Academy of Sciences*, pp. 159–172.
- Rosenberg E, Koren O, Reshef L, Efrony R, Zilber-Rosenberg I (2007) The role of microorganisms in coral health, disease and evolution. *Nature Reviews Microbiology*, **5**, 355–362.
- Round JL, Mazmanian SK (2009) The gut microbiota shapes intestinal immune responses during health and disease. *Nature Reviews Immunology*, **9**, 313–323.
- Russell SL, Gold MJ, Hartmann M *et al.* (2012) Early life antibiotic-driven changes in microbiota enhance susceptibility to allergic asthma. *EMBO Reports*, **13**, 440–447.
- Sadzikowski MR, Sperry JF, Wilkins TD (1977) Cholesterol-reducing bacterium from human feces. *Applied and Environmental Microbiology*, **34**, 355–362.
- Sender R, Fuchs S, Milo R (2016) Are We Really Vastly Outnumbered? Revisiting the Ratio of Bacterial to Host Cells in Humans. *Cell*, **164**, 337–340.
- Shapira M (2016) Gut Microbiotas and Host Evolution: Scaling Up Symbiosis. *Trends in Ecology & Evolution*, **31**, 539–549.
- Sharon G, Segal D, Ringo JM *et al.* (2010) Commensal bacteria play a role in mating preference of *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences*, **107**, 20051–20056.
- Simms EL, Triplett J (1994) Costs and Benefits of Plant Responses to Disease: Resistance and Tolerance. *Evolution*, **48**, 1973.

- Song SJ, Amir A, Metcalf JL *et al.* (2016) Preservation Methods Differ in Fecal Microbiome Stability, Affecting Suitability for Field Studies. *mSystems*, **1**, e00021-16.
- Sonnenburg ED, Smits SA, Tikhonov M *et al.* (2016) Diet-induced extinctions in the gut microbiota compound over generations. *Nature*, **529**, 212–215.
- Spor A, Koren O, Ley R (2011) Unravelling the effects of the environment and host genotype on the gut microbiome. *Nature Reviews Microbiology*, **9**, 279–290.
- Sprinz H, Kundel DW, Dammin GJ *et al.* (1961) The Response of the Germ-free Guinea Pig to Oral Bacterial Challenge with *Escherichia coli* and *Shigella flexneri*: With Special Reference to Lymphatic Tissue and the Intestinal Tract. *The American Journal of Pathology*, **39**, 681–695.
- Stevenson MM, Riley EM (2004) Innate immunity to malaria. *Nature Reviews Immunology*, **4**, 169–180.
- Strachan DP (1989) Hay fever, hygiene, and household size. *BMJ*, **299**, 1259–1260.
- Theis KR, Venkataraman A, Dycus JA *et al.* (2013) Symbiotic bacteria appear to mediate hyena social odors. *Proceedings of the National Academy of Sciences*, **110**, 19832–19837.
- Tierney L, Linde J, Müller S *et al.* (2012) An interspecies regulatory network inferred from simultaneous RNA-seq of *Candida albicans* invading innate immune cells. *Frontiers in Microbiology*, **3**, 1–14.
- Toivanen P, Vaahtovuori J, Eerola E (2001) Influence of Major Histocompatibility Complex on Bacterial Composition of Fecal Flora. *Infection and Immunity*, **69**, 2372–2377.
- Tollenaere C, Pernechele B, Mäkinen HS *et al.* (2014) A hyperparasite affects the population dynamics of a wild plant pathogen. *Molecular Ecology*, **23**, 5877–5887.
- Turnbaugh PJ, Backhed F, Fulton L, Gordon JI (2008) Diet-Induced Obesity Is Linked to Marked but Reversible Alterations in the Mouse Distal Gut Microbiome. *Cell Host & Microbe*, **3**, 213–223.
- Turnbaugh PJ, Hamady M, Yatsunenko T *et al.* (2009) A core gut microbiome in obese and lean twins. *Nature*, **457**, 480–484.
- van Valen L (1973) A new evolutionary law. *Evolutionary Theory*, **1**, 1–30.
- Venter JC, Remington K, Heidelberg JF *et al.* (2004) Environmental Genome Shotgun Sequencing of the Sargasso Sea. *Science*, **304**, 66–74.
- Videvall E (2018) *Plasmodium* parasites of birds have the most AT-rich genes of eukaryotes. *Microbial Genomics*, 1–9. doi: 10.1099/mgen.0.000150.
- Videvall E, Cornwallis CK, Ahrén D *et al.* (2017a) The transcriptome of the avian malaria parasite *Plasmodium ashfordi* displays host-specific gene expression. *Molecular Ecology*, **26**, 2939–2958.
- Videvall E, Cornwallis CK, Palinauskas V, Valkiūnas G, Hellgren O (2015) The Avian Transcriptome Response to Malaria Infection. *Molecular Biology and Evolution*, **32**, 1255–1267.
- Videvall E, Song SJ, Bensch HM *et al.* (2018) The development of gut microbiota in ostriches and its association with juvenile growth. *bioRxiv*, 270017.
- Videvall E, Strandh M, Engelbrecht A, Cloete S, Cornwallis CK (2017b) Direct PCR Offers a Fast and Reliable Alternative to Conventional DNA Isolation Methods for Gut Microbiomes. *mSystems*, **2**, e00132-17.

- Videvall E, Strandh M, Engelbrecht A, Cloete S, Cornwallis CK (2017c) Measuring the gut microbiome in birds: Comparison of faecal and cloacal sampling. *Molecular Ecology Resources*, 1–11. doi: 10.1111/1755-0998.12744.
- Vighi G, Marcucci F, Sensi L, Di Cara G, Frati F (2008) Allergy and the gastrointestinal system. *Clinical & Experimental Immunology*, **153**, 3–6.
- Vijay-Kumar M, Aitken JD, Carvalho FA *et al.* (2010) Metabolic Syndrome and Altered Gut Microbiota in Mice Lacking Toll-Like Receptor 5. *Science*, **328**, 228–231.
- de Visser JAGM, Zeyl CW, Gerrish PJ, Blanchard JL, Lenski RE (1999) Diminishing returns from mutation supply rate in asexual populations. *Science*, **283**, 404–406.
- Wahl M, Goecke F, Labes A, Dobretsov S, Weinberger F (2012) The Second Skin: Ecological Role of Epibiotic Biofilms on Marine Organisms. *Frontiers in Microbiology*, **3**, 1–21.
- Waters AP, Higgins DG, McCutchan TF (1991) *Plasmodium falciparum* appears to have arisen as a result of lateral transfer between avian and human hosts. *Proceedings of the National Academy of Sciences*, **88**, 3140–3144.
- Westerdahl H, Waldenström J, Hansson B *et al.* (2005) Associations between malaria and MHC genes in a migratory songbird. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 1511–1518.
- White NJ, Turner GDH, Medana IM, Dondorp AM, Day NPJ (2010) The murine cerebral malaria phenomenon. *Trends in Parasitology*, **26**, 11–15.
- Woese CR, Fox GE (1977) Phylogenetic structure of the prokaryotic domain: The primary kingdoms. *Proceedings of the National Academy of Sciences*, **74**, 5088–5090.
- Wu H-J, Wu E (2012) The role of gut microbiota in immune homeostasis and autoimmunity. *Gut Microbes*, **3**, 4–14.
- Wurtzel O, Sesto N, Mellin JR *et al.* (2012) Comparative transcriptomics of pathogenic and non-pathogenic *Listeria* species. *Molecular Systems Biology*, **8**, 1–14.
- Yang Z, Bielawski JP (2000) Statistical methods for detecting molecular adaptation. *Trends in Ecology & Evolution*, **15**, 496–503.
- Yilmaz B, Portugal S, Tran TM *et al.* (2014) Gut Microbiota Elicits a Protective Immune Response against Malaria Transmission. *Cell*, **159**, 1277–1289.
- Zehtindjiev P, Ilieva M, Westerdahl H *et al.* (2008) Dynamics of parasitemia of malaria parasites in a naturally and experimentally infected migratory songbird, the great reed warbler *Acrocephalus arundinaceus*. *Experimental Parasitology*, **119**, 99–110.
- Zilber-Rosenberg I, Rosenberg E (2008) Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiology Reviews*, **32**, 723–735.

# Acknowledgements

This PhD project was possible thanks to the **Department of Biology** at Lund University. It has been by financed by generous grants from **the Swedish Research Council** (grant 2010-5641) and a **Wallenberg Academy fellowship** to C.K.C., and by **the Swedish Research Council** (grant 621-2011-3548, grant 2016-03419) and **the Crafoord Foundation** (grant 20120630) to O.H. It has further been generously funded by grants from **the Royal Physiographic Society of Lund**, **the Helge Ax:son Johnson Foundation**, **the Längmanska Cultural Foundation**, **the Lund Animal Protection Foundation**, **the Lars Hierta Memorial Foundation**, **the Jörgen Lindström's Foundation**, **the Knut and Alice Wallenberg Foundation**, **Geneco Research School**, and **Heinrich Poll's genetic scholarship** to E.V.

Figure 2 has been reprinted with permission from **Springer Nature**: Kamada N, Chen GY, Inohara N, Núñez G (2013) Control of pathogens and pathobionts by the gut microbiota. *Nature Immunology*, 14, 685–690.

Figure 5 has been reprinted with permission from **Springer Nature**: Stevenson MM, Riley EM (2004) Innate immunity to malaria. *Nature Reviews Immunology*, 4, 169–180.

Figure 10C has been reprinted with permission from **Elsevier**: Cowman AF, Crabb BS (2006) Invasion of red blood cells by malaria parasites. *Cell*, 124, 755–766.



# Thank you

I've thoroughly enjoyed my time during the PhD, and in many ways I've considered myself lucky to be able to work with fascinating science and interesting research questions on a daily basis. The Department of Biology in Lund has been a wonderful place to work and study, primarily because of the people there. In this section, I want to share some of my personal thoughts and gratitude towards the many people who have contributed to this thesis or the scientific environment.

I've been fortunate to be surrounded by multiple senior scientists that have supported me in various ways throughout the PhD. Several of these seniors I have regarded as official or unofficial mentors. Their common ground has been a genuine interest in my development and growth as a scientist. They are listed in no particular order of importance.

My main supervisor, **Charlie Cornwallis**. Thank you for your patience, your support, and the invaluable advice over the years. It's been an intense period together with field work in South Africa, discussing microbiome data, and sending manuscripts back and forth. Your involvement in the ostrich microbiome project has been tremendous, including everything from planning of the proposal, caring for the animals and sampling, statistical advice, and writing papers. You have so many great qualities as a scientist, but the one I appreciate the most is probably your dedication.

My main supervisor, **Olof Hellgren**. Your enthusiasm and passion for science is very inspiring. One of the most valuable things a PhD student can have is a supervisor who is really interested in your work, and becomes excited over new results and discoveries. You have that, and I think that is a wonderful trait in a senior scientist. Your knowledge about the field of avian malaria is endless, which has of course been very valuable to a malaria novice like myself. Thank you for all the support and encouragement during my PhD, and for making a difficult but fun challenge to my bioinformatic skills in dual RNA-seq.

My co-supervisor, **Bengt Hansson**. You were one of the very first people who believed in my potential as a scientist. In fact, you believed I could do advanced bioinformatics long before I had any idea that I could, and it quickly became one of my major passions. Thank you for giving me so many opportunities and for always having that wonderful curiosity about science in general. Even though you were never directly involved in the

PhD projects themselves, you were instrumental in moulding that foundation of knowledge of which the PhD was built upon.

My PhD examiner, **Dennis Hasselquist**. During my time as an undergraduate student in Lund, one of my favourite courses was the Evolutionary Animal Ecology course, mostly because of your enthusiasm. I remember one lecture you had about sexually transmitted diseases in birds and I found it absolutely fascinating. Your door has always been open for me to come discuss various aspects of my PhD and academia in general, and for that I am very grateful. It has been incredibly valuable to have you around during my PhD not only as a highly committed and caring examiner, but also as a scientific mentor.

My departmental scientific mentor, **Dag Åhrén**. You know you've made a good choice in a departmental official mentor when you just want to meet and have lunch with this person all the time. What's even better is when that feeling is reciprocated and your mentor is genuinely interested in your PhD progress and development as a scientist. Dag, you are all that, and so much more. Your pedagogic skills are amazing, and you have such a big compassion and understanding for the person behind the scientist, it is very inspiring.

My Geneco mentor, **Mats Hansen**. Thank you for your genuine interest in my personal growth as a scientist and your continued support not only during the official mentor period, but still to this day. I have found it highly valuable to experience a scientific view from outside academia.

My Master of Science supervisor, **Staffan Bensch**. The master's project you supervised was not only interesting, it became very important for my initial scientific journey since it resulted in my first publication, largely because of you. Ever since, you've continued to support and believe in me during my entire PhD. Not many people know that you are probably the main reason why this particular PhD project happened in the first place. You have been one of the most important and encouraging people during my PhD. And for that I am very grateful.

My unofficial mentor, **Erik Svensson**. We have travelled and lived in South Africa together, cooked ostrich egg pancakes, and explored nature reserves together. You are a great source of zoological knowledge, and I've appreciated you teaching me about various birds and insects. What I think have been particularly valuable are the many interesting discussions we've had with various perspectives of science, academia, and evolutionary biology. I think that, instead of constantly defaulting to the old traditional way, academia in general, could benefit from a more open and diverse set of thinking. In this regard, you have been a source of inspiration for me in many ways.

I'd like to extend my thanks to several other supportive seniors in Lund, whom I've been positively influenced or inspired by: **Karin Rengefors, Helena Westerdahl, Lars Råberg, Maren Wellenreuther, Björn Canbäck, Honor Prentice, Jep Agrell, Åke**

**Lindström, Thomas Alerstam, Niklas Wahlberg, Johan Hollander, Torbjörn Säll, Jörgen Ripa, Jessica Abbott, Tobias Uller, Anders Brodin, Caroline Isaksson, Susanne Åkesson, Lars Pettersson, Ola Olsson, Jan-Åke Nilsson, and Anders Hedenström.**

I would also like to thank **the Research Education Board** (Forskarutbildningsnämnden) and **the Departmental Board** (Biologiska institutionens styrelse). In particular the head of the department **Christer Löfstedt**, and the director of studies, **Karin Rengefors** and **Per Lundberg**. It's been a pleasure to participate and contribute with a PhD student's perspective over all these years. I have learned a lot about university politics, departmental policies, and running a successful academic institution. I'm confident these experiences has better prepared me for a possible future in academia. Thank you also to the committee of **Geneco Research School** for including me as a board representative for PhD students. I'd also like to thank the other members of the **PhD students' council in Biology** (Biologiska doktorandrådet) for the many meetings during these years, and for assisting me when I organized the PhD student seminar.

**Christer Löfstedt**, I have always found it admiring how the Head of the Department manages to take the time to frequently stop and chat about everything and nothing with the PhD students. I honestly think you are one of several reasons we are getting so much praise from visiting international researchers regarding the social and friendly atmosphere in the Ecology building.

**Svante Pääbo**, thank you for accepting my invitation to come give a seminar in Lund and for being so incredibly accommodating during that whole process. It's very inspiring for us young academics to see someone with your career being so humble and down to earth.

**Inger Ekström**, it has always been a great pleasure to exchange information regarding web administration, LUCRIS, and LUP with you. I wish all departments could be so lucky as to have such a dedicated and colourful information officer as you are.

**Erling Jirle**, your interest and knowledge about all kinds of organisms is admiring. Thank you for some great conversations over the years, and for assistance with LUCRIS administration. **Kristina Arnebrant**, thank you for all assistance with LUCRIS.

**Anne Fogelberg**, you've been there for me from the start, when I've been sick, when I've been healthy. You've been incredibly patient when I'm struggling with the different grants and wading through the jungle of financial rules. I will miss your wonderful sense of humour and your genuine compassion about other people.

**Carl Sjökvist** and **Cecilia Thomasson**, the two magicians who can sort out any practical matters in an instant. And always with a smile! You have significantly simplified the practical aspects of my PhD.

**Anna Drews**, my amazing friend and office mate. You have meant so much during these years and I'm so grateful for all your support during both positive and negative times. I am lucky to be able to call you one of my closest friends, and I will truly miss all our wonderful lunches, dinners, and parties together.

**Jane Jönsson**, thank you for your friendship and everything you've done for me over the years. We've travelled together, lived together (in SA), crawled through caves, prepared äppelmust, and have lots of other great memories to look back at.

**Max Lundberg**, I actually don't remember when we first became friends because it feels like we've known each other for such a long time. For some reason we just seem to get along incredibly well, maybe we just understand each other, or maybe we're more similar than we think? Whatever the reason is, I've truly appreciated both our friendship and all our scientific endeavours at different conferences. And thank you for so patiently teaching impossible me about all the various bird species.

**Maja Tarka**, my academic big sister. I've looked up to you from the day I started working as research assistant, and I really appreciate you as a person and as a scientist. Thank you for some great times as office mates together.

**Martin Andersson** (in MEEL), you were such a fantastic friend during my MSc and in the beginning of my PhD. I was so happy for your new position in Kalmar, but simultaneously sad to see you leave. I've very much appreciated all our conversations and times together.

**Asghar Muhammad**, I still remember how you so elegantly handled an aggressive question during one of my first conferences, it was very inspiring. One of your many strengths is dedication to both the science and your colleagues.

**Arne Andersson**, you are such a fantastic person. For some reason I feel like we connected exceptionally well from the very start, and I'm so happy for that. I love your honesty and generosity. I will truly miss our coffee breaks and beer nights.

**Maria Strandh**, thank you for being such a wonderful and dedicated colleague and friend. Your help with setting up the laboratory aspects of the microbiome project has been invaluable.

**Lokesh Manoharan**, sometimes I feel like you're my mental doppelgänger. We view things so similarly. I love your easy-going nature and your compassion.

**Martin Stervander**, I have very much enjoyed your way of thinking, your scientific passion, and attention to details. I have appreciated our heated scientific discussions about various topics, and our common goal to continuously improve things.

**Utku Urhan**, I really enjoy your company and your personality, and you make me laugh all the time.

**Marco Klein Heerenbrink**, you have a fantastic way of seeing things, and just like me you sometimes get caught up in details. You've been a valuable friend since we met.

**Anna Nordén**, one of the highlights while working is hearing your wonderful laugh down the corridor. Thank you for spreading so much optimism and enthusiasm around you.

**Xi Huang**, thank you for being so generous and a very inspiring person to be around.

**Nayden Chakarov**, I love your enthusiasm and positive energy. Thank you for always believing in me more than I do.

And a large thank you to **Pablo Salmón, Hannah Watson, Johan Nilsson, Julio Neto, Martin N. Andersson, John Waller, John Kirwan, Beatriz Willink Castro, Tom Evans, Katrine Lund-Hansen, Ester Arévalo Sureda, Suvi Ponnikas** for being awesome and fun. There's been a lot of dinners, parties, pub evenings.. Thanks for all the laughs and memories.

**Arne Hegemann**, for showing me how to catch jackdaws and for taking care of the lab meetings when I was unable. **Julian Melgar**, for your assistance with the samples and for the lovely homemade dinners in South Africa. **Talatu Tende**, for bringing your Nigerian warmth and compassion to the cold Sweden. **Anna Stöckl**, for all the work and contribution to the Research Education Board. **Tomas Johansson**, for all your help with the MiSeq sequencing runs. **Markus Ringnér** for your help when one of my hard drives was failing. **Anu Helin**, for being a wonderful friend and for giving me a guided lab tour in Kalmar.

**Mads Fristrup Schou, Vincenzo Ellis, Philip Downing, Maria Svensson Coelho, Johan Bentzer, Markus Lindh, Jacob Roved, Hanna Sigeman, Kajsa Warfvinge, Nathalie Feiner, Jothi Kumar, Fredrik Andreasson, Linus Hedh, Gabriel Norevik, Dafne Ram, Emily O'Connor, Mikkel Willemoes, Ann-Kathrin Ziegler, Arif Ciloglu, Anna Runemark, Hanna Laakkonen, Reinder Radersma, Sandeep Kushwaha, Pallavi Chauhan**, for being such amazing colleagues and friends.

My colleagues in South Africa, **Maud Bonato, Naomi Serfontein, and Anel Engelbrecht**, thank you for all help with the ostriches, the sampling, and the dissections. Thank you also for being so welcoming and friendly while I've been visiting. I'm also grateful to **Adriaan Olivier, Zanell Brand, Stefan Engelbrecht, Schalk Cloete, Molatelo Mokoeler, and Tonny** for your assistance and advice regarding the ostrich microbiome project.

My colleagues in Lithuania, **Vaidas Palinauskas and Gediminas Valkiūnas**, it's been a pleasure collaborating with you and visiting your lab and department. Thank you for your hospitality and your enormous contribution to the malaria transcriptome project.

My colleagues in the USA, especially **Se Jin Song, Amnon Amir, Jamie Morton, Yoshiki Vázquez-Baeza, and Rob Knight**. Also thank you **Daniel McDonald, Chris**

**Callewaert, Justine Debelius, Tomasz Kościółek, Anupriya Tripathi, Jon Sanders, Luke Thompson, Greg Humphrey, Zhenjiang Xu, Stefan Janssen, and Qiyun Zhu** for taking the time to provide valuable advice, informative lectures, lab tours, and scientific discussions. And most importantly, thank you for making me feel so welcome at UCSD with all the lunches, dinners, parties, and beer nights. I had a fantastic time in San Diego and the Knight Lab.

All the wonderful and social visiting researchers who have had a large positive influence on my time in Lund, **Amparo Herrera-Dueñas, Luz García-Longoria, Sara Pardal, Michaël Moens, Josué Martínez-de la Puente, Martina Ferraguti, Paulo Pulgarin, Karina Ivanova, Kostas Sagonas, Katie Duryea, Barbara Lukasch, Cassandra Silverio, Rosa Sánchez Guillen, Marcos Lima**, and many others. We've had some really fantastic dinners, beer nights, fika gatherings, nature excursions, festival happenings, taco evenings, parties, coffee breaks, celebrations, Friday pubs, and nights out in Lund, Malmö, and Copenhagen. And some science too of course..

**Alice Evans** and **Dovilė Murauskaitė**, it was both fun and rewarding teaching you some bioinformatics and mentoring your genome projects. **Hanna Bensch**, thank you for all the help with the ostrich microbiome project and the follow-up project. I had a wonderful time with you in South Africa, and I've appreciated being your kitchen assistant.

**Maj Persson** and **Anna Persson**, I'm very grateful for the beautiful thesis cover.

**Alexander Suh, Jan Engler, and Luc Lens**, thank you for inviting and introducing me to your departments with incredible hospitality and generosity.

Thank you to colleagues previously at Lund for great company and discussions, **Cecilia Nilsson, Kristin Scherman, Sandra Sköld Chiriac, Dagmar Clough, Sissel Sjöberg, Andreas Nord, Keith Larson, Elin Sild, Magnus Ellström, Georg Andersson, and Ingrid Sassenhagen**.

The Friday Pub (**Pub Einar**), organized by **Magnus Ellström** and **Utku Urhan** during my time. It has been great to end the week with a few beers at this very social and fun event. Thank you to everyone who keeps the pub going strong by participating, I have truly enjoyed it.

Everyone participating in the **Journal Club** and the **Malaria Group** over the years.

**The Molecular Ecologist**, and its contributors who invited me to their inspiring community and gave me a wonderful forum to write about my thoughts in molecular ecology, bioinformatics, and genomics.

**Natursidan**, which provided me a great forum to write about popular science and nature-related news in Swedish, improving my popular science-writing while doing so.

**UPPMAX**, for allowing me to use their servers to analyse sequence data.

All my wonderful friends at the **Workshop on Genomics** in Český Krumlov, Czech Republic. **Scott Handley** and **Konrad Paszkiewicz**, thank you for the opportunities to teach at the workshop, it's been fantastic and memorable experiences. **Guy Leonard**, **Sophie Shaw**, **Josie Paris**, **Daniel McDonald**, **Rayan Chikhi**, there's no other people I'd rather have flaming shots at 3am on a Tuesday morning than with you guys. You are not only great friends but also very talented bioinformaticians. **Brian Haas**, **Julian Catchen**, **Bill Cresko**, **Mike Zody**, and **Chris Wheat**, thank you for being so inspirational and for your genuine interest in my academic career.

**MEEL** might just be one of the best lab groups there is. I have really enjoyed being a part of this group for so many years. With four semesters as lab meeting organizer, several years of journal club organizer, photo event organizer, Christmas party planner, web administrator, **MEEL** corner constructor, and **MEEL** Twitter account manager, I hope I have been able to give back just a little bit to this wonderful group of people. Thank you everyone in **MEEL** (none named and none forgotten) for these amazing years and inspiring lab meetings.

The steering committee of the **Geneco Research School**, thank you for creating such a fantastic organization for us early-career molecular ecologists. I have enjoyed many stimulating summer and winter meetings together with other **Geneco members**. Everyone in my **Geneco Mentor cohort**, you guys are awesome and I wish all of you a fantastic future whatever you choose to do in life.

I'm of course grateful for all my **co-authors**, who have significantly contributed to papers both within and outside the thesis.

Finally, I want to thank my parents **Börje** and **Maj-Lis**, my siblings **Marie**, **Tomas**, **Mats**, and my parents-in-law **Gun-Britt** and **Hans** for all the support and encouragement during these years.


**Daniel**, my wonderful husband. Your love, support, and belief in me have been invaluable. Thank you.







- I **Videvall E**, Strandh M, Engelbrecht A, Cloete S, Cornwallis CK. (2017) Measuring the gut microbiome in birds: Comparison of faecal and cloacal sampling. *Molecular Ecology Resources*. doi: 10.1111/1755-0998.12744.
- II **Videvall E**, Strandh M, Engelbrecht A, Cloete S, Cornwallis CK. (2017) Direct PCR offers a fast and reliable alternative to conventional DNA isolation methods for gut microbiomes. *mSystems*, 2: e00132-17.
- III **Videvall E**, Song SJ, Bensch HM, Strandh M, Engelbrecht A, Serfontein N, Hellgren O, Olivier A, Cloete S, Knight R, Cornwallis CK. (2018) The development of gut microbiota in ostriches and its association with juvenile growth. *bioRxiv*. 270017. doi: 10.1101/270017.
- IV **Videvall E**, Song SJ, Bensch HM, Strandh M, Engelbrecht A, Serfontein N, Hellgren O, Olivier A, Cloete S, Knight R, Cornwallis CK. (2018) Early-life mortality linked to gut dysbiosis in ostriches. Manuscript.
- V **Videvall E**, Cornwallis CK, Palinauskas V, Valkiūnas G, Hellgren O. (2015) The avian transcriptome response to malaria infection. *Molecular Biology and Evolution*, 32: 1255-1267.
- VI **Videvall E**, Palinauskas V, Valkiūnas G, Hellgren O. (2018) Avian transcriptome responses to high- and low-virulent malaria strains over the course of infection. Manuscript.
- VII **Videvall E**, Cornwallis CK, Ahrén D, Palinauskas V, Valkiūnas G, Hellgren O. (2017) The transcriptome of the avian malaria parasite *Plasmodium ashfordi* displays host-specific gene expression. *Molecular Ecology*, 26: 2939-2958.
- VIII Bensch S, Canbäck B, DeBarry JD, Johansson T, Hellgren O, Kissinger JC, Palinauskas V, **Videvall E**, Valkiūnas G. (2016) The genome of *Haemoproteus tartakovskyi* and its relationship with human malaria parasites. *Genome Biology and Evolution*, 8: 1361-1373.
- IX Hellgren O, Bensch S, **Videvall E**. (2017) *De novo* synthesis of thiamine (vitamin B1) is the ancestral state in *Plasmodium* parasites – evidence from avian haemosporidians. *Parasitology*. doi: 10.1017/S0031182017002219.
- X **Videvall E**. (2018) *Plasmodium* parasites of birds have the most AT-rich genes of eukaryotes. *Microbial Genomics*, 4: 1–9.



LUND UNIVERSITY

Faculty of Science  
Department of Biology  
ISBN 978-91-7753-577-5



978 91 7753 577 5