

LUND UNIVERSITY

Typification and emendation of Seirophora Poelt to include species segregated from **Teloschistes Norman**

Frödén, Patrik; Lassen, Per

Published in: Lichenologist

DOI: 10.1017/S002428290401432X

2004

Link to publication

Citation for published version (APA): Frödén, P., & Lassen, P. (2004). Typification and emendation of Seirophora Poelt to include species segregated from Teloschistes Norman. Lichenologist, 36(5), 289-298. https://doi.org/10.1017/S002428290401432X

Total number of authors: 2

General rights

Unless other specific re-use rights are stated the following general rights apply:

- Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the
- legal requirements associated with these rights

· Users may download and print one copy of any publication from the public portal for the purpose of private study You may not further distribute the material or use it for any profit-making activity or commercial gain
You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00

Typification and emendation of *Seirophora* Poelt to include species segregated from *Teloschistes* Norman

Patrik FRÖDÉN and Per LASSEN

Abstract: The type material of *Physcia magara* Kremp., which is the type species of *Seirophora* Poelt (*Teloschistaceae*), is shown to be heterogeneous. It is a mixture of *Teloschistes villosus* (Ach.) Norman and *Ramalina maciformis* (Delile) Bory. *Physcia magara* Kremp. is typified on the *T. villosus*-part. *Teloschistes* species with multiseriate hairs, spores with short septa, thick and strongly conglutinated cortex hyphae with thick walls, lacking cilia or rhizines, and which have a mainly northern hemispheric distribution, are transferred to *Seirophora austroarabica* (Sipman) Frödén, *Seirophora contortuplicata* (Ach.) Frödén, *Seirophora lacunosa* (Rupr.) Frödén, *Seirophora seirophora stenophylla* (Tav.) Frödén, and *Seirophora villosa* (Ach.) Frödén.

Key words: lichens, Seirophora, Teloschistaceae, Teloschistes, typification

Introduction

During our work with nomenclature and generic delimitation of Teloschistaceae, we had the opportunity to study the type material of Physcia magara Kremp., which is the type species of the genus Seirophora Poelt. The species was originally described by Krempelhuber in 1868. He based his description on material located in the Vienna herbarium (W) which had been collected by Kotschy and issued without description as Parmelia magara in his exsiccatum Iter Syriacum in 1855. In the protologue, Krempelhuber (1868) cited the two specimens he saw in W: nos 1012 and 1112. Stitzenberger (1890) included Physica magara in his African lichen flora without any special comments, but in the second supplement (1895), he stated that it was very closely related to Teloschistes villosus (Ach.) Norman. In his world monograph of Teloschistes, Hillmann (1930) concluded that Physica magara was most probably a

member of Teloschistaceae, but due to the scarce material, he was uncertain whether it belonged to Teloschistes or a new genus. The reasons for assigning it to the *Teloschistaceae* were the polarilocular spores, the K+ red reaction of the apothecia and the interspersed granules in the epithecium. He strongly opposed Stitzenberger's (1895) view on its relation to T. villosus, and in fact accused him of not having seen any material at all of Physica magara. Poelt (1983) erected a new genus for this taxon named Seirophora. including only the type species. Poelt did not explicitly record what material he studied, but it appears that he studied only the single specimen collected by Kotschy (no. 1112) located in M, which he chose as lectotype (erroneously called holotype), and an additional, sterile specimen collected by Schweinfurth on the Libyan coast in 1883 (also in M). Nevertheless, due to its peculiar strands of prosoplectenchymatic, chondroid tissue embedded in the medulla combined with typical apothecial characters of Teloschistaceae, he decided it was distinct enough to form a genus of its own in Teloschistaceae. Kärnefelt (1989) followed the views of Poelt,

P. Frödén (corresponding author) and P. Lassen: Botanical Museum, Lund University, Ö. Vallg. 18, SE-22361 Lund, Sweden.

but noted the close affinity to *T. lacunosus* (Rupr.) Savicz, thereby incorporating Stitzenberger's view, as *T. lacunosus* is closely related to *T. villosus*.

Material and Methods

Material of Seirophora and Teloschistes from the following herbaria have been studied: ASU, BM, Bratt priv. hb., C, E, G, GB, H, H-ACH, H-NYL, HBG, LD, Litterski priv. hb., M, MIN, S, UPS, and W. Morphological characters were studied with a dissecting binocular microscope with $\times 40$ magnification. Anatomical structures were studied with a Zeiss Axioscope light microscope with and without interference contrast. Macro photographs were taken with a Canon Eos 300D digital camera with a 50 mm macro lens, and micro photographs were taken with an Olympus DP 11 digital camera. The photographs were subsequently processed in Photoshop 7. Measurements of the spores (×1000 magnification) were carried out without interference contrast on hand cut sections mounted in distilled water. The cortex hyphae (\times 1000 magnification) were first treated with a 10% KOH solution, which then was replaced by distilled water. The type material of Physcia magara and material of Ramalina maciformis have been investigated with spot tests; using KOH, C, and Pd.

Taxonomy

The type material

According to a note on the designated lectotype specimen, it had been in Krempelhuber's personal herbarium and was transferred to M in 1883 after his death. Krempelhuber (1868) only mentioned that he studied the two specimens in W and does not refer to any material in his private herbarium. The lectotype also lacks the original, partly printed, exsiccatum label. However, the text on the handwritten label of the lectotype reads the same as in the original label, save for "Parmelia magara El Arysch" at the bottom, which is lacking (Fig. 1). Most probably the lectotype is a part taken from the original specimen no. 1112 in W. The fact that specimen no. 1012 is in better condition than no. 1112 also suggests that material has been removed from it. If this assumption is correct, the typification made by Poelt was indeed made on the original material, although it would have been more

appropriate if the lectotype was chosen from one of the syntypes in W. Specimen no. 1112 in W is thus to be regarded as an isolectotype and specimen no. 1012 as a syntype.

The identity of Physcia magara

The lectotype consists of two thalli (Fig. 2), clearly separated, but treated as belonging to the same taxon by Poelt (1983) and Kärnefelt (1989). The original material in W is composed of several thalli, but was also treated as one species by Krempelhuber (1868) and Hillmann (1930). The larger, sterile thallus of the lectotype (left in Fig. 2) as well as the other sterile type material, on which the descriptions of the morphology and anatomy are based, has a reddish to yellowish grey colour, but is K -, which was noted by Stitzenberger (1895), Hillmann (1930), Poelt (1983), and Kärnefelt (1989). For the most part, it has no colour reactions, but in part the medulla is K+ yellow turning orange-red after a while. In addition, it is also Pd+ orange-red. These colour reactions indicate either norstictic acid, stictic acid, or salazinic acid. Furthermore, it is corticated all around, without hairs, and has reddish chondroid strands in the medulla. The smaller thallus of the lectotype (right in Fig. 2, Fig. 3) and the other fertile material of the types hold all the apothecia and is thus the base for including Seirophora in Teloschistaceae. The upper surface bears short hyaline hairs. The lower side is cracked, exposing the medulla with remaining cortex strands forming a reticulate pattern. These cortex strands are superficially similar to the chondroid strands of the sterile material, but are white, not free from the cortex, and are of different composition. The K+ red reaction of the apothecium disc was reported by Hillmann (1930), Poelt (1983), and Kärnefelt (1989). It lacks the K+ and Pd+ reactions of the medulla completely. The characteristics of the two parts are summarized in Table 1. In our view, the two parts of the lectotype and the rest of the type material clearly represent two different taxa and it is also clear that Krempelhuber's

The Kotschig . Her Syriacum 1855. Physeia Magara (Katshi) Walls speinov. Nº MIZ Preseit ad El Brysch in confinies Regipti et Byrsae Paris Desortis Th. Kotschy. Iter Syriacum 1855. N: 1112 Physisa Magara (hotsily) Krphik Crescit ad El arytoh in confiniis legyptich grine, alt. ped. Die locis defertis. 1. Tarmetia magara el arytoh!

FIG. 1. Physcia magara, labels of the lectotype (Kotschy 1112, M) and the isolectotype (Kotschy 1112, W).

(1868) protologue is based on two taxa. The original drawings of the protologue, most of which are referable to thalli in the syntype, also confirm this (Fig. 4).

The morphology, anatomy, and chemical reactions indicate that the larger thallus of the lectotype as well as the other sterile material of the types belong to *Ramalina maciformis* (Delile) Bory. This species was described from Egypt (Delile 1813), not far from where *Physcia magara* was collected. The sterile material of the types agree well with the available material of *R. maciformis* in LD. *Ramalina maciformis* usually contains norstictic acid+bourgeanic acid or salazinic acid+bourgeanic acid, but acid deficient strains occur (Krog & Østhagen 1980).

Follmann (1976) has also reported partly acid deficient specimens from Israel. The Libyan material cited by Poelt (1983) shows strong colour reactions in the medulla with K and Pd. The smaller thallus of the lectotype as well as the other fertile material of the isotype and syntype, belong to Teloschistes villosus (Ach.) Norman. As the type consists of two different taxa which is also reflected in the protologue, an additional typification needs to be made. Thus, Physcia magara is here typified on the smaller, fertile T. villosus part (Fig. 3 and on the right in Fig. 2) of the lectotype (M). Physcia magara is antedated by Parmelia villosa Ach. (Acharius 1803), and is reduced to synonymy with the latter.



FIG. 2. Physcia magara (Kotschy 1112, M), the two thalli of the lectotype.



FIG. 3. Physcia magara (Kotschy 1112, M), the fertile part on the right (in Fig. 2) chosen here as lectotype.

292

Character	Sterile part (R. maciformis)	Fertile part (T. villosus)
Colour	Yellowish grey to reddish all around	Grey on upper side, white on lower
Surface	Naked	Villose on upper side, naked on lower
Cortex	Corticated all around	Upper side corticated, lower cracked exposing the medulla with remaining cortex strands forming a reticulate pattern
Chondroid strands	Present, reddish, separated from cortex	Not present, lower cortex forming a reticulate pattern
Spot test	Partly PD+ orange red, K+ yellow turning red	No colour reactions on thallus, apothecia K+ red

TABLE 1. Comparison between the two parts of the type material of Physcia magara



FIG. 4. Physcia magara, Krempelhuber's original drawings. The two Figs. 2 represent Ramalina maciformis, Fig. 2a and 2b represent Teloschistes villosus.

Emendation of Seirophora

As was suggested in Frödén *et al.* (2004), some species should be separated from *Teloschistes*. With the new typification of *Seirophora*, the name becomes available for this group. *Seirophora* differs from *Teloschistes* s. str. in a number of characters, the principal ones being summarized in Table 2. The species of *Seirophora* have no cilia or rhizines. All species of *Teloschistes* produce these, although in some cases only sparingly so, as for example in *T. cymbalifer* (Eschw.) Müll. Arg. All species of *Seirophora* produce multiseriate, complex hairs, consisting of strongly conglutinated hyphae (Fig. 5A).

Character	Seirophora	Teloschistes s. str.
Cilia/rhizines	Absent	Present in all species
Spores	Short septa (2–4 µm in water)	Long septa $(4-10 \mu\text{m in water})$
Hairs	Simple and complex (multiseriate) sometimes branched	Mostly simple, consisting of a single row of cells, rarely complex and then more loosely conglutinated
Cortex hyphae	Strongly conglutinated after treatment with KOH, thick (3–7 μ m) with thick walls (<i>c</i> . 1–2 μ m)	More lax and separating after treatment with KOH, thin $(1.5-3 \ \mu m)$ with thin walls (c. 0.5 $\ \mu m$)
Conidia	Bacilliform to narrowly ellipsoid	Bacilliform, bacilliform to narrowly ellipsoid, or bifusiform
Chemistry*	Chemosyndrome A	Chemosyndrome A, A+1, A+2, A+3, A+4, A+5, A3, A3+2, or A3+3
Distribution	Mainly northern hemispheric	Mainly southern hemispheric

TABLE 2. Comparison between Seirophora and Teloschistes s. str.

*Chemosyndromes are referable to Søchting & Frödén (2002).

Simple hairs made up of single rows of cells are also produced. The species of Teloschistes either lack hairs or mainly produce simple hairs, single or loosely conglutinated, but in some species such as T. puber (Ach.) Almb. the complex type is also formed, although shorter and less conglutinated than in Seirophora (Fig. 5B and 5C). Seirophora species have spores with short septa (2–4 μ m in water); in *Teloschistes* they are $4-10 \,\mu\text{m}$ (Fig. 6). In T. hypoglacus (Nyl.) Zahlbr., which has quadrilocular spores, the septa are only c. $1.5-2 \,\mu\text{m}$, but the total length between the end locules is 9-10 µm. Although both genera have a prosoplectenchymatic cortex, the cortical hyphae in Seirophora, measured after treatment with KOH, are thicker $(3-7 \,\mu m)$ with thicker walls (c. 1–2 μ m) than in *Teloschistes* (1·5–3 μ m, and walls c. $0.5 \,\mu\text{m}$) (Fig. 7). The cortical hyphae in Seirophora remain virtually unchanged after the treatment while they become somewhat lax and separated in Teloschistes. All species of Seirophora investigated have narrowly ellipsoid to bacilliform conidia and chemosyndrome A (Søchting & Frödén 2002). The variation in Teloschistes is much greater; bacilliform, bacilliform to narrowly ellipsoid, and bifusiform conidia occur and several chemosyndromes have been found (Søchting & Frödén 2002).

Seirophora species occur only in the northern hemisphere, except for the outlying populations of S. villosa (Ach.) Frödén in Peru and Chile. They occur over large parts of Asia, southern parts of Europe, northern Africa, and western North America up to the Canadian Arctic and Greenland. The centre of diversity for Teloschistes, on the other hand, is in southern Africa, southern Australia and New Zealand, and South America. While Seirophora has a Laurasian distribution, Teloschistes s. str. has a distribution pattern typical of groups of organisms originating in Gondwanaland (see Galloway 1991 and references therein). This is probably due to the area of origin of the two genera as well as their preference for rather dry areas, but with elevated local air humidity. Apparently, most species of Seirophora and Teloschistes do not cross the tropical and subtropical regions and establish themselves in the suitable areas on the other side of these regions. Only three widespread species of Teloschistes occur in the areas derived from Laurasia, and these are also the most widespread Teloschistes species in the southern Hemisphere.

DNA-data from the nuclear ribosomal ITS1-5.8S-ITS2 region also fully support the separation of *Seirophora* from *Teloschistes* (P. Frödén, unpublished data).



FIG. 5. Seirophora and Teloschistes, hairs. A, multiseriate hairs of S. lacunosa (V. P. Savicz: Lichenotheca Rossica decas 5 no 44, Kazakhstan, Kumaschkino, Keller & Lejsle s. n., LD); B, simple hairs of T. puber (South Africa, Melkbosch strand, Garside 5040a, LD); C, loosely conglutinated hairs of T. puber (same specimen as B). Scales: A, B & C=5 μm.



FIG. 6. Seirophora and Teloschistes, spores. A, S. lacunosa (Spain, Andalusia, Punta del Sabinar, Søchting 9649, C); B, T. flavicans (Mexico, 32 miles SE of Gusave, Nash 12262, ASU). Scales: A & B=5 μm.

The species taxonomy of *Seirophora* is in need of a revision but this is beyond the scope of the present study. Morphological and molecular work is underway and detailed descriptions of the taxa and keys will be presented in a future paper.

Seirophora Poelt emend. Frödén

Flora **174:** 440 (1983); type: *Seirophora magara* (Kremp.) Poelt [=*Parmelia villosa* Ach.]

Seirophora aurantiaca (R. Br.) Frödén comb. nov.

Borrera aurantica R. Br., Chloris Melvilliana: 47 (1823).—Teloschistes arcticus Zahlbr., Catal. Lich. Univ. 7: 311 (1931), nomen novum for Borrera aurantica R. Br.



FIG. 7. Seirophora and Teloschistes, cortex. A, S. lacunosa (V. P. Savicz: Lichenotheca Rossica decas 5 no 44, Kazakhstan, Kumaschkino, Keller & Lejsle s. n., LD); B, T. exilis (Equador, Chimborazo, 5 km SE of Pungala, Arvidsson, Lindqvist & Lindström 6480, GB). Scales: A & B=5 μm.

(non Teloschistes aurantiacus (Lightf.) Norman); type: Winter Harbour, 23 July 1820 (BM!—lectotypus selected here); other original material: Melville Island, leg. Sabine (E!); Melville Island (BM!, H-NYL 32840!). Borrera aurantiaca R. Br. was described by Robert Brown in Chloris Melvilliana (1823), which is an independent paginated preprint of his work published a year later (Brown

1824). As the combination Teloschistes aurantiacus (Lightf.) Norman had already been made, Zahlbruckner (1931) gave it a new name, T. arcticus Zahlbr. Lambert (1966), who rediscovered the species, wrote "the type specimen" in the legend to Fig. 1, which is a picture of S. aurantiaca (called T. arcticus), but this is apparently a typographic error. He is really referring to Fig. 2, which is the type of T. nodulifer (Nyl.) Hillmann. The only material of S. aurantiaca that is cited, apart from his own material, is a specimen from H-NYL (no. 32840). This specimen is possibly type material, but it is definitely not the specimen depicted in Fig. 1. Most probably Fig. 1 in Lambert (1966) is one of his own specimens and he was not intentionally making a typification of Borrera aurantiaca. The lectotype designated here has been in Robert Brown's personal herbarium and was collected at Winter Harbour on Melville Island.

Seirophora austroarabica (Sipman) Frödén comb. nov.

Teloschistes austroarabicus Sipman, Willdenowia **32**: 128 (2002); type: Oman, Dhofar, Jabal Nuss at Ras Nuss, 17°14'N, 55°12'E, 300 m, dry, leeward NW slopes, on Boscia arabica, 7 x 1998, N. Kilian & P. Hein NK5545a (B—holotypus not seen); Oman, Dhofar, Jabal Nuss E of Jabal Samhan, N of village Hadbin, coord. 17°14'N, 55°13'E, alt. 640 m rocky slopes with monsoon-affected vegetation, epiphyte on Euphorbia balsamifera, 7 x 1998, P. Hein & N. Kilian PH5479a (B!—paratypus).

Seirophora californica (Sipman) Frödén comb. nov.

Teloschistes californicus Sipman, Willdenowia 23: 312 (1993); type: Mexico, Baja California, near km 22 on road from Ciudad Constitucion to San Carlos (Route 22), 24°59'N, 111°51'W, on branches in desert scrub on sandy plain with Acacia constricta, Fouquieria diguetii, etc., 30 m, 8 i 1989, Sipman 25056, LICHENOTHECA LATINOAMERICANA a museo botanico berolinensi edita fasc. 2, curavit H. Sipman anno 1993, No. 98 (B—holotypus not seen; LD!, S!, and UPS!—isotypi).

Seirophora contortuplicata (Ach.) Frödén comb. nov.

Parmelia contortuplicata Ach., Synops. Meth. Lich: 210 (1814).—Teloschistes contortuplicatus (Ach.) Clauzade &

Rondon, in Vězda, *Sched. ad Lich. Sel. Exs.*: nr. 149 (1962); type: Mons Cenis (H-ACH 1303!—lectotype selected here), France Mons Cenis (UPS L00728!— isotype).

Seirophora lacunosa (Rupr.) Frödén comb. nov.

Ramalina lacunosa Rupr., Mém. Acad. Sci. St-Pétersb. 6-me ser. t. 6 (Sci nat., t. 5): 235 (1845).—Teloschistes lacunosus (Rupr.) Savicz, Acta Inst. Bot. Acad. Sci. URSS 2(2): 313 (1934); type: "Ad terram nudam, argillaceam deserti propre Barchotskoi, Mys." (LE? not seen).

Seirophora scorigena (Mont.) Frödén comb. nov.

Evernia scorigena Mont., apud Barker-Webb & Berthelot, Hist. Natur. Iles Canar., t. 3, 2. part: 97 (1840).—Teloschistes scorigenus (Mont.) Vain., Catal. Welwitsch Afric. Plants vol. 2: 407 (1901); type: "Super scorias vulcanias in lingua terræ vulgo Punta de Melenara dicta insul Canariæ nec alibi legerunt clar. Webb, Berthelot et Despréux" (FI or PC? —not seen).

Seirophora stenophylla (Tav.) Frödén comb. nov.

Teloschistes villosus v. stenophyllus Tav., Revista Biol. 4: 142 (1964).—Teloschistes stenophyllus (Tav.) Sipman, Willdenowia 32: 130 (2002); type: Cabo verde, S. Vicente, Monte Verde, on the rocks, at about 750 m. s. m. K. Byström, 10 v 1959 (LISU 862—holotype not seen).

Seirophora villosa (Ach.) Frödén comb. nov.

Parmelia villosa Ach., Meth. Lich.: 254 (1803).— Teloschistes villosus (Ach.) Norman, Nyt Mag. Naturvid. 7(3): 17 (1852 preprint), Nyt Mag. Naturvid. 7(3): 229 (1853); type: Lusitania (H-ACH 1452!—lectotype selected here).

Physcia magara Kremp., Verh. zool.-bot. Gesellsch. Wien 18: 323 (1868).—Seirophora magara (Kremp.) Poelt, Flora 174: 440 (1983); type: Crescit ad El Arysch in confiniis Aegypti et Syriae, locis desertis, Kotschy Iter Syriacum no 1112 (M!—lectotype selected here [see discussion above]; W!—isolectotype); Crescit ad El Arysch in confiniis Aegypti et Syriae, locis desertis, Kotschy Iter Syriacum no 1012 (W!—syntypus).

Physcia villosa f. brevior Nyl., Actes Soc. Linn. Bordeaux 25: 61 (1864). — Teloschistes brevior (Nyl.) Hillmann, Flora 69: 320 (1930); type: Aegyptus, In Lycio afro propre Bir Krer et alibi. G. Ehrenberg (H-NYL 32749!—lectotype selected here). Thanks are due to G. Zijlstra in Utrecht (U) for help with literature, the curators of the herbaria for supplying the material, O. Vitikainen (H) for assisting with the typifications, M. Westberg and two anonymous referees for comments on the manuscript, and I. Kärnefelt for putting the lectotype of *Physcia magara* at our disposal.

References

- Acharius, E. (1803) Methodus qua omnes detectos Lichenes. Stockholm.
- Brown, R. (1823) Chloris melvilliana. A list of plants collected in Melville island (latitude 74°–75°N. longitude 110°–112°W.) in the year 1820; by the officers of the voyage of discovery under the orders of Captain Parry. With characters and descriptions of the new genera and species. London.
- Brown, R. (1824) A list of plants collected in Melville Island, by the officers of the expedition; with characters and descriptions of the new species. A supplement to the appendix of Capt. Parry's first voyage, No 11: 261–310. London.

Delile, A. (1813) Flore d'Égypte. Paris.

- Follmann, G. (1976) Zur Nomenklatur der Lichenen 3. Über Desmazeria Mont. (Ramalinaceae) und andere kritische Verwandtschaftskreise. Philippia 3: 85–89.
- Frödén, P. Ryan, B. D. & Kärnefelt, I. (2004) Teloschistes. In Lichen Flora of the Greater Sonoran Desert Region Vol. 2 (T. H. Nash III, B. D. Ryan, P. Diedrich, C. Gries & F. Bungartz, eds). Tempe, AR: Lichens Unlimited (in press).
- Galloway, D. (1991) Phytogeography of Southern Hemispheric lichens. In *Quantitative Approaches to*

Phytogeography (P. L. Nimis & T. J. Crovello, eds): 233–262. Dordrecht: Kluwer Academic Publishers.

- Hillmann, J. (1930) Studien über die Flechtengattung Teloschistes. Hedwigia **69:** 303–343.
- Kärnefelt, I. (1989) Morphology and phylogeny in the *Teloschistales. Cryptogamic Botany* 1: 147–203.
- Krempelhuber, A. v. (1868) Exotische flechten. Verhandlungen der kaiserlich-königlichen zoologischbotanischen Gesellschaft in Wien 18: 303–330.
- Krog, H. & Østhagen, H. (1980) The genus Ramalina in the Canary Islands. Norwegian Journal of Botany 27: 255–296.
- Lambert, J. D. H. (1966) Teloschistes articus Zahlbr. Reaffirmed in the Canadian Arctic. Bryologist 69: 358–361.
- Poelt, J. (1983) Musterbeispiele analoger Lagerdifferenzierrung bei Flechten: Almbornia, Speerschneidera, Seirophora gen.nov. Flora 174: 439–445.
- Søchting, U. & Frödén, P. (2002) Chemosyndromes in the lichen genus *Teloschistes (Teloschistaceae, Lecanorales) Mycological Progress* 1(3): 257–266.
- Stitzenberger, E. (1890) Lichenæa Africana. Jahresberichten der St. Gallischen Naturwissenschaftlichen Gesellschaft 1888/1889, 1890/1891: 105–249. St. Gallen.
- Stitzenberger, E. (1895) Supplementa ad lichenæam africanam 2. Jahresberichten der St. Gallischen Naturwissenschaftlichen Gesellschaft 1893/1894: 215– 264. St. Gallen.
- Zahlbruckner, A. (1931) Catalogus lichenum universalis 7. Leipzig: Borntraeger.

Accepted for publication 7 June 2004