

On the effects of *Streptococcal* NAD⁺-glycohydrolase and Streptolysin O on macrophages

ELSA WESTERLUND

FACULTY OF MEDICINE | LUND UNIVERSITY





On the effects of *Streptococcal* NAD⁺-glycohydrolase and Streptolysin O on macrophages

Elsa Westerlund



LUND
UNIVERSITY

DOCTORAL DISSERTATION

by due permission of the Faculty of Medicine, Lund University, Sweden.
To be defended on October 16 2019 at 13:00 in Segerfalksalen, BMCA10,
Sölvegatan 19, Lund.

Faculty opponent

Thomas Henry, PhD

Centre for Infectiology Research (CIRI), Lyon, France

Organization LUND UNIVERSITY Section for Immunology Department of Experimental Medical Science SE-221 84 LUND SWEDEN Author: Elsa Westerlund	Document name Doctoral Dissertation	
	Date of issue 2019-10-16	
	Sponsoring organization	
Title On the effects of <i>Streptococcal</i> NAD ⁺ -glycohydrolase and Streptolysin O on macrophages		
Summary <p>The human pathogen <i>Streptococcus pyogenes</i> (GAS) causes both superficial infections, such as strep throat, and invasive infections, such as necrotizing fasciitis, and is responsible for about half a million deaths each year. The number of GAS infections have increased since the 1980's, due to the emergence of a M1T1 strain that has become widely disseminated. Increased expression of Streptolysin O (SLO) and NAD⁺-glycohydrolase (NADase) has been correlated with the emergence of this strain. This thesis has focused on how these two virulence factors affect cytokine release from macrophages, an innate immune cell important for the host's defense against GAS.</p> <p>In Paper I we describe a novel function for NADase: inhibition of IL-1β release after NLRP3 inflammasome activation. This inhibitory effect was mediated by extracellularly located NADase, which represents a novel functional niche for this enzyme.</p> <p>In Paper II, we explore this effect further and show that NADase inhibits an unconventional IL-1β release pathway that is dependent on the P2X7 receptor and membrane permeabilization. Interestingly, we see that IL-1β release in response to GAS is independent of the pore-forming protein GSDMD.</p> <p>In Paper III we show that pro-IL-1β is ubiquitinated and degraded during GAS infection. Pro-IL-1β ubiquitination requires the presence of SLO and results in heterotypic linkage types. The broad PI3K inhibitor 3-MA rescues pro-IL-1β degradation.</p> <p>Paper IV focuses on NADase and the observation that NADase binds macrophages in the absence of SLO, in contrast to what has been shown previously. Recombinant NADase binds macrophages in the absence of other bacterial proteins and induces cytokine release that requires TLR4 and CD14 and is dependent on MyD88 and TRIF.</p>		
Key words Innate immunity, macrophage, <i>Streptococcus pyogenes</i> , IL-1 β , NAD ⁺ -glycohydrolase		
Classification system and/or index terms (if any)		
Supplementary bibliographical information		Language English
ISSN and key title 1652-8220		ISBN 978-91-7619-833-9
Recipient's notes	Number of pages 84	Price
	Security classification	

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature  Date 2019-09-10

On the effects of *Streptococcal*
NAD⁺-glycohydrolase and
Streptolysin O on macrophages

Elsa Westerlund



LUND
UNIVERSITY

Cover photo: *Streptococcus pyogenes* colonies on a blood agar plate, taken by Elsa Westerlund

Copyright pp 1-84 Elsa Westerlund

Paper 1 © The Authors

Paper 2 © The Authors

Paper 3 © The Authors

Paper 4 © by The Authors (Manuscript)

Department of Experimental Medical Science
Lund University, Faculty of Medicine
Doctoral Dissertation Series 2019:104

ISBN 978-91-7619-833-9

ISSN 1652-8220

Printed in Sweden by Media-Tryck, Lund University
Lund 2019



Media-Tryck is an environmentally certified and ISO 14001:2015 certified provider of printed material. Read more about our environmental work at www.mediatryck.lu.se

MADE IN SWEDEN 

Table of contents

Abbreviations	7
Popular scientific summary	11
Populärvetenskaplig sammanfattning	13
Introduction to the immune system.....	15
Recognition of pathogens and danger signals.....	17
Surface-associated PRRs	18
TLR4 and LPS, recognition and signaling	18
Intracellular PRRs.....	19
Inflammasomes	19
The NLRP3 inflammasome.....	20
Non-canonical and alternative inflammasomes and macrophage hyperactivation	22
Downstream effects of inflammasome activation	23
Microbial inflammasome modulation strategies	24
Ubiquitination.....	25
Ubiquitin and its linkage types	25
Ubiquitin addition, removal and recognition.....	25
Regulation of inflammasome activation by ubiquitin.....	26
The P2X7 receptor.....	27
Receptor structure, genetics and activation	27
Genetic variants	28
Macropore formation	29
Modulation of activation.....	30
Downstream effects.....	30
IL-1 β and its release pathways	32
Non-vesicular release (Unconventional type I release).....	33
GSDMD pores and membrane permeabilization.....	33
Non-vesicular release by other proteins.....	34
Vesicular release (Unconventional type III release)	35
Secretory lysosomes	35

Microvesicles and exosomes	36
Secretory autophagy	37
IL-1 β release as a spectrum reflecting signal strength	37
Group A <i>Streptococcus</i>	38
Pathogenesis	39
Adherence and colonization	39
Invasion and evasion.....	40
M1T1 GAS	41
The evolution of epidemic M1T1 GAS	41
SLO, β -NAD ⁺ -glycohydrolase and cytolysin-mediated translocation	42
The host response to GAS	44
Cytokines and PRRs	44
Adaptive immune responses.....	45
Other host factors influencing infection outcome.....	45
The role of IL-1 β during GAS infection	46
Present investigation	48
Original papers and manuscripts.....	48
Summary and discussion of papers	49
Paper I: Inhibition of Inflammasome-Dependent Interleukin 1 β	
Production by Streptococcal NAD ⁺ -Glycohydrolase: Evidence for	
Extracellular Activity.....	49
Paper II: The Secreted Virulence Factor NADase of Group A	
<i>Streptococcus</i> Inhibits P2X7 receptor-mediated Release of IL-1 β	50
Paper III: Streptolysin O Induces Ubiquitination and Degradation of	
pro-IL-1 β	52
Paper IV: <i>Streptococcal</i> β -NAD ⁺ -glycohydrolase binds to and induces	
cytokine release from macrophages in the absence of Streptolysin O	54
Acknowledgements.....	57
References	59

Abbreviations

3-MA	3-methyladenine
ABC	ATP-binding cassette
ADP	Adenosine di-phosphate
ADP-r	ADP-ribose
AIM2	Absent in melanoma 2
AMP	Anti-microbial peptide
ALR	AIM2-like receptor
APC	Antigen-presenting cell
ASC	Apoptosis speck-like protein containing a CARD
ATP	Adenosine tri-phosphate
BCR	B cell receptor
CARD	Caspase activation and recruitment domain
CD	Center of differentiation
CDC	Cholesterol-dependent cytolysin
cGAS	cyclic GMP-AMP synthase
CLR	C-type lectin receptor
CMT	Cytolysin-mediated translocation
CovRS	Control of virulence regulatory system
DAMP	Damage-associated molecular pattern
DC	Dendritic cell
DUB	Deubiquitinase enzyme
FADD	Fas-associated protein with a death domain
FGF-2	Fibroblast growth factor 2

GAC	Group A carbohydrate
GAS	Group A Streptococcus (<i>Streptococcus pyogenes</i>)
GSDMD	Gasdermin D
GSK	GlaxoSmithKline
HAMP	Homeostasis-altering molecular process
HGT	Horizontal gene transfer
IFS	Immunity factor for <i>Streptococcus pyogenes</i> NADase
IKK	I κ B kinase
IL	Interleukin
IL-1R	IL-1 receptor
IL-1Ra	IL-1 receptor antagonist
IFN	Interferon
IRAK	IL-1 receptor associated kinase
IRF	IFN regulatory factor
K	Lysine
K ⁺	Potassium
LBP	LPS-binding protein
LDH	Lactate dehydrogenase
LPS	Lipopolysaccharide
LRR	Leucine-rich repeat
MAPK	Mitogen-activated protein kinase
Mg ²⁺	Magnesium
MHC-II	Major histocompatibility complex II
MMP	Matrix metalloproteinase
MVB	Multivesicular bodies
MyD88	Myeloid differentiation primary response 88
β -NAD ⁺	β -nicotinamide adenine dinucleotide
NADase	β -NAD ⁺ -glycohydrolase
NAIP	NLR family of apoptosis inhibitory proteins

NAM	Nicotinamide
NEK7	NIMA-related kinase 7
NF	Necrotizing fasciitis
NF- κ B	Nuclear factor kappa-light-chain-enhancer of activated B cells
NLR	Nucleotide-binding domain, LRR-containing receptor
NLRC	NLR and CARD-containing receptor
NLRP	NLR and PYD-containing receptor
P2R	Purinergic receptor
P2X7R	P2X7 receptor
PAMP	Pathogen-associated molecular pattern
PI	Propidium iodide
PI(4,5)P ₂	Phosphatidylinositol 4,5-bisphosphate
POP	PYD-only protein
PRR	Pattern-recognition receptor
PYD	Pyrin domain
RIPK	Receptor-interacting serine/threonine-protein kinase
RIG-I	Retinoic acid-inducible gene I
RLR	RIG-I-like receptor
ROS	Reactive oxygen species
SAg	Superantigen
SLO	Streptolysin O
SLS	Streptolysin S
SNP	Single nucleotide polymorphism
Spe	Streptococcal pyrogenic exotoxin
STING	Stimulator of IFN gene
STSS	Streptococcal toxic shock syndrome
TAK1	TGF- β -activated kinase 1
TCR	T cell receptor
TIR	Toll-IL-1 receptor

TIRAP	TIR-containing adaptor protein
TGF- β	Transforming growth factor β
TLR	Toll-like receptor
TNF	Tumor necrosis factor
TRAF6	TNF-associated factor 6
TRAM	TRIF-related adaptor molecule
TRIF	TIR domain-containing adaptor-inducing IFN- β
UBD	Ubiquitin-binding domain
wt	Wild-type

Popular scientific summary

Our immune system faces a multitude of challenges daily: it not only needs to notice differences between our own cells and those of invaders, it is also required to recognize whether a microorganism is one of our harmless symbionts or if it has the potential to cause inflammation and disease. To fight breaches in our defense we need patrolling soldiers ready to battle a wide range of enemies at a moment's notice; to resolve conflicts we need agents that are specially trained for each case. Our body has solved these issues by splitting the immune system into two parts: the innate and the adaptive. Cells of the innate immune system are trained to recognize characteristics common to a wide range of bacteria and viruses, enabling them to quickly mount a protective response and alert the adaptive immune system. The adaptive cells are specially trained to recognize specific pathogens and able to make clones of themselves when activated; the whole troop is then deployed to the site of combat.

Numerous studies on pathogenic microorganisms have shown us that we are part of a continuous arms race, but it is seldom clear who has the upper hand. This thesis has studied a small part of this arms race, where a bacterium known as Group A *Streptococcus* (GAS, Latin name *Streptococcus pyogenes*) uses two weapons, so-called virulence factors, to dampen the levels of alarm signals released by macrophages, a type of innate immune cell.

GAS gives rise to a wide range of diseases, from common and mild strep throat and impetigo to potentially life-threatening and invasive infections such as necrotizing fasciitis (known as the “flesh-eating disease”) and toxic shock syndrome. With this repertoire it's not surprising to learn that GAS is high up on WHO's list of infection-related deaths worldwide. Antibiotic resistance in GAS is not (yet) a problem, but the invasive diseases often progress faster than the effect of antibiotics. There's been an increase in invasive GAS disease cases since the 1980's. Bacteria isolated from these unfortunate patients revealed that a lot of infections were caused by the same strain of GAS that had spread rapidly around the world. Genetic comparisons of this new strain and older ones showed that one of the differences between them was that the new strain produced increased levels of the two virulence factors Streptolysin O (SLO) and NADase.

Previous studies on SLO showed that it is able to kill macrophages by making large holes in them, but that macrophages on the other hand use a warning system called

the inflammasome to detect these holes and release a warning signal called IL-1 β to attract other immune cells. Our studies have shown that NADase works to dampen the amount of IL-1 β released, by somehow inhibiting a macrophage surface molecule called P2X7. We also discovered that SLO can cause breakdown of IL-1 β inside the macrophage. That both SLO and NADase affected IL-1 β was very interesting, since this signal molecule has been shown to be very important in combating GAS. This could mean that the bacterium uses SLO and NADase to prevent activation of the immune system. More research into this area would give us valuable information about the infection process, but it would also award us with increased knowledge of how our immune cells work.

Populärvetenskaplig sammanfattning

Vårt immunförsvar står inför många olika prövningar varje dag: det måste inte bara kunna göra skillnad på kroppens egna celler och inkräktare, utan behöver också kunna skilja på mikroorganismer som är våra ofarliga hyresgäster och de som har potential att orsaka sjukdomar. När våra barriärer rämnar behövs patrullerande soldater som är redo att slåss och slå larm vid första anblick av hot, men för att lösa konflikterna behövs också en trupp agenter som är specialtränade för varje fall. Vår kropp har löst detta genom att dela upp immunförsvaret i två delar: det medfödda och det adaptiva. Cellerna som tillhör det medfödda immunförsvaret har tränats för att känna igen varningssignaler gemensamma för många olika sorters bakterier och virus, vilket gör att de snabbt kan börja oskadliggöra hoten och skicka varningssignaler till det adaptiva immunförsvaret. När en adaptiv cell, specialtränad för att känna igen just denna inkräktare, aktiveras börjar den kлона fler kopior av sig själv och den samlade truppen ger sig sedan ut i kroppen för att bekämpa hotet.

Om det är något vi lärt oss av många års studier av sjukdomsframkallande bakterier är det att vi befinner oss i en ständig kapprustning, där det sällan är klart vem som har överhanden. Den här avhandlingen handlar om en del av denna kapprustning – hur bakterien *Streptococcus pyogenes* (oftast kallad Grupp A Streptokocker (GAS)) använder sig av två vapen, så kallade virulensfaktorer, för att dämpa varningssignaler från makrofager, en del av det medfödda immunförsvaret.

GAS ger upphov till infektioner av väldigt olika karaktär, från milda sjukdomar såsom halsfluss och svinkoppor till de potentiellt livshotande (invasiva) infektionerna nekrotiserande fasciit (mer känd som köttätande sjukdom) och toxiskt chocksyndrom. Med en sådan bred sjukdomsrepertoar är det inte konstigt att GAS finns med i toppen på WHO:s lista över infektionsrelaterade dödsorsaker världen över. Som tur är går det fortfarande att behandla de flesta GAS-infektioner med antibiotika, men när det gäller de invasiva infektionerna är sjukdomsförloppet ofta för snabbt för att det ska hinna verka. Sedan 1980-talet har det skett en ökning i antal rapporterade fall av invasiva infektioner. Bakterier tagna från dessa patienter visade att många av sjukdomsfallen orsakats av en och samma bakteriestam som snabbt spridits världen över. Genom genetiska jämförelser av denna stam och äldre stammar har man kommit fram till att det som skiljer dem åt är bland annat att den nya stammen producerar mer av de två virulensfaktorerna Streptolysin O (SLO) och NADase.

Tidigare studier har visat att SLO kan döda makrofager genom att göra stora hål i dem, men makrofagerna kan å andra sidan upptäcka dessa hål genom ett varningssystem som kallas inflammasomen och snabbt utsöndra en signalsubstans, IL-1 β , som lockar till sig andra celler ur immunförsvaret. Våra studier har visat att NADAs kan minska mängden IL-1 β som utsöndras, och att detta sker genom att NADAs på något sätt hämmar en av makrofagens ytmolekyler som kallas P2X7. Vi har också sett att SLO kan göra så att IL-1 β bryts ner inne i cellen istället för att utsöndras. IL-1 β har tidigare visats spela en viktig roll i att dämpa GAS-infektioner, så att det var just den signalsubstansen som både SLO och NADAs påverkade var väldigt intressant, eftersom det skulle kunna innebära att bakterien använder dem för att undgå att upptäckas av immunförsvaret. Vidare forskning i exakt hur SLO och NADAs hämmar IL-1 β kommer ge viktig information om infektionsförloppet i stort men också göra att vi fördjupar våra kunskaper om hur våra immunceller fungerar.

Introduction to the immune system

Our first line of defense against invading pathogens consists of the physical and anatomical barriers that must be overcome in order to gain access to the host, *e.g.* our skin, the mucociliary apparatus of our respiratory tract and the low pH in our stomach. If these barriers are successfully breached, the next obstacle is our immune cells, classically divided into innate and adaptive. Innate immune cells recognize conserved signals relating to pathogens and danger molecules and are armed with antimicrobial defense systems such as phagocytosis and antimicrobial peptides (AMPs), enabling them to rapidly mount a response.¹ They are also tasked with initiating activation of adaptive immunity, leading to a highly specific immune response and the generation of persisting memory cells.²

The innate immune system includes professional phagocytes and antigen-presenting cells (APCs) such as macrophages, dendritic cells (DCs) and neutrophils.¹ They are aided by non-professional phagocytes, *e.g.* epithelial cells, and by the AMP-containing granules found in natural killer cells, eosinophils and basophils.¹ In addition, circulating macromolecules such as complement proteins and lipopolysaccharide (LPS)-binding protein (LBP) also participate in host defense and make up humoral innate immunity.^{1,3} Recognition of pathogens is mediated by a number of germline-encoded receptors that recognize conserved bacterial and viral structures.⁴ Upon activation (which will be dealt with in molecular detail in the following chapters) innate cells release cytokines and chemokines to recruit and activate circulating immune cells to the site of infection.²

Innate immune cells such as macrophages are divided into subsets depending on location, function and developmental stage. Macrophage precursors, monocytes, are derived from hematopoietic stem cells and can be found in the spleen or circulating in the blood, awaiting recruitment signals.³ Upon sensing danger, they can migrate into the tissue and differentiate into mature macrophages, joining specialized tissue-resident macrophages such as alveolar macrophages in the lung and Kupffer cells in the liver.³ Tissue-resident macrophages are derived from embryonic progenitors and their capacity for self-renewal makes sure that their numbers are constantly replenished.⁵ Macrophages can also be viewed of as being on a spectrum, where on one end M1 macrophages are involved in the response against bacteria and viruses and on the other M2 macrophages are involved in wound and tissue healing and

resolving immune responses.³ However, there is great plasticity and overlap between these subsets.

While the strength of innate immunity lies in its speed, the T and B cells of adaptive immunity instead have the potential to recognize virtually any antigen. Through a complicated process involving random rearrangements of gene segments, highly specific T cell receptors (TCR) and B cell receptors (BCR) are generated.⁶ After a rigorous selection process to weed out cells expressing *e.g.* self-reacting receptors, T and B cells migrate from primary lymphoid organs (thymus and bone marrow) to secondary lymphoid organs (spleen and lymph nodes) where they await activation by APCs or soluble antigens, leading to proliferation and differentiation.^{2,6}

Activated T cells migrate to effector sites and are roughly divided into cytotoxic or helper subsets. Cytotoxic T cells express center of differentiation (CD) 8 (CD8⁺ T cells) and are involved in killing infected host cells, while helper T cells express CD4 (CD4⁺ T cells) and aid in activating B cells, phagocytes and other T cells by releasing specific cytokines.⁶ Depending on environmental cues, such as cytokines released by APCs, CD4⁺ T cells differentiate into distinct subsets with specific functions and roles in influencing the immune response, *e.g.* by releasing a specific repertoire of cytokines.²

The humoral part of adaptive immunity is mediated by antibodies that can bind and neutralize pathogens and mediate immune cell activation.⁷ Activated B cells can differentiate into short-lived plasma cells which produce a high number of low-affinity antibodies, or they can establish germinal centers, where long-lived plasma cells generate high-affinity antibodies of different isotypes, which differ in their ability to *e.g.* activate the complement pathway and bind to APCs.⁸

The antigen-specific T and B cells that persist after a primary infection is resolved are called memory cells and ensure efficient activation of the immune system if the pathogen is encountered again, thus creating a library of all the challenges our body has faced. Despite the longevity and specificity of the adaptive immune response it is only shared among vertebrates; other multicellular organisms rely solely on innate immunity, highlighting its fundamental role in our defense system.¹

This thesis focuses on the interactions between macrophages and the human pathogen Group A *Streptococcus* (GAS; *Streptococcus pyogenes*), with particular interest in how production and release of the cytokine interleukin (IL) 1 β (IL-1 β) is regulated. As promised above, the following sections describe the events taking place when pathogens or danger signals are detected. It is followed by sections on ubiquitination and IL-1 β release pathways and ends with a description of GAS before going into our present research and results.

Recognition of pathogens and danger signals

In 1989, Charles Janeway proposed a model for how innate immunity distinguishes self from non-self and activates adaptive immune responses in response only to the latter: a set of receptors expressed by all innate cells, pattern-recognition receptors (PRRs), that would be able to recognize pathogen-specific structures, pathogen-associated molecular patterns (PAMPs).⁹ In the decade after this, a few seminal studies proved this to be true. The discovery that humans express a functional and genetical homologue of the *Drosophila melanogaster* protein Toll,¹⁰ important for the fly's antifungal responses,¹¹ gave us the first PRR but not its PAMP. The PAMP, as well as the gene encoding that first human Toll-like receptor (TLR), was resolved a year later when it was reported that a defective *tlr4* gene prevented a response against LPS.¹² Since then a multitude of studies have identified additional PRRs and their ligands, leading to a modification of Janeway's first hypothesis: PRRs are not limited to recognizing PAMPs but also react to signs of cell damage (damage-associated molecular patterns [DAMPs]) and disruption of homeostasis (homeostasis-altering molecular processes [HAMPs]).¹³ Recent efforts have been made to understand how the innate response is modulated, *e.g.* how are the commensal bacteria in our gut – which also express PAMPs – distinguished from pathogens? The presence of virulence factors that activate several types of PRRs are thought to be one clue, compartmentalization another; detection of PAMPs in the gut lumen is normal, while their presence in the lamina propria indicates that the epithelial lining has been breached and that an immune response needs to be initiated.²

The PRRs discovered so far can be divided into TLRs, C-type lectin receptors (CLRs), nucleotide-binding domain, leucine-rich repeat (LRR)-containing receptors (NLRs), retinoic acid-inducible gene-I (RIG-I)-like receptors (RLRs), absent in melanoma 2 (AIM2)-like receptors (ALRs) and stimulator of interferon (IFN) gene (STING) and cyclic GMP-AMP synthase (cGAS).¹⁴ NLRs, RLRs, ALRs, STING and cGAS survey the cytoplasm while CLRs are present in the plasma membrane; TLRs mediate signaling from both plasma membrane and intracellular endosomes and are the most well-studied class of PRRs.^{14,15}

Surface-associated PRRs

A common fate after a PRR recognizes its ligand is activation of the transcription factor nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B), which can lead to expression and release of several inflammatory cytokines.¹⁵ For CLRs such as Dectin-1 this happens in response to glucans from fungi and mycobacteria,¹⁴ while cell-surface TLRs generally recognize microbial membrane components, as exemplified by the LPS-TLR4 and flagellin-TLR5 interactions.¹⁶ TLR activation occurs after the formation of homo- or heterodimers; the ability of TLR2 to recognize diverse ligands from bacteria, viruses and fungi is thought to be due to its ability to form heterodimers with TLR1 and TLR6.¹⁵ Pattern recognition depends on the horseshoe-like extracellular domain made up of LRRs which is connected to a transmembrane domain and a cytosolic Toll-IL-1 receptor (TIR) domain required for recruitment of adaptor molecules to mediate signaling.¹⁷ Two sets of adaptor proteins mediate the downstream signaling events following TLR activation. As the events following TLR4 activation can be viewed as a prototypical TLR signaling cascade, the following section will describe this in detail.

TLR4 and LPS, recognition and signaling

As LPS is not a direct ligand for TLR4, several coreceptors and adaptor proteins are required for initiating the signaling cascade downstream of LPS sensing.¹⁸ LBP binds LPS and transfers it to the coreceptor CD14, which in turn mediates transfer to MD-2 which is bound to TLR4, initiating receptor dimerization.¹⁴ The dimerization mediates recruitment of the TIR-containing adaptor molecules TIR-containing adaptor protein (TIRAP) and myeloid differentiation primary response 88 (MyD88).^{19,20} MyD88, used by all TLRs except TLR3, forms a complex termed the myddosome with the serine/threonine kinase IL-1 receptor-associated kinase 4 (IRAK4) and IRAK1, leading to their activation.¹⁴ IRAK1, together with the E3 ubiquitin ligase tumor necrosis factor (TNF)-associated factor 6 (TRAF6), recruits and ubiquitinates a complex which includes transforming growth factor- β (TGF- β)-activated kinase 1 (TAK1).¹⁷ This allows for activation of the I κ B kinase (IKK) complex, which frees NF- κ B from its inhibitor and permits it to travel to the nucleus and induce gene expression of proinflammatory genes such as cytokines.¹⁵ The TAK1 complex also activates mitogen-activated protein kinase (MAPK) family members, leading to induction of genes controlled by the transcription factor activator protein 1, which controls processes such as cell differentiation and death.¹⁴

TLR4 is unique among the TLR family in that it can signal both from the surface and from endosomes. After CD14 mediates endosomal uptake of TLR4,²¹ the adaptor proteins TIR domain-containing adaptor-inducing IFN- β (TRIF) and TRIF-related adaptor molecule (TRAM) are recruited.^{15,19,22} TRIF interacts with TRAF6,

which recruits receptor-interacting serine/threonine-protein kinase 1 (RIPK1), leading to activation of the TAK1 kinase and the transcription factors NF- κ B and IFN regulatory factor 3 (IRF3), the former inducing expression of proinflammatory genes such as pro-IL-1 β , important for antibacterial responses, and the latter resulting in production of type I IFNs, important in *e.g.* antiviral responses.^{4,15}

Intracellular PRRs

PRRs located in endosomes or the cytosol generally detect the presence of intracellular pathogens. For example, TLR3, TLR7 and TLR9 respond to endosomal DNA and single- or double-stranded RNA, *e.g.* released by pathogens residing in endosomes, or liberated after phagocytic uptake and destruction.¹⁴ TLR9 and TLR7 induce a MyD88-dependent pathway that, apart from activating NF- κ B, also induces release of type I IFN *via* the transcription factor IRF7.^{4,17} TLR3 instead induces type I IFNs through a pathway shared with endosomal TLR4.

RLRs induce NF- κ B and IRFs in response to cytosolic viral RNA, while STING and cGAS recognize cytosolic DNA and trigger production of both type I IFNs and proinflammatory cytokines.^{14,23} The NLRs NOD1 and NOD2 induce proinflammatory genes upon detection of bacterial cell wall degradation products,¹⁵ while activation of other NLRs, and some ALRs, leads to formation of inflammasomes and production of IL-1 β ,²⁴ which will be described in detail below.

Inflammasomes

The term inflammasome was coined in 2002 to describe high-molecular weight complexes that mediate activation of inflammatory caspases.²⁵ What seemed like a simple system – a sensor, an adaptor and caspase-1 – has blossomed into a diverse field of sensors, activation signals and regulatory mechanisms. Inflammasome formation occurs when the activated sensor protein oligomerizes, allowing recruitment of the adaptor protein ASC (apoptosis-associated speck-like protein containing a CARD [caspase activation and recruitment domain]) *via* their mutual pyrin domains (PYD).²⁶ This induces ASC polymerization and the formation of a filamentous structure termed the ASC speck or pyroptosome, which serves to amplify downstream signals.²⁷ The ASC speck can also be released extracellularly to increase inflammation.²⁸ Caspase-1 is recruited to ASC *via* its CARD, leading to proximity-induced autoprocessing of caspase-1 and downstream responses: cleavage and activation of pro-IL-1 β , pro-IL-18 and the pore-forming protein Gasdermin D (GSDMD), induction of an inflammatory form of cell death called

pyroptosis and release of signaling molecules called eicosanoids from intestinal epithelial cells.^{24,29}

Five different sensor proteins have been described to form inflammasomes: the NLR proteins NLR and PYD containing receptor 1B (NLRP1B), NLRP3 and NLR and CARD containing receptor 4 (NLRC4), the ALR AIM2, and pyrin.³⁰ Proteins such as NLRP6, NLRP12 and IFN γ -inducible protein 16 have also been suggested to form inflammasomes, but the details surrounding them have not been elucidated.^{31,32}

Although not the first described, the AIM2 inflammasome best fits the classical view of PRR-PAMP activation in that AIM2 directly binds its ligand, dsDNA of host or microbial origin, and recruits ASC and caspase-1.^{33,34} In contrast, activation of NLRP1B, the first inflammasome sensor identified,²⁵ is more complicated. In response to *e.g.* lethal factor from *Bacillus anthracis*, N-terminal cleavage and partial proteasomal degradation of NLRP1B releases a C-terminal part responsible for caspase-1 activation.^{35,36} The NLRC4 inflammasome relies on NLR family of apoptosis inhibitory proteins (NAIPs) to recognize intracellular flagellin and the rod and needle subunits of type III secretion systems from Gram-negative bacteria, leading to NAIP-NLRC4 colocalization and inflammasome formation.³⁷⁻³⁹ Pyrin, the newest addition to the inflammasome family, recognizes inactivation of the GTPase RHOA caused by bacteria such as *Clostridium difficile* and can thus be classified as a HAMP sensor.^{13,40} Despite being the most well-studied, several questions still surround NLRP3 inflammasome activation and regulation, presented in detail below.

The NLRP3 inflammasome

NLRP3 contains an N-terminal PYD, a central oligomerization domain and a C-terminal LRR involved in regulation of activation.^{26,41} NLRP3 inflammasome activation has been suggested to augment Alzheimer's and Parkinson's disease and inflammatory diseases such as atherosclerosis and type 2 diabetes.³⁰ In addition, mutations in *nlrp3* give rise to cryopyrin-associated periodic syndromes, a group of hereditary autoinflammatory diseases presenting with symptoms such as recurring fever and skin rashes.³⁰

What separates it from other inflammasomes is that NLRP3 activation is a two-step process. An initial priming signal is required to license NLRP3 for activation, involving induction of post-translational modifications and an increase in production of NLRP3 and pro-IL-1 β .⁴¹ The activation signal can come from a wide range of stimuli, *e.g.* DAMPs such as adenosine triphosphate (ATP)⁴² and uric acid⁴³ and PAMPs such as pore-forming toxins⁴⁴ and nucleic acids.⁴⁵ Lastly, NLRP3 can be involved in non-canonical⁴⁶⁻⁴⁸ and alternative⁴⁹ inflammasome formation.

Priming and posttranslational modifications of NLRP3 inflammasome components

Transcriptional upregulation of *nlrp3* is controlled by NF- κ B *e.g.* after TLR activation and is dependent on MyD88, TRIF, Fas-associated protein with a death domain (FADD) and caspase-8.^{50,51} To complement this process, which can take several hours,⁵¹ TLR activation also induces non-transcriptional, or transient, priming independently of protein synthesis, but dependent on proteasome function.⁵² Transient priming has been shown to involve IRAK1 associating with NLRP3-ASC or a pathway dependent on TRIF, IRAK1, FADD, caspase-8 and RIPK1.^{50,53–55} Reactive oxygen species (ROS) production also serves as a signal for non-transcriptional priming.^{56,57}

In addition, the NLRP3 inflammasome is regulated by posttranslational modifications. While it has been well established that NLRP3 deubiquitination promotes priming and activation,^{56,58–63} a recent report shows the opposite,⁶⁴ indicating differential roles depending on cell type or residue modified. NLRP3 phosphorylation has likewise been shown to be both beneficial⁶⁵ and inhibitory^{66,67} for inflammasome activation. Similarly, ASC speck formation can be both enhanced^{68,69} and decreased^{70,71} by ubiquitination, while its phosphorylation⁷² is beneficial for inflammasome activation. Lastly, caspase-1 ubiquitination has been shown to increase inflammasome activation.⁷³

Activation and termination

Over the years, several stimuli have been suggested to induce NLRP3 inflammasome formation: NLRP3 relocation to mitochondrial membranes, mitochondrial damage, cathepsin B in the cytosol and potassium (K⁺) efflux.⁴¹ Attempts to unify these pathways have not yet resulted in a conclusive model, but indicate that a drop in intracellular K⁺ might be the common denominator and that NLRP3 thus reacts to cellular stress.⁴³ A suggested mechanism for this is a conformational change in NLRP3 upon K⁺ efflux that promotes activation.⁷⁴ However, the exact mechanism has not been identified and the million-dollar question of how NLRP3 (or an unknown adaptor protein) senses K⁺ efflux is still considered to be open. Recently, NIMA-related kinase 7 (NEK7) was discovered to be essential for NLRP3 activation downstream of K⁺ efflux.^{75–77} The mechanism is largely unknown but involves NEK7 binding to the LRR and oligomerization domains of NLRP3, independently of NEK7 kinase activity.^{75,78} Figure 1 attempts to illustrate the events surrounding canonical NLRP3 inflammasome activation.

Several mechanisms are in place to regulate inflammasomes. As mentioned above, posttranslational modification of inflammasome components can prevent or decrease activation. Another way is to employ PYD-only proteins (POPs) or CARD-only proteins – both present in humans but not mice – which bind ASC or caspase-1, respectively, to inhibit their recruitment.⁷⁹ IL-10 released after prolonged TLR activation leads to a decrease in NLRP3 expression, which is thought to

prevent aberrant activation upon chronic stimulation.⁸⁰ In addition, prolonged caspase-1 activation leads to self-cleavage and destabilization, thus terminating its own activity.⁸¹ Finally, inflammasomes can be terminated by induction of autophagy and subsequent degradation of inflammasome components.⁷⁰




Figure 1. Canonical activation of the NLRP3 inflammasome. Activation of the NLRP3 inflammasome is a two-step process. The first step, priming, induces transcription of *nlrp3* and *il-1β* after NF-κB activation and also induces posttranslational modifications of NLRP3 and ASC. The second signal, K⁺ efflux, promotes assembly and oligomerization of NEK7, NLRP3 and ASC and recruitment of caspase-1, which cleaves and activates pro-IL-1β and GSDMD. GSDMD-N forms a pore in the membrane, mediating IL-1β release.

Non-canonical and alternative inflammasomes and macrophage hyperactivation

The non-canonical inflammasome pathway is initiated when caspase-11 in mice or caspase-4 and -5 in humans recognize and directly bind cytosolic LPS *via* their CARD domain, thus enabling recognition of bacteria that have escaped vacuoles and phagosomes.^{47,48,82} Non-canonical inflammasome activation can result in both pyroptosis and IL-1β release, although the latter is not mediated directly by caspase-11.⁴⁶ Instead, activated caspase-11 can induce pyroptosis and K⁺ efflux after cleaving GSDMD, leading to NLRP3 inflammasome activation and IL-1β release.^{83,84} Pyroptosis and downstream NLRP3 activation can also be induced by

caspase-11-dependent cleavage and activation of the pore-forming channel Pannexin-1, leading to K⁺ efflux, ATP release and activation of the P2X7 receptor.⁸⁵

Alternative inflammasome activation refers to the involvement of additional proteins other than NLRP3, ASC and caspase-1 or alternative activation stimuli or pathways. For example, stimulating human monocytes with LPS simultaneously primes and activates the NLRP3 inflammasome through a pathway involving TLR4, TRIF, RIPK1, FADD and caspase-8 in addition to NLRP3, ASC and caspase-1.⁴⁹ Caspase-8 affects inflammasome activation in additional ways: by mediating priming together with FADD,⁵⁰ functioning as a scaffold for NLRP3 activation in response to dsRNA⁵⁵ or enhancing caspase-1 activation.⁵⁰

Cells that secrete IL-1 β with retained viability, measured by absence of lactate dehydrogenase (LDH) release, are termed hyperactivated.^{86,87} Hyperactivation can be seen after canonical and non-canonical inflammasome activation in response to oxidized lipids commonly released from dying cells and certain bacterial peptidoglycans and requires GSDMD for IL-1 β release.^{87,88}

Downstream effects of inflammasome activation

Cleavage and maturation of IL-1 β and IL-18 is important in mounting inflammatory responses, *e.g.* recruiting and activating other immune cells in response to pathogens.³¹ Both cytokines lack a sequence targeting them to the conventional secretion pathway and must thus rely on unconventional release,⁸⁹ which has been suggested to occur through several pathways and which will be discussed in later sections. Recently, it was shown that *Salmonella* activates the NLRC4 inflammasome in intestinal epithelial cells, leading to cell expulsion and death and release of eicosanoids and IL-18; protective against infection but also causing diarrhea and fluid loss.²⁹

GSDMD cleavage by caspase-1 or caspase-11 induces an inflammatory form of cell death termed pyroptosis, characterized by the formation of a pyroptosome and cell lysis after plasma membrane permeabilization; commonly measured by release of LDH and staining with the membrane impermeable dye propidium iodide (PI).^{83,88,90} GSDMD cleavage frees an N-terminal domain which oligomerizes on the plasma membrane and forms 10-15nm pores causing cell lysis,^{91,92} but which have also been suggested to mediate IL-1 β release.^{83,90} To avoid harming neighboring cells upon extracellular release, GSDMD-N only binds the inner leaflet of the plasma membrane.⁹¹ Pyroptosis is important in the defense against certain pathogens: intracellular bacteria are left without a niche and released DAMPs can attract and activate other immune cells.²⁴

The downstream effects of inflammasome activation were previously thought to be coupled, leading to a hypothesis that cytokine release is a consequence of membrane

permeabilization and subsequent cell death.^{93,94} However, several studies challenge this idea and propose that pyroptosis and cytokine release are mechanistically separate events. For example, inflammasome activation in neutrophils⁹⁵⁻⁹⁷ as well as the alternative inflammasome described above⁴⁹ result in IL-1 β and IL-18 release without pyroptosis. NLRC4 activation is another example: direct recruitment of caspase-1 to NLRC4 results in pyroptosis, while cytokine release depends on the presence of ASC.^{27,98} In addition, in hyperactivated cells IL-1 β release is mediated through GSDMD pores, which cause membrane permeabilization as evidenced by PI influx.⁸⁸ However, this does not lead to cell lysis, suggesting that membrane permeabilization and lysis are separable events. The mechanisms underlying separation of GSDMD-dependent IL-1 β release and lysis have not been elucidated but are thought to involve a reduction in GSDMD pore size or numbers in hyperactivated cells,⁸⁸ allowing for a newly described membrane repair system to prevent lysis.⁹⁹ However, DiPeso *et al.* suggest that cell lysis and cell death can be separated: cell death is characterized by cell swelling and loss of cell movement and mitochondrial activity after GSDMD-dependent membrane permeabilization, while LDH release indicates cell lysis.¹⁰⁰ Further, this study suggests that glycine can block cell lysis but not cell death, indicating that previous studies might reflect dead cells with retained membrane integrity.¹⁰⁰

Microbial inflammasome modulation strategies

Although there are examples of increased microbial survival and spread after inflammasome activation,¹⁰¹ the numerous strategies used to inhibit inflammasomes indicate their importance in mounting a protective immune response.¹⁰² Some pathogens affect inflammasome components directly: enteropathogenic *Escherichia coli* (*E. coli*) prevent NLRP3 deubiquitination,¹⁰³ human papilloma virus induces pro-IL-1 β degradation,¹⁰⁴ YopM from *Yersinia pestis* inhibits caspase-1¹⁰⁵ and a viral POP inhibits ASC recruitment.¹⁰⁶ *Yersinia* YopE and YopT¹⁰⁷ and *Pseudomonas aeruginosa* ExoS¹⁰⁸ instead interfere indirectly with caspase-1 activity through unexplored pathways involving Rho GTPases and actin polymerization. Another strategy is to avoid activation entirely: *Salmonella enterica* downregulates NLRC4 expression,¹⁰⁹ Gram-negative bacteria can produce modified LPS⁴⁷ and *Staphylococcus aureus* modifies its cell wall to prevent lysosomal degradation and NLRP3 detection.¹¹⁰ Lastly, the recent increase in research on our commensal microflora has revealed anti-inflammatory effects of microbial metabolites¹⁰² and a role for inflammasomes in exacerbating inflammatory bowel disease.³¹ Surely, continued research into the inflammasome field is likely to reveal additional strategies by which pathogenic microbes avoid and exploit this important machinery.

Ubiquitination

At any time, the inside of a cell is teeming with activity. Diverse processes such as DNA repair, endocytosis, protein degradation, signal transduction and cell-cycle regulation can and will happen simultaneously, illustrating the importance of systems to regulate this and prevent mayhem. The small protein ubiquitin – 76 amino acids long and aptly named after its abundance in eukaryotic cells – is one tool used to convey many of these signals; its addition or removal from a target protein makes up a code that is only beginning to be deciphered.

Ubiquitin and its linkage types

Covalent attachment of ubiquitin to its target is mediated by a C-terminal glycine residue which forms a bond with a lysine (K) on the target protein.¹¹¹ One or several target protein residues can be modified, creating monoubiquitination or multi-monoubiquitination, respectively.¹¹¹ Ubiquitin itself contains seven lysines that can be targeted for ubiquitin addition, thus creating chains that can be linked *via* the same lysine, homotypic, or *via* different residues, heterotypic.¹¹² In addition, the N-terminal methionine (Met) can also be ubiquitinated, creating what is called a linear chain.¹¹² Ubiquitination of two residues on one ubiquitin protein generates branched chains and adds further complexity to the code, as does the realization that ubiquitin can be modified at several sites by *e.g.* phosphorylation and acetylation, creating a staggering number of the potential combinations.¹¹³ While the functions of branched and modified chains remain largely unexplored, more is known about K48-, K63- and Met1-linked chains. K48 chains are the most abundant and commonly signal for transport to and degradation by the proteasome; K63 chains can be a signal for autophagic degradation but may also lead to protein transport and kinase activation; Met1 chains have been found to be involved in regulating NF- κ B signaling.^{112,113}

Ubiquitin addition, removal and recognition

Ubiquitination is a three-step process mediated by as many enzymes. First, ubiquitin-activating or E1 enzymes use ATP to form ubiquitin adenylate, thus activating the ubiquitin.¹¹⁴ Activated ubiquitin is then transferred to a ubiquitin-conjugating or E2 enzyme; lastly, interactions with a ubiquitin ligase or E3 enzyme

facilitates transfer of ubiquitin to its target protein.¹¹¹ The job of activating ubiquitin is performed by two known E1 enzymes, while about 40 different E2 and over 600 E3 enzymes work together to confer specificity to the reaction: E2 enzymes determine the chain length and linkage type and E3 enzymes recognize and bind target proteins.¹¹¹ As mentioned above, ubiquitin removal is part of the code. Deubiquitinase enzymes (DUBs) are tasked with removing ubiquitin: cleaving off whole chains, partial chains or single ubiquitin proteins.¹¹⁵ One important role for DUBs is to maintain ubiquitin homeostasis, *e.g.* by removing ubiquitin molecules on proteins destined for degradation.¹¹⁶

Ubiquitin recognition, by DUBs as well as other proteins mediating *e.g.* degradation and transport, is mediated by ubiquitin-binding domains (UBDs), of which there are currently more than 150 described.¹¹⁵ Ubiquitin-binding proteins can create specificity by differentially spacing their UBDs, as the distance between ubiquitin molecules differs depending on chain type: K48 linkage creates a compact chain while K63 results in an elongated version.¹¹⁵

Regulation of inflammasome activation by ubiquitin

As mentioned previously, NLRP3 deubiquitination is thought to mostly be a positive signal for inflammasome activation. The K48- and K63-linked chains are added by several E2 and E3 enzymes; similarly, multiple DUBs have been implicated in their removal,^{59,60,62,63} illustrating the importance of regulating inflammasome activation. Ubiquitin-dependent degradation of pro-IL-1 β is mediated by K48 chains¹¹⁷ while its aberrant activation is mediated by K63-linked chains.^{118,119} The E2 enzyme UBE2L3 was recently shown to add K48-linked chains to pro-IL-1 β ,¹¹⁷ and the DUBs A20¹¹⁹ and POH1¹¹⁸ remove K63-linked chains. Caspase-1 is also activated by K63 ubiquitination,⁷³ but the E2 and E3 ligases involved are currently unknown. ASC ubiquitination can be K63- or Met1-linked; both types can increase speck formation,^{68,69} but K63 chains can also lead to degradation^{70,71} which can be counteracted by the DUB USP50.⁷¹ In summary, ubiquitination plays an important role in regulation cell processes in general and inflammasomes in particular. Each published study increases our appreciation of (and perhaps frustration with) the complex interactions between the ubiquitin code, the proteins interpreting it and the downstream effects it sets in motion.

The P2X7 receptor

Extracellular ATP is an important DAMP sensed by purinergic receptors (P2Rs), divided into P2X and P2Y subfamilies.¹²⁰ While ATP and derivatives of it are the only identified ligands for the ion channel P2X receptors, the G-protein coupled P2Y receptors recognize other nucleotides as well.¹²⁰ P2X receptors are formed as homo- or heterotrimers on the plasma membrane, where ATP binding leads to calcium (Ca^{2+}) and sodium influx and K^+ efflux.¹²¹ The seven P2X proteins identified so far (P2X1-7) are expressed by several immune cells, in many cases with unknown functions.^{121,122} P2X7 is not such a case. Its effects have mostly been studied in macrophages, DCs and microglia, but it is also expressed by and affects maturation and activation of T and B cells.¹²³

Receptor structure, genetics and activation

Like its family members, P2X7 consists of intracellular N- and C-termini, two transmembrane domains and a bulky extracellular domain.¹²⁴ The crystal structure of P2X4 – which shows 41% sequence homology to P2X7 – revealed that a subunit can be likened to a dolphin rising out of the membrane, with three ATP binding sites located in pockets formed by the “head” of one subunit and the “dorsal fin” of another (Figure 2).¹²⁵ Filling the pockets with ATP leads to a conformational change which opens up the receptor complex like a pore.¹²⁵

Unlike its family members, the trimeric P2X7 receptor (P2X7R) has low affinity for ATP and requires millimolar levels for activation, which led many to doubt its physiological relevance as an ATP receptor.^{126,127} However, recent studies have illustrated a dramatic increase in ATP levels at inflammatory sites, indicating that the low sensitivity of P2X7R prevents its aberrant activation.¹²⁸ Another unique aspect of each P2X7 subunit is its elongated cytoplasmic tail which shows sequence homology to the TNF receptor and LBP, although the implications of this has not been investigated further.^{127,129} The cytoplasmic tail is important for receptor oligomerization and plasma membrane localization¹³⁰ and has been suggested to be involved in permeabilization of the plasma membrane, so-called macropore formation, upon P2X7R activation.¹²⁴




Figure 2. Crystal structure of the zebrafish trimeric P2X4 receptor. A trimeric P2X4 receptor without the intracellular C- and N-termini was crystallized with ATP bound. By Hattori *et al*,¹²⁵ Protein Data Bank ID 4DW0.

Genetic variants

There are nine human and three mouse splice variants of P2X7, which all affect activation and receptor formation differently.¹³¹ For both human and mouse, the original, full-length variant is denoted P2X7A.¹³¹ The human C-terminal truncation variant P2X7B forms ion channels but not macropores,¹³² similar to the murine variants P2X7I3B and I3C, which in addition show decreased surface expression.¹³³ In contrast, the murine P2X7K variant, with a different exon 1 as compared to P2X7A, shows increased sensitivity to activation and macropore formation.¹³⁴ P2X7K can also be activated in response to extracellular NAD⁺, while sensing of this ligand instead only lowers the activation threshold for the P2X7A variant.^{135,136} Membrane-bound adenosine di-phosphate (ADP)-ribosyltransferases mediate the transfer of ADP-ribose (ADP-r) from NAD⁺ to P2X7, thus inducing activation or sensitization.^{135,137} T cells have been found to preferentially express P2X7K while macrophages express P2X7A,¹³⁶ explaining why NAD⁺ stimulation leads to P2X7R activation and cell death in T cells but not macrophages.^{135,138}

The discovery of murine P2X7 splice variants led to the realization that the two available and commonly used P2X7-deficient mice, one developed by

GlaxoSmithKline (GSK)¹³⁹ and one by Pfizer,¹⁴⁰ are not full receptor knock-outs. Neither strain expresses the P2X7A variant of the receptor. However, as the GSK strain was generated by disrupting P2X7 exon 1,¹³⁹ and the P2X7K splice variant contains a different first exon, P2X7K escapes deletion in the GSK strain.¹³⁴ Interestingly, the GSK mouse even exhibits enhanced P2X7-responses in some tissues and cells.^{134,141} The Pfizer strain was generated by disrupting the C-terminal exon 13A, but receptor splice variants encoding the alternative exons 13B and 13C escape deletion in this strain.^{133,140} The P2X713B and C receptors do however show decreased function.¹³³ This may offer an explanation to previously reported phenotypical differences between P2X7-deficient mice.¹³¹

To add another level of complexity, single nucleotide polymorphisms (SNPs) also affect P2X7R activity. In humans, gain-of-function mutations may offer protection against some infections but increase the risk of inflammatory disorders, while the reverse is suggested for loss-of-function mutations.¹³¹ Although several SNPs have been reported in mice, only one has been fully characterized.¹³¹ BALB/c mice are more sensitive to ATP compared to C57Bl/6 mice, which is due to a single amino acid difference in the C-terminal tail: in BALB/c mice residue 451 is a proline, in C57Bl/6 it is a leucine.¹⁴² Further studies have shown that the 451L variant, which is found in several mouse strains, confers decreased membrane permeabilization, ion flux and cell death.^{131,142,143}

Macropore formation

In addition to opening up the P2XR ion channels, ATP stimulation of macrophages also causes membrane permeabilization that mediates uptake of molecules up to 900 Da.¹⁴⁴ As these openings allowed the passage of molecules considerably larger than did the P2XR ion channels, they were referred to as macropores.¹²² When P2X7R was discovered it was shown that this receptor could mediate the macropore functions,¹²⁷ and subsequent studies have shown that macropores may also form upon *e.g.* P2X4 and P2X2 activation.¹²⁴ The molecular identity of the P2X7 macropore is a contested subject.¹²⁴ Currently there are two opposing main hypotheses: 1. the macropore really is only a dilated P2X7R pore¹⁴⁵ or 2. additional pore-forming accessory proteins, such as pannexin-1, are recruited and make up the macropore.¹⁴⁶ However, it has been shown that pannexin-1 knock-out cells still show macropore-mediated functions after ATP stimulation,¹⁴⁷ and no other candidate has been proposed. For many years, it has been commonly accepted in the field that macropore formation happens upon prolonged stimulation of P2X7R, a view that has recently been challenged. Indeed, new data indicate that P2X7R activation induces an instant and stable opening of the channel, allowing the passage of large molecules. Contrary to previous beliefs, no dilation of the channel was observed, and it was suggested that previous conclusions about dilation may have

been a result of the interpretation of whole cell-population kinetics rather than kinetics of the pore itself.^{148,149} The C-terminal tail of P2X7 was thought to be required for uptake of large molecules.^{132,142,150,151} Recently however, Karasawa *et al.* showed macropore formation by C-terminal deletion mutants expressed in liposomes.¹⁵² These macropores did not form in cholesterol-containing liposomes,¹⁵² in agreement with a previously reported inhibitory effect of cholesterol on P2X7R-mediated uptake of large molecules.¹⁵³ Complementing the deletion mutants with C-terminal residues allowing palmitoylation¹⁵⁴ counteracted the effect of cholesterol,¹⁵² suggesting that the C-terminal tail may not be required for pore formation *per se* but may have a regulatory effect depending on the lipid composition of the membrane.

Modulation of activation

In addition to NAD⁺, P2X7R can be activated by the AMP LL-37¹⁵⁵ and amyloid- β ,¹⁵⁶ although it is not known whether these substances induce cellular release of ATP¹⁴⁷ which then activates P2X7R, or whether these molecules are actual P2X7R ligands. The antibiotic polymyxin B,¹⁵⁷ LPS,⁸⁵ the antihistamine clemastine,¹⁵⁸ and P2X4R^{159,160} all lower the P2X7R activation threshold by unknown mechanisms. P2X4R can be coimmunoprecipitated with P2X7R and has been shown to interact with the P2X7 C-terminus,¹⁶¹ but reports contradict on whether they interact as homo- or heterotrimers and how this interaction modulates activation.^{162,163}

In addition to lipids, P2X7R activation is regulated by a feedback loop mediated by matrix metalloproteinase 2 (MMP-2). After its P2X7-dependent release, MMP-2 cleaves P2X7 to terminate its activity, thus preventing detrimental effects of sustained receptor activation.¹⁶⁴ P2X7R activation is also negatively modulated by divalent cations such as magnesium (Mg²⁺) and zinc, through chelation of ATP or through poorly understood allosteric inhibition of the receptor itself.¹⁶⁵ Several P2X7R inhibitors have been developed; although structurally different, some of them bind to the same site where they allosterically prevent pore opening.¹⁶⁶ However, due to the polymorphic nature of P2X7, the inhibitory concentrations of these inhibitors vary between both species and individuals, complicating their therapeutic potential.¹³¹

Downstream effects

The most well-known downstream effect of P2X7R activation is formation of the NLRP3 inflammasome, thought to be due to P2X7-dependent K⁺ efflux.^{42,167} However, a recent study demonstrated that the NLRP3-activating K⁺ efflux in fact

is mediated by TWIK2 (two-pore domain weak inwardly rectifying K⁺ channel 2) after cation influx by P2X7R activation.¹⁶⁸ P2X7 has been reported to directly associate with NLRP3,¹⁶⁹ but the importance of this observation on inflammasome activation has not been investigated.

P2X7R activation also leads to nonclassical release of e.g. IL-1 β ,¹⁷⁰ transglutaminase-2,¹⁷¹ cathepsins^{172,173} and MMP-9¹⁷⁴ through its effects on membrane blebbing¹⁷⁵⁻¹⁷⁷ and release of exosomes^{178,179} and secretory lysosomes.¹⁸⁰ In addition, P2X7R can influence cell migration through induction of cytokines and chemokines,¹⁸¹ promote shedding of surface molecules and induce plasma membrane phosphatidylserine exposure, apoptosis and phagocytosis.^{131,182,183} Interestingly, Janks *et al.* recently showed that several of these downstream events depend on P2X7R-dependent activation of a chloride channel,¹⁸⁴ in contrast to a previous study reporting inhibitory effects of chloride on P2X7R activation and NLRP3 activation.¹⁸⁵

These diverse downstream effects reflect the many-faceted role of P2X7R during infection and inflammation. On one hand its activation can exacerbate the inflammatory response during sepsis and in response to some viruses, but on the other its inhibition during certain bacterial, fungal or protozoan infections leads to deleterious effects.¹²³ The same is true for cancer, where both pro- and anti-tumor effects are seen when manipulating P2X7R activation.¹²¹ In contrast, inflammatory diseases such as diabetes and neurodegenerative diseases such as multiple sclerosis are alleviated by P2X7 inhibition, making it an attractive therapeutic target.¹⁸⁶

IL-1 β and its release pathways

IL-1 β and its 10 IL-1 family members possess highly conserved gene structures and are mostly clustered on human chromosome 2, suggesting that the genes arose from duplication of a common ancestor gene.¹⁸⁷ IL-1 β and IL-1 α were the first cytokines to be identified; initially thought to be the same protein, two studies published in 1984 demonstrated different sequences of IL-1 cDNA.^{188,189} Prior to this, Dinarello *et al.* were first in demonstrating the presence of the fever-mediating protein leukocytic pyrogen in humans, a protein that later was shown to be IL-1 β .^{190,191} Since then, a host of studies have attributed additional functions to these inflammatory cytokines and demonstrated their importance in the inflammatory response. In short, IL-1 β induces production of cytokines and chemokines, promotes influx of immune cells by increasing expression of adhesion molecules on endothelial cells and promotes differentiation and expansion of T helper cells.¹⁹² Its downstream effects are mediated by IL-1 receptors (IL-1R), of which there are 10 family members.¹⁹³ IL-1 β binds primarily to IL-1R1, leading to *e.g.* NF- κ B activation through binding of MyD88 to its TIR domain, similar to TLR signaling.¹⁹³

Dysregulated IL-1 β release can wreak havoc in our bodies, illustrated by its association with several autoinflammatory diseases and the success in treating them with anakinra, a recombinant form of the naturally occurring IL-1 receptor antagonist (IL-1Ra), which binds IL-1R1 with high affinity and prevents its activation.¹⁹¹ IL-1Ra is not the only endogenous mechanism that keeps IL-1 β at bay: high levels of IL-10 in the gut keeps macrophages virtually unresponsive to inflammasome activating stimuli,¹⁹⁴ and type I IFNs reduce the amount of pro-IL-1 β via IL-10 production and/or direct inhibition of inflammasome activation.¹⁹⁵ Indeed, for certain patients with multiple sclerosis, IFN- β is an efficient treatment, possibly due to its role in quenching IL-1 β generation.¹⁹¹

Further underlining the importance of keeping the production of IL-1 β under tight control, its release is regulated by several checkpoints. First, baseline IL-1 β expression is low and can be increased after NF- κ B activation, *e.g.* after TLR or IL-1R activation.⁵¹ Second, gene expression results in production of the inactive precursor pro-IL-1 β that requires proteolytic cleavage in order to induce IL-1R signaling.¹⁹⁶ A few years after the cloning of the IL-1 β gene, the enzyme mediating the activating cleavage of the pro-IL-1 β protein was identified to be caspase-1.¹⁹⁷ However, the presence of active IL-1 β in caspase-1^{-/-} mice¹⁹⁸ demonstrated the

presence of additional proteases: host-derived MMP-9, granzyme A and neutrophil proteinase-3^{192,199} and microbe-derived proteases from GAS²⁰⁰ and *C. albicans*²⁰¹ are examples of these. Third, pro-IL-1 β can be targeted for autophagosomal²⁰² or proteosomal degradation, the latter after its ubiquitination.²⁰³ Last, IL-1 β lack the N-terminal leader sequence that allows for release of proteins through the so-called conventional secretion pathway. Proteins that do contain this leader sequence are, upon translation, targeted to the ER lumen from which they are transported to the Golgi, packaged into secretory vesicles and finally released through vesicle fusion with the plasma membrane.²⁰⁴ The only member of the IL-1 family of proteins to be secreted through this pathway is IL-1Ra.¹⁹² Several nonconventional pathways have been suggested to govern IL-1 β release, but externalization of its other family members is a largely unexplored area.

Rabouille *et al.* recently proposed a classification system for unconventional release of leaderless proteins: type I release involves pore-mediated translocation across plasma membranes, type II is mediated by ATP-binding cassette (ABC) transporter proteins in yeast and type III involves membrane-bound organelles.²⁰⁵ These organelles can release their contents either by fusing with the plasma membrane, similarly to the classical release pathway, or by vesicle lysis after their release into the extracellular space.²⁰⁶ Several type I and III pathways have been put forth as alternatives for IL-1 β release. Figure 3 summarizes the proposed unconventional IL-1 β release pathways.

Non-vesicular release (Unconventional type I release)

Non-vesicular, or type I,²⁰⁵ release of IL-1 β suggests that pro-IL-1 β cleavage occurs in the cytosol and that the active cytokine then is transported over the membrane by a transporter protein or diffused through a pore. Recent research has focused on the pore-forming protein GSDMD, but other mechanisms have also been proposed.

GSDMD pores and membrane permeabilization

GSDMD pore characterization revealed that its diameter, 10-15nm,⁹² is large enough to allow passage of IL-1 β with a diameter of 4.5nm²⁰⁷ as well as other cytosolic proteins of similar size, such as IL-18, suggesting a passive process.²⁰⁸ In caspase^{mlt} mice, which lack caspase-1 protease activity, caspase-8 drives GSDMD-independent LDH-, IL-1 α - and IL-1 β release, possibly also reflecting a passive process.²⁰⁹ In contrast, Monteleone *et al.* showed that IL-1 β cleavage is sufficient for release, which can be rapid and GSDMD-dependent or slow and GSDMD-independent.²¹⁰ In this study it was found that pro-IL-1 β cleavage led to exposure of

a specific motif that enabled relocation to and electrostatic interaction with plasma membrane ruffles enriched with phosphatidylinositol 4,5-bisphosphate (PI[4,5]P₂), suggesting an active process regardless of the presence of GSDMD.²¹⁰ On a similar note, lanthanides, a group of chemical elements, prevent Nigericin-mediated PI influx and pyroptosis, but do not affect GSDMD-dependent IL-1 β release, also suggesting an active process.²¹¹

Several other proteins or substrates that cause membrane permeabilization may also mediate IL-1 β release, such as MLKL (necroptosis-associated mixed lineage kinase domain like pseudokinase),²¹² the shellfish toxin maitotoxin²¹³ and the bee venom melittin.²¹⁴ The organic compound punicalagin has been found to block membrane permeabilization/pore formation but not NLRP3 activation, as illustrated by the intracellular accumulation of mature IL-1 β and its subsequent release after punicalagin washout.²¹⁵ The target(s) of punicalagin remain unclear.

Non-vesicular release by other proteins

Prior to the discovery of GSDMD, other pore-forming proteins were suggested to mediate IL-1 β release. An early suggestion was ABC transporters, which facilitate translocation of numerous compounds,^{216–218} but the results of several of these studies were complicated by the realization that the inhibitors used commonly have off-target effects.^{219,220} Similarly, based on studies using inhibitors, pannexin-1 was suggested as a route across the membrane during IL-1 β release.^{221,222} However, subsequent studies using pannexin-1 knock-out mice have shown that this pore-forming protein is dispensable for IL-1 β release.^{147,223} As discussed earlier, P2X7R activation can induce several of the type III pathways discussed above, but P2X7-dependent unconventional release has also been suggested to occur through the P2X7R macropore.²⁰⁴ However, IL-1 β is larger²⁰⁷ than the reported diameter of the macropore,¹⁴⁵ making it unlikely that this is the direct conduit for release.

Fibroblast growth factor 2 (FGF-2) is externalized by an intriguing mechanism. Similar to IL-1 β , PI(4,5)P₂ recruits FGF-2 to the plasma membrane²²⁴ where it oligomerizes and inserts itself, thus creating its own release pore.²²⁵ The mechanism for disassembly and release is not fully elucidated but involves capture by cell surface heparan sulfates.²²⁶ Although IL-1 β itself so far has not been shown to possess pore-forming abilities, it is possible that PI(4,5)P₂ could mediate its release through other proteins.




Figure 3. Unconventional IL-1 β release pathways. IL-1 β release has been suggested to occur through plasma membrane translocation by accessory proteins, after uptake into secretory autophagosomes or through the plasma membrane pores formed by GSDMD after inflammasome activation. In addition, IL-1 β has been suggested to be taken up into exosomes and secretory lysosomes or to be captured in microvesicles.

Vesicular release (Unconventional type III release)

Secretory lysosomes

Secretory lysosomes are lysosomal organelles which have evolved to not only degrade proteins but to also mediate their release by fusing with the plasma membrane.²²⁷ Used by *e.g.* CD8⁺ T cells to deliver granzymes and neutrophils to release chemoattractants, secretory lysosomes depend on Ca²⁺ influx to induce microtubule-dependent migration to the plasma membrane.^{206,228} IL-1 β has been shown to colocalize with the lysosomal markers cathepsin D and LAMP-1, and to show similar secretion kinetics as the former, suggesting a role for a secretory lysosomal release pathway^{179,180} Further studies showed that lysosomal release was blocked by inhibiting the enzymes phospholipase A2 and C, which are activated by Ca²⁺ influx after P2X7R activation by ATP.¹⁸⁰ This early model proposed that pro-IL-1 β processing by caspase-1 may take place in the lysosome²²⁹ and that caspase-

1 and IL-1 β may be protected from lysosomal degradation by interacting with the vesicular membrane and/or by the decrease in lysosomal pH that occurs upon migration of lysosomes to the plasma membrane.²³⁰ However, recent studies demonstrate that caspase-1 activation and pro-IL-1 β processing occurs at the cytosolic ASC speck,²³¹ and that mice with faulty secretory lysosomes still release IL-1 β ,⁹⁴ indicating that early results might have been misinterpreted by *e.g.* contamination of ASC specks during subcellular fractionation and purification of lysosome-containing fractions.²³²

Microvesicles and exosomes

Activation of P2X7R leads to formation of 200nm-1 μ m microvesicles derived from the plasma membrane through a process called blebbing, which has been suggested to trap IL-1 β and mediate its release.^{175,233} Microvesicles can in theory contain any cytosolic protein in the vicinity to the bleb, and their formation and release has been suggested to be dependent on Ca²⁺ influx, calpains and Rho kinases after P2X7R activation.^{176,177} In addition to IL-1 β , microvesicle cytosols have been suggested to contain caspase-1, cathepsin D and high-mobility group protein 1 and to express P2X7R and major histocompatibility complex II (MHC-II) on their surface.^{234,235} As for the proposition of the involvement of secretory lysosomes, the microvesicular release theory has received some critique: there seem to be differential requirements for microvesicle and IL-1 β release after P2X7R activation,¹⁷⁸ and IL-1 β release is independent of Ca²⁺ influx.²³⁶

Smaller than microvesicles, exosomes also contain cytosolic proteins and can be formed by inward budding of endosomes; they are then contained in so-called multivesicular bodies (MVB) in the cytosol and are released when the MVBs fuse with the plasma membrane.²³⁷ Qu *et al.* demonstrated that ATP stimulation leads to P2X7R- and inflammasome-dependent release of exosomes containing IL-1 β and cathepsin D and expressing MHC-II.^{178,238} However, in a later study the same group found that exosome release was independent of caspase-1 activity,²³⁸ raising concerns about its importance for release of mature IL-1 β .

While intact microvesicles and exosomes can deliver their cargo to neighboring cells by fusion with the plasma membrane or endocytosis, they can also bind to surface receptors such as integrins and induce downstream signaling pathways or release their content extracellularly by lysing.²³⁷ It is not clear whether lysis of IL-1 β -containing vesicles occurs actively or not; the presence of P2X7R on their membranes has led to speculation that receptor activation by extracellular ATP could induce vesicle lysis.²³⁹

Secretory autophagy

Like lysosomes, autophagosomes can be repurposed to mediate secretion instead of degradation, a well-known pathway in yeast that only recently has been explored in eukaryotic cells.²⁴⁰ However, whether autophagy inhibits²⁰² or promotes^{241,242} IL-1 β release is debated and has been proposed to depend on type and strength of stimulus. A mechanism of IL-1 β release through secretory autophagy has been described: HSP90-dependent translocation of IL-1 β into the space between the two membranes of an autophagophore, which then fuses directly with the plasma membrane or with a secretory lysosome, forming a so-called amphisome.²⁴² Translocation was suggested to involve IL-1 β unfolding and was dependent on a sequence motif previously implicated in chaperone-mediated autophagy, a pathway used to target specific proteins for degradation.²⁴²

IL-1 β release as a spectrum reflecting signal strength

The release pathways described for IL-1 β have been studied in different cell types from different species, with differing experimental settings and using different methods of analysis, making it difficult to compare the suggested pathways and their physiological relevance. In an attempt to unify results, these pathways have been proposed to reflect the intensity of the activating signal.^{204,206} Initial signals would lead to secretory lysosomal or exosomal release, which is more tightly controlled but less efficient. Packaging of IL-1 β into vesicles would also allow for degradation if the signal is short-lived, *e.g.* through autophagy.²⁰² A persistent threat would call for activation of non-vesicular release pathways, which are effective but potentially threaten cell viability. Related to this, the rate of IL-1 β release can also vary depending on redox state; cells with upregulated antioxidant systems display a reduced rate of IL-1 β release due to buffering of ROS produced upon activation stimuli.²⁴³ Thus, each cell is able to activate several pathways in order to properly regulate the magnitude of the inflammatory response and to prevent the harmful effects of aberrant IL-1 β release.

Group A *Streptococcus*

Streptococcus pyogenes, commonly known as Group A *Streptococcus* (GAS), is a pathogen capable of causing a wide range of diseases in its only host: humans.²⁴⁴ Since up to 20% of school-age children and 25% of adults in contact with school-age children carry GAS asymptomatically, it is considered a facultative rather than obligate pathogen.^{244,245} Relatively mild diseases such as throat infection (pharyngitis, “Strep throat”) or skin infection (impetigo) are found at one end of the spectrum and represent the majority of cases; over 600 million pharyngitis cases are estimated yearly.²⁴⁶ At the other end lies invasive but rare infections such as necrotizing fasciitis (NF, “flesh-eating disease”) and Streptococcal toxic shock syndrome (STSS), complicated to treat and associated with mortality rates of 15-25%.²⁴⁷ About 517 000 deaths are caused by GAS each year, placing it among the top 10 infectious causes of human mortality.²⁴⁶ In a majority of cases, GAS-related death is due to post-infection sequelae such as rheumatic heart disease and acute glomerulonephritis, characterized by autoimmune damage to heart and kidney, respectively.²⁴⁸ A relatively unexplored aspect of GAS infections are their effect on our nervous system: they are linked with rapid onset of obsessive-compulsive and tic disorders in children (pediatric autoimmune neuropsychiatric disorders associated with streptococcal infections)²⁴⁹ and directly activate pain-sensing neurons during NF, leading to a decreased immune response.²⁵⁰ Although the number of GAS infections have decreased globally due to increased living conditions and hygiene, there has been a resurgence in invasive infections since the 1980’s, due to the emergence and subsequent world-wide dissemination of strains with increased fitness and/or virulence.²⁵¹ The incidence of GAS infection follows a seasonal pattern, with rates peaking during winter and spring,²⁵² and a recent study suggests an association between invasive GAS infections and influenza A infection.²⁵³

GAS is Gram-positive and characterized from other species in *e.g.* a throat culture by complete lysis of red blood cells (β -hemolysis) on blood agar plates and presence of a group A carbohydrate (GAC) on its surface, a classification system developed by Rebecca Lancefield based on differing immunological responses to streptococcal cell wall carbohydrates.^{248,254} As it is the only member of the *Streptococcus* genus expressing a GAC, the term GAS has become synonymous with the species *Streptococcus pyogenes*. Further serotyping is based on genetic differences in the

surface-expressed M protein, a major virulence factor that currently exists in over 200 variants.²⁵⁵ Other cell-surface proteins such as T and R proteins can also be used to distinguish strains from each other.²⁴⁸ Once identified, a GAS infection is treated with antibiotics; although increased resistance to certain antibiotic types is being reported, GAS is amazingly still sensitive to penicillin.²⁵⁵ Notably, disease progression in invasive infections can be too overwhelming for antibiotic treatment to be effective: 8-23% of patients with invasive disease die within 7 days of infection even when antibiotics are administered, and the number is even higher if STSS is developed.²⁵⁵ There is currently no vaccine against GAS, despite many years of research and several clinical trials. Diversity among strains and their geographic distribution is one complicating factor, incomplete understanding of adaptive immune responses and their role in inducing post-infection sequelae another.²⁵⁶ As an example, molecular mimicry between the M protein and myosin allows antibodies to the former to cross-react with the latter, which may contribute to disease progression in rheumatic heart disease.²⁵⁷

Pathogenesis

GAS expresses an impressive arsenal of virulence factors: proteins, enzymes and carbohydrates that are used during an infection to ensure optimal adherence, survival and spread. Recent research has revealed that virulence factors may play different roles depending on GAS location, leading to a realization that the interplay between virulence factors and the cues dictating their regulation may be more complicated than previously thought.

Adherence and colonization

The most crucial step in GAS pathogenesis is adhesion to surfaces such as our skin and nasopharyngeal mucosa, which constantly work to remove foreign objects by exfoliation or mucus and saliva flow.²⁴⁸ Adhesion is thought to occur in two steps: weak, hydrophobic interactions between bacterial lipoteichoic acid and host cell surfaces,²⁵⁸ followed by specific adhesion mediated by bacterial surface molecules, such as the M protein and the hyaluronic acid capsule which may bind CD46 and CD44 on keratinocytes, respectively.^{259,260} The extracellular matrix is also an important adhesion site illustrated by the presence of several fibronectin-binding proteins.²⁵⁸ GAS competes with the local microflora but can also cooperate with it to form mixed-species biofilms, adherent bacterial aggregates associated with chronicity in other types of bacterial infections.²⁶¹ GAS biofilms and their role in pathogenesis are poorly studied, but presence of biofilm in biopsies from NF

patients was associated with increased bacterial load and tissue destruction, indicating a role in disease severity.²⁶²

While GAS as a species is versatile in its ability to infect different tissues, specific strains often show a preference to infect either skin or throat. Genetic studies have shown that the sequences of the M protein-encoding gene *emm* and other *emm*-like genes can be grouped depending on tissue tropism: throat specialists, skin specialists and generalists without preference.²⁴⁴

Invasion and evasion

GAS is typically considered an extracellular pathogen, but it has become clear that it can invade and reside in epithelial cells²⁶³ by *e.g.* escaping autophagy.²⁶⁴ Detection of intracellular GAS in removed tonsils²⁶⁵ and skin infection biopsies²⁶⁶ suggests that recurrent and persistent infections are due to this reservoir, which is not targeted by antibiotics. Intracellular survival and replication²⁶⁷ can also occur in macrophages, by prevention of phagolysosomal fusion²⁶⁸ or acidification.²⁶⁹

While opsonophagocytosis might not be an immediate death threat, it is still a crucial part of the host defense system and GAS has evolved several ways of inhibiting it. M protein²⁷⁰ and the hyaluronic capsule²⁷¹ inhibit or prevent opsonization by complement proteins, IgG-degrading enzymes prevent antibody opsonization,²⁷² the cell wall proteinase SpyCEP inhibits neutrophil recruitment by cleaving chemoattractants such as IL-8^{273,274} and cytolysins such as streptolysin O (SLO) and S (SLS) induce cell damage and death by forming large pores.²⁷⁵⁻²⁷⁷ The additional immune-modulatory effects of SLO are discussed later on.

The systemic and detrimental inflammatory response during STSS is thought to be due to streptococcal pyrogenic exotoxins (Spe). These superantigens (SAg) act as a bridge between the β -chain of TCRs and MHC-II molecules, causing antigen-independent activation and proliferation of a large number of T cells.²⁴⁸ Kasper *et al.* suggest that the biological role of SAg is to establish nasopharyngeal infection, as the presence of SAg led to a dramatic increase in colonization of mice expressing human MHC-II, arguing that nonspecific T cell activation might be an unfortunate side effect of SAg expression after colonization.²⁷⁸ In accordance with this, immunization against SpeA protects against nasopharyngeal infection in a T cell-dependent manner.²⁷⁹

The GAS genome contains multiple regulatory systems that control the expression of virulence factors and thus the transition from colonization to infection, depending on *e.g.* bacterial density and environmental cues such as extracellular Mg²⁺ and LL-37 for the control of virulence regulatory system (CovRS).^{280,281} This two-component system regulates 10-15% of the genome, for example it negatively regulates the expression of capsule and SLO and positively regulates SpeB, a

secreted cysteine protease which cleaves both host and microbial proteins.²⁸² Invasive and hypervirulent GAS isolates commonly exhibit inactivating mutations in CovRS, allowing for increased expression of virulence factors and resistance to defense mechanisms.^{283,284} These mutations are almost always *de novo* mutations that occur in the host, as strains with inactive CovRS are impaired in establishing infection due to hyperencapsulation and thus also impaired in spreading to other hosts.^{285,286} Invasive infections are therefore thought of as evolutionary dead-ends, arising due to colonization of and local adaptation to sites that normally are sterile.²⁸⁷

M1T1 GAS

A widespread M1T1 strain that has been circulating since the 1980's has been under much scrutiny. An epidemiologic study of invasive GAS cases in USA showed that while several M types cause invasive infections, M1 strains dominated,²⁸⁸ and analysis of invasive M1 isolates from Canadian patients showed that they were all M1T1 clones.²⁸⁹ However, an Australian study suggests that M1T1 strains are not inherently more invasive than other strains and that the apparent prevalence in invasive disease instead is due to the overall abundance of this strain in the population.²⁹⁰ This is in line with what was discussed above regarding hypervirulence; however, as the majority of studies on M1T1 strains have focused on why they cause invasive infections, their (eventual) fitness advantage remains unexplored.

The evolution of epidemic M1T1 GAS

By sequencing the entire genome of 3615 M1T1 isolates found worldwide, and by comparing their sequences with a pre-epidemic reference strain, Nasser *et al.* repeated and solidified the findings of a previous study showing that emergence of the epidemic strain was precluded by three gene transfer events.^{291,292} The first two were *via* bacteriophages: first the extracellular virulence factor streptococcal DNase D2 was acquired, followed by SpeA1, a SAg.²⁹¹ Replacing a single nucleotide in SpeA1 gave rise to SpeA2, which *in vitro* shows increased binding to MHC-II.²⁹³ Finally, a M12 strain is thought to be the donor during a horizontal gene transfer (HGT) event leading to the acquisition of a 36 kb region which increases production of the virulence factors SLO and β -nicotinamide adenine dinucleotide (β -NAD⁺, NAD⁺)-glycohydrolase (NADase).^{291,292} This region differs from pre-epidemic strains by a few SNPs, two of which are located in the promoter region for the *nga* gene encoding NADase; these two plus a SNP in the coding region of *nga* are sufficient and necessary for increased expression of SLO and NADase.²⁹⁴ A similar HGT event is thought to precede the recent increase in M89 prevalence, as some of

these emerging M89 strains express the same promotor variant as the epidemic MIT1 strains.^{294,295} M1 and M89 strains expressing increased levels of SLO and NADase have been shown to resist phagocytic killing and cause increased disease in several mouse models,^{294,296–298} indicating that the two virulence factors are important in both bacterial survival and disease progression.

NADase expression in M1 strains compared to pre-epidemic strains has been noted in clinical isolates in different parts of the world.^{299,300} In a portion of these isolates, mutations conferring little or no NADase activity have become established, indicating that NADase lacking enzymatic activity is under positive selection.^{300–302} The role of inactive NADase in bacterial fitness or disease progression has not been fully elucidated, although Riddle *et al.* suggest that NADase-inactive strains show tissue tropism while NADase-active strains are generalists.³⁰¹

SLO, β -NAD⁺-glycohydrolase and cytolysin-mediated translocation

The relationship between SLO and NADase begins with their transcription. The *nga* gene is located upstream of *slo* and the two genes are transcribed as a single operon,^{303–305} although *slo* has also been reported to have its own promoter.³⁰⁶ The next part of their close association involves stability. NADase binds SLO and stabilizes both toxins upon secretion, preventing their breakdown and enhancing SLO-mediated cytotoxicity.^{307,308} As this is independent of NADase activity, it is thought to be one reason for the positive selection of NADase.³⁰⁷ The final part of cooperation and co-dependence between the two virulence factors is in regard to their effects on host cells.

SLO belongs to the cholesterol-dependent cytolysin (CDC) family, which includes pore-forming toxins expressed by several types of bacteria. CDC pores can be up to 30 nm in diameter and are made up of about 50 monomers which oligomerize into a pre-pore complex before insertion, a process that requires membrane cholesterol.³⁰⁹ However, SLO membrane binding and pre-pore formation can occur in the absence of cholesterol; a carbohydrate-binding motif allows SLO to bind galactose-containing residues on red blood cells^{310,311} and epithelial cells.³¹² Additionally, NADase mediates SLO binding to epithelial cells in a manner independent of both cholesterol and carbohydrates.^{312,313} Pore formation induces apoptosis in keratinocytes³¹⁴ and macrophages,³¹⁵ but can also cause an inflammatory form of cell death termed oncosis, characterized by loss of membrane integrity, organelle swelling and leakage of intracellular contents.³¹⁶ The reason why infected mice often fare better in its absence^{315–317} are not limited to reduced cell death: SLO can reduce cytokine release,^{318,319} protect from phagocytosis³²⁰ and impair neutrophil degradation.³²¹

In a seminal paper from 2001, Madden *et al.* described a new function for SLO: translocation of NADase into the host cell cytosol.³⁰⁴ The process was termed cytolysin-mediated translocation (CMT) and was the first injection system described for Gram-positive bacteria. CMT occurs only when GAS adheres to host cells and requires secretion of SLO and NADase from the same bacterium.³⁰⁴ NADase then binds an unknown receptor *via* its N-terminal domain,³¹² which requires the presence of SLO.³¹² NADase binding promotes translocation, as it mediates SLO binding even in the absence of cholesterol or carbohydrates.³¹³ Other CDCs can become CMT competent by fusing them to an N-terminal domain unique to SLO and by adding residues from the SLO membrane-binding domain; both are required but not sufficient by themselves.^{313,322} Similarly, fusing the N-terminal domain and last C-terminal residues from NADase to other proteins enables their translocation.³²³ Translocation was first thought to occur through the SLO pore, but as subsequent studies showed that pore formation in fact is dispensable^{322,324} it is currently unknown how NADase travels across the membrane.

NADase was discovered in 1957, when it was shown that fractions of SLO purified from bacterial supernatant hydrolyzed the important coenzyme NAD⁺ into nicotinamide (NAM) and ADP-ribose (ADP-r),³²⁵ a process that can lead to energy store depletion and cell death for both pro- and eukaryotic cells.³⁰⁸ This toxicity thwarted early attempts to express NADase in, and purify it from, *E. coli* until immunity factor for *Streptococcus pyogenes* NADase (IFS) was discovered; a small cytoplasmic protein which blocks the active site pocket in NADase,^{305,326,327} thus protecting the bacterium from the effects of its toxin. Transcribed in the same operon as *nga* and *slo*, *ifs* is often truncated and nonfunctional in NADase-inactive mutants.³⁰⁵ As the ADP-r-cyclase and -transferase abilities previously ascribed to NADase purified from bacterial supernatant^{299,328} were absent when studying a cloned version, NADase is now thought to be a strict glycohydrolase.³²⁹ Both NAD⁺ cleavage products affect eukaryotic cells: NAM inhibits NF- κ B activation and thus prevents the production of pro-inflammatory cytokines^{330,331} and ADP-r induces Ca²⁺ signaling, which has multiple downstream effects.^{332,333}

Enzymatically active NADase enhances SLO-mediated toxicity by NAD⁺ and ATP depletion,^{304,308,334,335} while inactive NADase induces necrosis dependent on activation of c-Jun N-terminal kinase (JNK), a MAPK.³³⁶ However, intracellular delivery of enzymatically NADase alone also induces toxicity.³³⁷ NADase affects cytokine release by modulating the activity of the stress-related enzyme PARP-1,³³⁸ activated by SLO pore formation: active NADase induces release of the TLR4-binding DAMP high mobility group box 1³³⁹ (HMGB1), inactive NADase induces release of IL-8 and TNF- α .³³⁵ Together, SLO and NADase promote intracellular survival by inhibiting phagolysosomal acidification²⁶⁹ and autophagic killing,³⁴⁰ but NADase has also been shown to promote intracellular survival independently of

SLO.³³⁷ Lastly, SLO and NADase are involved in regulating IL-1 β release after NLRP3 inflammasome activation, discussed below and in Paper I and II.

The host response to GAS

Whether it is adhering to barriers such as epithelial cells or spreading into deeper tissue, GAS does not go unnoticed by its host. The continued arms race between host and pathogen has led to development of several strategies to ensure proper removal of the pathogen, among them opsonization by complement proteins and subsequent phagocytosis. Additional host responses are discussed below.

Cytokines and PRRs

Despite not being competent for phagocytosis, non-immune cells such as keratinocytes are not defenseless against invading GAS: K⁺ efflux induced by pore-forming toxins triggers p38 MAPK activation and IL-8 release from epithelial cells,^{341,342} recruiting *e.g.* neutrophils which limit GAS spread through production of AMPs and ROS.³⁴³ Professional phagocytes such as macrophages and DCs are also important in limiting GAS dissemination, evidenced by increased bacterial numbers and decreased survival when either cell type is depleted.^{344,345} Disease control is mediated by production of IL-6, TNF- α and IL-12 through induction of Th17 cells,³⁴⁶ recruitment of macrophages³⁴⁷ and protection during skin infection,³⁴⁸ respectively and in addition to activation of innate cells. Lastly, IFN- β production is crucial for preventing hyperinflammation by controlling neutrophil influx.^{349,350}

MyD88 plays a central role in launching defense mechanisms in response to GAS. Its deletion leads to a decrease in cytokine levels^{349,351–353} and in upregulation of DC activation markers³⁵³; although phagocytosis is unaffected, MyD88^{-/-} mice exhibit increased susceptibility, pathology and bacterial numbers.³⁵² MyD88 deficiency in humans leads to predisposition to invasive pyogenic infection at early age.³⁵⁴ Despite this, the upstream receptors and their ligands have not been fully elucidated. TLR4 is one proposed candidate involved, as SLO has been suggested to activate TLR4³⁵⁵, TLR4^{-/-} mice display decreased mucosal antibody responses to GAS infection³⁵⁶ and human *tlr4* polymorphisms are linked with predisposition to GAS infections.³⁵⁷ On the other hand, TLR2/4/9 single or triple knockout cells from mice only exhibit a slight reduction in cytokine production *in vitro*.^{351,353} Similarly, TLR2^{-/-} mice have been reported to be both susceptible to³⁵⁸ or unaffected by³⁴⁷ *in vivo* GAS infection compared to wild-type (wt) mice. GAS nucleic acids have been suggested as TLR13^{358,359} or STING³⁴⁹ ligands, but a recent study was unable to

detect cytosolic GAS DNA during infection and instead suggest that the M protein induces STING-dependent IFN- β induction via an unidentified receptor.³⁶⁰

Adaptive immune responses

The adaptive response to GAS is less characterized than the innate. On one hand, mice deficient in T- and B-cells exhibit similar clearance and survival rates as wt mice,^{361,362} but on the other anti-M protein antibodies provide protection against infection in mice and humans,³⁶³ suggesting that humoral adaptive immunity is important in preventing infections. Additionally, as children have higher infection rates than adults³⁶⁴ and MyD88-deficient individuals are only susceptible to infection when young, adaptive immunity developed during life seems to be able to complement innate-mediated protection.³⁵⁴ In a study comparing immune responses children and adults, it was found that adults displayed higher levels of IFN- γ and IgG3 (the IgG subclass that most efficiently mediates complement activation and opsonophagocytosis) during GAS infection.³⁶⁵ Thus, it is possible that differential immune responses may partly explain why adults and children present with very different diseases after GAS infection.³⁵⁴

The T cell response against GAS is dominated by Th1 and Th17 cells, with low numbers of Th2 cells, in both mice and humans.^{365,366} Mouse studies have also revealed that the T helper cell response against GAS is affected by route of infection. Intravenous or subcutaneous infection results in formation of antigen-specific Th1 cells, while Th17 cells dominate after intranasal infection.³⁴⁶ It has been shown that Th17 numbers are reduced in TLR2^{-/-} mice in response to GAS³⁵⁶ and that Th17 cell expansion upon GAS infection is dependent on production of IL-6 and TGF- β .^{346,367,368} In addition, transfer of CD4⁺ T cells from mice immunized with GAS into naïve hosts can promote bacterial clearance from the nasal cavity, but only if the transferred T cells can produce IL-17.³⁶⁷ This indicates a protective role for Th17 responses during mucosal GAS infection, but antigen-specific Th17 cells have also been shown to promote detrimental central nervous system infiltration and blood-brain barrier leakage during GAS infection.³⁶⁹

Other host factors influencing infection outcome

In mice, sex affects susceptibility to GAS infection: males generally exhibit exacerbated symptoms and increased mortality compared to females.^{362,370} Another host factor that can influence the outcome of infection is MHC-II; some haplotypes are associated with increased susceptibility to sepsis while others confer protection, in humans³⁷¹⁻³⁷⁴ and mice³⁶¹ alike. Those with protective haplotypes are better at controlling bacterial and immune cell proliferation and cytokine production, preventing the extensive tissue destruction and bacteremia seen in susceptible

haplotypes.^{361,362,371–375} Susceptibility in mice has been mapped to four additional loci, and analysis of differentially expressed genes in these loci identified IL-1 β as a key regulator of disease outcome.³⁷⁶ The implications of this are discussed below.

The role of IL-1 β during GAS infection

After identifying IL-1 β as a key susceptibility regulator in mice infected with GAS, Chella Krishnan *et al.* compared *il-1 β* expression in several different mouse strains and found that decreased survival was associated with high *il-1 β* expression.³⁷⁶ A link between high IL-1 β levels and detrimental effects during GAS infection has been reported previously: type I IFN receptor^{-/-} mice produce increased levels of IL-1 β , which leads to increased neutrophil influx and cytokine levels and results in systemic hyperinflammation and decreased survival.³⁵⁰ A similar phenotype is seen in mice deficient in NF- κ B signaling; here hyperinflammation is rescued by deletion of IL-1R.³⁷⁷ The presence of a neutralizing anti-IL-1 β antibody also reduces wound size in a GAS skin infection model.³⁷⁸ However, IL-1 β is not only detrimental to an infected host: IL-1R^{-/-} and IL-1 β ^{-/-} mice or mice infected in the presence of anakinra also show decreased survival and increased lesion size and bacterial numbers,^{200,350,377,379} illustrating that a balanced IL-1 β response is beneficial for disease outcome. This is also reflected in humans, as patients receiving anakinra have a 300-fold higher risk of contracting invasive GAS infections, while infections by other pathogens are unchanged.²⁰⁰ Despite the clear implications of removing IL-1 β or its receptor during GAS infection, results are less clear when it comes to the role of specific inflammasome components *in vivo*. Harder *et al.* found no difference in susceptibility between NLRP3^{-/-} mice and wt during intraperitoneal infection, despite lower serum levels of IL-1 β .⁴⁴ In contrast, caspase-1/11^{-/-} mice were more susceptible than wt mice in a subcutaneous infection model,³⁵⁰ suggesting that there might be differences in the response depending on infection route.

Several GAS proteins have been described to induce IL-1 β release. Most recent is SLS, suggested to play a partial role in IL-1 β release during GAS infection,³⁷⁸ but the mechanism and inflammasome components involved were not expanded on. Another GAS protein shown to activate NLRP3 is the ADP-ribosylating enzyme SpyA, leading to decreased bacterial survival in host cells.³⁷⁹ In addition, recombinant M1 protein induces NLRP3-dependent IL-1 β release after endocytosis, and infecting macrophages with GAS strains lacking M1 results in a partial reduction in IL-1 β levels compared to wt GAS.³⁸⁰ Last in this list but the first GAS inflammasome activator to be described, SLO-dependent activation of NLRP3⁴⁴ is dependent on K⁺ efflux³⁸¹ and requires SLO pore formation.³⁸² Increased IL-1 β levels in response to a partially active SLO mutant are thought to be due to decreased cytotoxicity, allowing for plasma membrane repair and a sustained response.³⁸² We have recently shown that GAS can regulate IL-1 β release through NADase-

dependent inhibition³⁸³ (Paper I) of a release pathway involving P2X7 and membrane permeabilization (Paper II).³⁸⁴ In contrast to previously reported functions of NADase, inhibition of P2X7-dependent IL-1 β release is mediated by the extracellularly located fraction of this toxin (Paper I).³⁸³ In addition to its inflammasome-activating properties, SLO also seems to be able to regulate IL-1 β levels, by inducing ubiquitination and degradation of pro-IL-1 β which could be rescued by the broad autophagy inhibitor 3-methyladenine (3-MA) (Paper III).³⁸¹ This suggests that GAS has evolved several mechanisms of regulating IL-1 β release (Paper I-III).

Present investigation

Original papers and manuscripts

Paper I: Inhibition of Inflammasome-Dependent Interleukin 1 β Production by *Streptococcal* NAD⁺-Glycohydrolase: Evidence for Extracellular Activity

Dóra Hancz, Elsa Westerlund, Benedicte Bastiat-Sempe, Onkar Sharma, Christine Valfridsson, Lena Meyer, John F. Love, Maghnus O'Seaghda, Michael R. Wessels, Jenny J. Persson.

mBio. 2017 Jul 18;8(4). pii: e00756-17. doi: 10.1128/mBio.00756-17

Paper II: The Secreted Virulence Factor NADase of Group A *Streptococcus* Inhibits P2X7 receptor-mediated Release of IL-1 β

Elsa Westerlund, Christine Valfridsson, Daisy X Yi, Jenny J. Persson.

Front. Immunol. 18 June 2019. doi: 10.3389/fimmu.2019.01385

Paper III: Streptolysin O Induces Ubiquitination and Degradation of pro-IL-1 β

Dóra Hancz, Elsa Westerlund, Christine Valfridsson, Getachew Melkamu Aemero, Benedicte Bastiat-Sempe, Pontus Orning, Egil Lien, Michael R. Wessels, Jenny J. Persson.

J Innate Immun. 2019 Mar 19:1-12. doi: 10.1159/000496403

Paper IV: *Streptococcal* β -NAD⁺-glycohydrolase binds to and induces cytokine release from macrophages in the absence of Streptolysin O

Elsa Westerlund, Lena Meyer, Christine Valfridsson, Dóra Hancz, Onkar Sharma, Michael R. Wessels and Jenny J. Persson.

Manuscript

Summary and discussion of papers

Paper I: Inhibition of Inflammasome-Dependent Interleukin 1 β Production by Streptococcal NAD⁺-Glycohydrolase: Evidence for Extracellular Activity

Background and aim

The two virulence factors SLO and NADase are genetically and functionally linked: SLO protein stability requires co-expression of NADase³⁰⁷ and SLO can mediate translocation of NADase into host cell cytosols.³⁰⁴ Their combined effects include cell death and phagolysosomal escape,²⁶⁹ and strains expressing them show increased disease symptoms and are overrepresented in invasive infections.^{289,299}

The inflammatory cytokine IL-1 β is important during GAS infection – without it, bacterial numbers increase and disease symptoms worsen.^{200,350,379} The NLRP3 inflammasome can be activated by at least three different GAS proteins,^{44,379,380} one of which is SLO and the pores it forms.^{44,382} However, the role of NADase during SLO-dependent inflammasome activation inflammasome has not been studied previously. We set out to investigate this using a macrophage infection model and a set of isogenic bacterial mutants.

Key findings

- NADase inhibits IL-1 β release after SLO-dependent NLRP3 activation.
- NLRP3 activation and inhibition of IL-1 β release are executed by extracellularly located bacteria.
- Differences in IL-1 β release are not due to differential induction of cell death or priming.
- Caspase-1 activation levels are similar in absence or presence of NADase, suggesting an inhibitory effect of NADase downstream of inflammasome activation.
- Inhibition of IL-1 β release is performed by non-translocated NADase, revealing a novel role and functional niche for the enzyme.

Discussion

After the discovery that cell adherent GAS may perform CMT, SLO-mediated translocation of NADase into the cell cytosol,³⁰⁴ much effort has been invested in elucidating its mechanism and understanding the effects of intracellularly located

NADase. During an infection however, it is likely the case that NADase is both translocated and present extracellularly. By taking advantage of the fact that translocation occurs only upon secretion from the same bacterium³⁰⁴ and by using a strain not competent for translocation,²⁶⁹ our study is the first to show that extracellular NADase has a specific function during macrophage infection: inhibition of IL-1 β release. We also found that NADase cannot affect IL-1 β release induced by SLO when only pure SLO and NADase proteins are used (unpublished data), implying that an infection context and possibly other host-pathogen interactions are required for this event to occur. It remains to be investigated whether this immune regulatory property is common to all NAD⁺-hydrolase enzymes expressed by other bacterial species, such as the newly described NAD⁺ and NADP⁺ glycohydrolase expressed by *Mycobacterium tuberculosis*.^{385,386}

Patients receiving the IL-1R antagonist anakinra have a significantly increased risk of specifically contracting invasive GAS diseases, illustrating the importance of IL-1 signaling for infection control. In contrast, in a study of susceptibility to GAS among mouse strains, high levels of IL-1 β correlated with decreased survival,³⁷⁶ and yet another indicated that uncontrolled IL-1 β levels cause lethal hyperinflammation and tissue damage.³⁵⁰ This indicates that IL-1 β responses need to be finely balanced for optimal host survival and bacterial clearance. Our findings suggest that GAS have evolved a way to limit IL-1 β levels during infection, possibly to skew the immune response in its favor. However, the contribution of NADase-dependent inhibition of IL-1 β during GAS infection still requires further investigation before we can fully understand its effects on disease progression.

While several pathogens have the ability to interfere with the generation of IL-1 β , *e.g.* by preventing inflammasome formation and activity,^{103,105} our study indicates that NADase inhibits IL-1 β release *per se*. Several non-classical IL-1 β release pathways have been described, among them release *via* exosomes, secretory lysosomes or through GSDMD pores.²⁰⁶ It is possible that NADase targets one or several of these pathways to prevent IL-1 β release. In summary, our results describe a previously unknown role for NADase and functionality of the toxin in a novel location. Inhibition of IL-1 β release may mediate regulation of the immune response, potentially leading to increased bacterial spread in the host.

Paper II: The Secreted Virulence Factor NADase of Group A *Streptococcus* Inhibits P2X7 receptor-mediated Release of IL-1 β

Background and aim

The inflammatory cytokine IL-1 β lacks a leader sequence directing it to the classical ER-Golgi secretion pathway and thus has to rely on unconventional protein release pathways.²⁰⁶ Several pathways have been suggested, including packaging into

secretory lysosomes or exosomes, active membrane transporters or GSDMD pores.²⁰⁶ Activation of the purinergic receptor P2X7 leads to Ca²⁺ and K⁺ flux, which has been implicated both in NLRP3 inflammasome activation and unconventional protein release.¹⁷⁰ Innate immune cells such as macrophages activate the NLRP3 inflammasome and release IL-1 β in response to K⁺ efflux induced by GAS and its pore-forming toxin SLO.⁴⁴ We have previously shown that extracellularly located NADase can counteract this response by inhibiting IL-1 β release through an unknown mechanism (Paper I).³⁸³ The focus of Paper II was to investigate this pathway further.

Key findings

- In absence of functional NADase, GAS induces a P2X7-dependent IL-1 β release pathway.
- P2X7-dependent IL-1 β release occurs independently of extracellular ATP and is not affected by NAD⁺ cleavage products.
- Protein degradation or vesicular release does not differ in absence or presence of functional NADase.
- P2X7-dependent IL-1 β release is dependent on permeabilization of the plasma membrane.
- GSDMD is not involved in P2X7-dependent or -independent IL-1 β release in response to GAS.

Discussion

As the presence of NADase or PEGs only partially inhibits IL-1 β release, our results suggest that this cytokine is externalized through two pathways in response to GAS, one of which involved the P2X7 receptor and membrane permeabilization. This partial inhibition could represent a way for the host to create redundancy – by releasing IL-1 β through several non-classical pathways, immune system activation in response to GAS is ensured. Release through several pathways has also been suggested to reflect signal strength; in response to a weak signal, one pathway is activated, while another might be switched on as the signal intensity increases.²⁰⁴ In addition, maintaining IL-1 β at appropriate levels could also be beneficial for the bacterium. Invasive infections caused by complete lack of IL-1 signaling are detrimental for the host, but GAS which in these cases are located in normally sterile sites may also be less likely to spread to new individuals.²⁸⁷ Host mortality likely also limits the spread of GAS in a population suggesting that lethal GAS infections may be an evolutionary dead end for the bacterium.²⁸⁷

The P2X7 receptor is well-known for its importance in ATP-mediated NLRP3 activation, but has also been implicated in the regulation of unconventional protein

release pathways.¹⁷⁰ Our data suggest P2X7-mediated IL-1 β release is not vesicular or autophagic, but indicates a requirement for plasma membrane permeabilization. In light of the recent discovery that GSDMD pores mediate IL-1 β release in response to several inflammasome-activating stimuli,²⁰⁶ we were thus surprised to find that GAS-induced IL-1 β release is completely independent on GSDMD. Activation of the P2X7 receptor has been reported to lead to opening of a large membrane pore, the so-called P2X7 macropore. This structure has been suggested to mediate unconventional release of proteins,³⁸⁷ but whether the macropore is formed by the trimeric P2X7 receptor itself or by accessory proteins is debated.¹²⁴ However, as the estimated size of the pore formed by P2X7 is smaller than IL-1 β ,^{145,207} direct release through P2X7 itself is questionable and accessory proteins are likely involved.¹²⁴ Our data supports this hypothesis, although we were not able to specifically identify additional proteins involved. In addition, our data does not allow us to predict the sequence of events surrounding P2X7-dependent IL-1 β release, *i.e.* whether P2X7 and eventual accessory proteins are involved in an upstream regulatory process or if they are involved in the actual release. The C-terminal tail of P2X7 harbors predicted binding motifs for multiple proteins and lipids¹²⁹ and is important for membrane localization¹³⁰ and permeabilization.³⁸⁸ It is thus attractive to speculate that P2X7-dependent functions can be dictated or modified by different binding partners to these domains, including the release of IL-1 β in response to GAS infection.

In summary, our results indicate that NADase acts to inhibit an IL-1 β release pathway dependent on the P2X7 receptor and membrane permeabilization. While the exact mechanism is still unknown, our study sheds light on IL-1 β release in response to a setting where several stimuli are present and pinpoints P2X7 as a future research interest in relation to GAS infections.

Paper III: Streptolysin O Induces Ubiquitination and Degradation of pro-IL-1 β

Background and aim

Addition or removal of the small protein ubiquitin can modify the activation and function of a target protein as well as mark it for degradation or secretion.¹¹² The presence of one or several ubiquitin moieties, and whether the chains they form are linear or branched, determines which of the diverse fates a target protein meets.¹¹² An example of how ubiquitination affects cellular processes relates to NLRP3 inflammasome activation and subsequent IL-1 β release. Caspase-1 ubiquitination enhances its activity⁷³ while NLRP3 activity instead requires deubiquitination, preventing its proteasomal degradation.^{60,62} ASC speck formation can be both enhanced⁶⁸ and inhibited⁷¹ by ubiquitination, as can pro-IL-1 β levels; either leading

to aberrant activation^{118,119} or degradation.^{117,203} During our investigations into NADase-dependent inhibition of IL-1 β ,^{383,384} we found that pro-IL-1 β was ubiquitinated in macrophages in response to GAS infection and that this was dependent on SLO, thus prompting us to further investigate this phenomenon.

Key findings

- SLO-expressing GAS induce ubiquitination of pro-IL-1 β .
- SLO pore formation is required but not sufficient for ubiquitination of pro-IL-1 β .
- SLO-dependent ubiquitination occurs independently of inflammasome activation or inflammasome proteins.
- Pro-IL-1 β ubiquitination is of mixed linkage specificity.
- Pro-IL-1 β is degraded upon infection with SLO-expressing GAS.

Discussion

SLO activates the NLRP3 inflammasome,⁴⁴ leading to release of IL-1 β which plays an important role in the host response against GAS.²⁰⁰ In this study, we describe a novel role for SLO in regulating the levels of this cytokine by ubiquitination and degradation of pro-IL-1 β .

Pro-IL-1 β ubiquitination has previously been shown to be of the K63 or K48 linkages, leading to increased release^{118,119} or degradation,¹¹⁷ respectively. We found that the ubiquitination pattern induced in response to GAS contained a mixture of these linkages, but we have not determined the exact composition of the ubiquitin pattern, *i.e.* whether the two types of linkages are part of the same chain or stem from separate ubiquitination sites. In addition, we have not investigated whether the basal pro-IL-1 β ubiquitination seen after LPS priming is responsible for one linkage type and GAS infection the other, or whether LPS-induced ubiquitination is required for the SLO-induced process. Although the ubiquitination pattern is heterogeneous, the overall signal seems to be related to degradation, which can be blocked by the autophagy inhibitor 3-MA. However, whether pro-IL-1 β degradation actually proceeds through the autophagy pathway is uncertain at this point; 3-MA is a broad PI3K inhibitor that can interfere with several processes in addition to autophagy, and Bafilomycin, an inhibitor that prevents the formation of the autophagolysosome, did not prevent pro-IL-1 β degradation.

A lysine at position 133 in pro-IL-1 β has been identified as a site for K63-linked ubiquitination that supports increased release of mature IL-1 β ,¹¹⁹ but whether this ubiquitination site is of relevance for our observations remains to be determined. Future studies will be aimed at defining the residues involved in ubiquitination in response to GAS, and specifically mutating these residues would aid in the

investigation of the role and importance of pro-IL-1 β ubiquitination during infection. It would also be of interest to determine whether the mature, released cytokine is ubiquitinated and whether this could affect binding to IL-1R and subsequent signaling.

Presence of the inflammasome components NLRP3, ASC or caspase-1 do not seem to be required for pro-IL-1 β ubiquitination induced by GAS, nor does inflammasome activation. In contrast, ubiquitination is increased in macrophages lacking some of the inflammasome components, indicating that they may play a role in regulating the ubiquitination processes. Indeed, it has been suggested that inflammasome proteins may be involved in non-inflammasome events, such as regulation of metabolic pathways.³⁸⁹

Several pathogens exploit the ubiquitin system in general, including its role in inflammasome activation; *E. coli* prevents NLRP3 deubiquitination¹⁰³ and a ubiquitin ligase from human papilloma virus induces IL-1 β degradation.¹⁰⁴ Whether SLO-mediated ubiquitination belongs to this category or if pro-IL-1 β degradation is a regulatory mechanism induced by the host is a question for further studies.

Another question that remains to be answered is the identity of the ubiquitin ligase responsible for pro-IL-1 β ubiquitination, and whether it comes from GAS or the host. The requirement of an infection setting, *i.e.* that no ubiquitination occurs when recombinant SLO alone is sensed by the cell, suggests that additional bacterial factors are involved, possibly including a bacterial E3-ligase.

Another involved factor might be translocated NADase, and the striking similarities between the requirements for ubiquitination and CMT provides some support for this speculation. However, studying this possibility is complicated by the fact that NADase stabilizes SLO and that full SLO stability requires that NADase expresses a domain needed for its translocation.³⁰⁷ To investigate the involvement of NADase in SLO-mediated ubiquitination, a translocation mutant with retained SLO stability would need to be constructed.

Paper IV: *Streptococcal* β -NAD⁺-glycohydrolase binds to and induces cytokine release from macrophages in the absence of Streptolysin O

Background and aim

Much effort has gone into elucidating the mechanism behind CMT, the SLO-dependent translocation of NADase into host cell cytosols. For CMT to occur, the co-association of SLO and NADase at the cell surface requires that NADase binds to a yet unknown receptor; a binding event that also requires the presence of SLO.³¹² During our previous studies, we unexpectedly found NADase activity associated

with the membrane even in the absence of SLO, which prompted our further investigation.

Key findings

- NADase associates to the plasma membrane in absence of SLO during GAS infection of macrophages.
- Recombinant NADase binds to a cell surface protein independently of other bacterial proteins.
- Recombinant NADase induces MyD88- and TRIF-dependent cytokine release from macrophages expressing TLR4 and CD14.
- TLR4 and CD14 are not required for NADase binding to macrophages.
- Inactive recombinant NADase harbouring a G330D mutation exhibit decreased binding to macrophages and does not induce cytokine release.

Discussion

SLO has previously been suggested to induce TLR4 activation, leading to production of inflammatory cytokines.³⁵⁵ We also propose involvement of this receptor in GAS pathogenesis in the downstream signaling pathways induced upon NADase binding to macrophages. However, as NADase does not seem to bind directly to TLR4 or CD14 it is currently unclear how the receptor is activated: does NADase induce release of an endogenous TLR4 agonist or does NADase binding to one or several of the surface proteins involved in TLR4 signaling? Cross-linking of NADase when present on the macrophage surface and subsequent immunoprecipitation using an anti-NADase antibody might let us identify the different components involved in NADase binding and cytokine induction.

NADase binding to epithelial cells involves an N-terminal region with a putative carbohydrate-binding motif.³¹² Although binding to macrophages seems to depend on a cell-surface protein, it is still possible that the carbohydrate-binding motif is important for the interaction to occur. Recombinant NADase carrying a mutation in the carbohydrate-binding motif would thus be of interest to our continued studies. A screen of surface molecules expressed on macrophages but not epithelial cells could narrow down the list of possible candidates for NADase binding. In addition, future studies should investigate whether the interaction between NADase and macrophages is unique or whether it extends to *e.g.* other innate immune cells such as DCs and neutrophils and, importantly, to cells of human origin.

Interestingly, in a subpopulation of MIT1 isolates, specific mutations rendering NADase enzymatically inactive have been fixed, suggesting that the presence of an inactive NADase is under positive evolutionary pressure.^{300–302} Strains expressing inactive NADase have been suggested to belong to the so-called specialist strains,

i.e. strains showing a preference for causing infections at either throat or skin, while strains expressing active NADase are generalists, equally likely to infect both tissues.³⁰¹ The pressure to express enzymatically inactive NADase is thought to partly be due to its importance in the stability and function of SLO,³⁰⁷ but might also be due to unknown roles in bacterial fitness and/or virulence. Our study suggests that active NADase induces cytokine release, while the inactive form of the protein does not, and it remains to be determined whether host or bacterium benefits from the induction of pro-inflammatory cytokines. Elevated cytokine release could be important in restricting bacterial dissemination through influx and activation of immune cells, but could also increase the permeability of tissues and vasculature, which may facilitate bacterial spread into deeper tissues and improve bacterial establishment and disease progression. Conversely, decreased cytokine release might similarly also benefit both bacterium and host. To further elucidate this newly described function of NADase and its possible role in immune regulation by GAS, studies comparing NADase-active and -inactive strains and the immune responses generated against them would be an interesting place to start.

Acknowledgements

I'd like to thank the people without whom this thesis would not have been possible. Prepare for some well-deserved superlatives!

Jenny, for taking me in as a master student all those years ago and for convincing me to battle my impostor syndrome and stay on as a PhD student. Your never-ending passion for, and curiosity in, the inflammasome field has been very inspiring and has converted me to the inflammasome-loving cult. :) Thank you for providing great support and guidance while simultaneously letting me try my own wings, I've grown so much with you as my supervisor. Our tendency to finish papers and applications in the nick of time might not have done wonders to our stress levels, but we always made it in the end so why should we change? :) I'm very happy to be able to continue working with you after my PhD, I'm not ready to leave just yet!

Dóra, for being an inspiration in terms of scientific knowledge and ability to never give up (no matter the number of failed Western blots). It's been great to follow your evolution personally as well as scientifically during these years. **Lena**, for great collaboration on our project and always being so caring. You are missed in the lab!

Christine, the back-bone of the JP/FC labs, for all your invaluable help with experiments, organization and for helping me reach things in the storage room. :) I can't imagine how we'd manage without you and your knowledge. Thank you for always taking time to give advice, whether about science or life in general.

Fredric, for great scientific input and discussion during our joint group meetings and for your enthusiasm about natural sciences in general and infection immunology in particular. Our meetings would have been a lot duller without your tangents about everything from lipid bilayer structures to the last common ancestor.

The FC group, I can't think of better people to be stuck with in our "cosy" P2 lab. **Julia**, for being a role-model in terms of balancing personal life and science and living sustainably. Our well-organized trip to Lyon is one I will always remember, and I'm very happy that Sweden won you over. **Elin**, for our in-depth discussions on everything from science to Game of Thrones-theories, which I hope we'll keep up even though we won't be seeing each other every day. I'll miss having you in my office! **Úna, Esther, Kristina and Tori**, for brightening up the lab with your cheerful personalities, inspiring me with your ambition and for keeping me company during "early" lunches.

Daniel, for your passion for research and for all our lunch conversations. Your visits to our office were always a welcome break. I look forward to visiting you in Philly! **Madde**, for all your advice about science and life and for making everyone feel welcome. I miss our running sessions and races! **Mo**, for your positive mood and for always being up for dinner or a party. Thank you for working hard to make our environment better scientifically as well as socially. **Thorsten**, for all the innebandy games and lunch talks. Your knowledge has been very inspiring (and slightly intimidating). **Gudrun**, tack för all hjälp och alla råd. Utan dig skulle D14 inte vara detsamma!

All other former and present members of D14, for making me feel at home and for creating a great environment to work in. Thank you for all the scientific feedback as well as all the after-works, fikas and for always being up for dancing at dissertation parties. I expect no less during mine!

Björn, Jenny, Manja, Martina, Sara, Sarah, for helping me take my mind off immunology with board games and parties. I'm lucky we all decided to study in Lund at the same time.

Mom, pappa, for encouraging my interest in science from an early age and for always being enthusiastic and supportive of what I do.

Viktor, for always being able to make me laugh. I couldn't have done this without you.

References

1. Turvey SE, Broide DH. Innate immunity. *J Allergy Clin Immunol.* 2010;125(2):191-197. doi:10.1007/978-1-4419-5774-0_16
2. Iwasaki A, Medzhitov R. Control of adaptive immunity by the innate immune system. *Nat Immunol.* 2015;16(4):343-353. doi:10.1038/ni.3123
3. Murray PJ, Wynn TA. Protective and pathogenic functions of macrophage subsets. *Nat Rev Immunol.* 2011;11(1474-1741 (Electronic)). doi:10.1038/nri3073
4. Jain A, Pasare C. Innate Control of Adaptive Immunity: Beyond the Three-Signal Paradigm. *J Immunol.* 2017;198(10):3791-3800. doi:10.4049/jimmunol.1602000
5. Hussell T, Bell TJ. Alveolar macrophages: plasticity in a tissue-specific context. *Nat Rev Immunol.* 2014;14(2):81-93. doi:10.1038/nri3600
6. Bonilla FA, Oettgen HC. Adaptive Immunity. *J Allergy Clin Immunol.* 2013;1-2(2):20-29. doi:10.1016/B978-0-323-08593-9.00003-6
7. Paul WE. Bridging innate and adaptive immunity. *Cell.* 2011;147(6):1212-1215. doi:10.1016/j.cell.2011.11.036
8. Hoffman W, Lakkis FG, Chalasani G. B cells, antibodies, and more. *Clin J Am Soc Nephrol.* 2016;11(1):137-154. doi:10.2215/CJN.09430915
9. Janeway CAJ. Approaching the asymptote? Evolution and revolution in immunology. *Cold Spring Harb Symp Quant Biol.* 1989;54 Pt 1(9):1-13. <http://www.ncbi.nlm.nih.gov/pubmed/24141854>.
10. Janeway CA, Medzhitov R, Preston-Hurlburt P. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature.* 1997;388(6640):394-397. doi:10.1038/41131
11. Lemaitre B, Nicolas E, Michaut L, Reichhart J, Hoffmann JA. The Dorsoventral Regulatory Gene Cassette *spätzle/Toll/cactus* Controls the Potent Antifungal Response in *Drosophila* Adults. *Cell.* 1996;86(imd):973-983.
12. Poltorak A, He X, Smirnova I, et al. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in *Tlr4* gene. *Science (80-).* 1998;282(5396):2085-2088. doi:10.1126/science.282.5396.2085
13. Liston A, Masters SL. Homeostasis-altering molecular processes as mechanisms of inflammasome activation. *Nat Rev Immunol.* 2017;17(3):208-214. doi:10.1038/nri.2016.151
14. Brubaker SW, Bonham KS, Zanoni I, Kagan JC. *Innate Immune Pattern Recognition: A Cell Biological Perspective.* Vol 33.; 2015. doi:10.1146/annurev-immunol-032414-112240
15. Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: Update on toll-like

- receptors. *Nat Immunol.* 2010;11(5):373-384. doi:10.1038/ni.1863
16. Kawai T, Akira S. Toll-like Receptors and Their Crosstalk with Other Innate Receptors in Infection and Immunity. *Immunity.* 2011;34(5):637-650. doi:10.1016/j.immuni.2011.05.006
 17. Kawasaki T, Kawai T. Toll-like receptor signaling pathways. *Front Immunol.* 2014. doi:10.3389/fimmu.2014.00461
 18. Medzhitov R. Toll-like Receptors and Innate Immunity. *Nat Rev Immunol.* 2001;1(1):49-54. doi:10.1038/35100529
 19. Hornig T, Barton GM, Flavell RA, Medzhitov R. The adaptor molecule TIRAP provides signalling specificity for Toll-like receptors. *Nature.* 2002;420(6913):329-333. doi:10.1038/nature01180
 20. Kawai T, Adachi O, Ogawa T, Takeda K, Akira S. Unresponsiveness of MyD88-deficient mice to endotoxin. *Immunity.* 1999;11(1):115-122. doi:10.1007/s11103-011-9767-z.Plastid
 21. Zanoni I, Ostuni R, Marek LR, et al. CD14 controls the LPS-induced endocytosis of Toll-like receptor 4. *Cell.* 2011;147(4):868-880. doi:10.1016/j.cell.2011.09.051
 22. Yamamoto M, Sato S, Hemmi H, et al. Role of Adaptor TRIF in the MyD88-Independent Toll-Like Receptor Signaling Pathway. *Science (80-).* 2003;301:640-643.
 23. Motwani M, Pesiridis S, Fitzgerald KA. DNA sensing by the cGAS–STING pathway in health and disease. *Nat Rev Genet.* 2019. doi:10.1038/s41576-019-0151-1
 24. Broz P, Dixit VM. Inflammasomes: mechanism of assembly, regulation and signalling. *Nat Rev Immunol.* 2016;16(7):407-420. doi:10.1038/nri.2016.58
 25. Martinon F, Burns K, Tschopp J. The Inflammasome: A molecular platform triggering activation of inflammatory caspases and processing of proIL- β . *Mol Cell.* 2002;10(2):417-426. doi:10.1016/S1097-2765(02)00599-3
 26. Martinon F, Mayor A, Tschopp J. The inflammasomes: guardians of the body. *Annu Rev Immunol.* 2009;27:229-265. doi:10.1146/annurev.immunol.021908.132715 [doi]
 27. Dick MS, Sborgi L, Ruhl S, Hiller S, Broz P. ASC filament formation serves as a signal amplification mechanism for inflammasomes. *Nat Commun.* 2016;7:11929. doi:10.1038/ncomms11929
 28. Baroja-Mazo A, Martín-Sánchez F, Gomez AI, et al. The NLRP3 inflammasome is released as a particulate danger signal that amplifies the inflammatory response. *Nat Immunol.* 2014;15(8):738-748. doi:10.1038/ni.2919
 29. Rauch I, Deets KA, Ji DX, et al. NAIP-NLRC4 Inflammasomes Coordinate Intestinal Epithelial Cell Expulsion with Eicosanoid and IL-18 Release via Activation of Caspase-1 and -8. *Immunity.* 2017;46(4):649-659. doi:10.1016/j.immuni.2017.03.016
 30. Guo H, Callaway JB, Ting JP. Inflammasomes: mechanism of action, role in disease, and therapeutics. *Nat Med.* 2015;21(7):677-687. doi:10.1038/nm.3893
 31. von Moltke J, Ayres JS, Kofoed EM, Chavarria-Smith J, Vance RE. Recognition of bacteria by inflammasomes. *Annu Rev Immunol.* 2013;31:73-106. doi:10.1146/annurev-immunol-032712-095944
 32. Anand PK, Malireddi S, Lukens JR, et al. NLRP6 negatively regulates innate immunity and host defence

- against bacterial pathogens. *Nature*. 2012;488(1476-4687 (Electronic)):389-394. doi:10.1038/nature11250
33. Hornung V, Ablasser A, Charrel-Dennis M, et al. AIM2 recognizes cytosolic dsDNA and forms a caspase-1-activating inflammasome with ASC. *Nature*. 2009;458(7237):514-518. doi:10.1038/nature07725
34. Fernandes-Alnemri T, Yu JW, Datta P, Wu J, Alnemri ES. AIM2 activates the inflammasome and cell death in response to cytoplasmic DNA. *Nature*. 2009;458(7237):509-513. doi:10.1038/nature07710
35. Chavarria-Smith J, Vance RE. Direct proteolytic cleavage of NLRP1B is necessary and sufficient for inflammasome activation by anthrax lethal factor. *PLoS Pathog*. 2013;9(6):e1003452. doi:10.1371/journal.ppat.1003452
36. Sandstrom A, Mitchell PS, Goers L, Mu EW, Lesser CF, Vance RE. Functional degradation: A mechanism of NLRP1 inflammasome activation by diverse pathogen enzymes. *Science (80-)*. 2019;1330(March):eaau1330. doi:10.1126/science.aau1330
37. Zhao Y, Yang J, Shi J, et al. The NLRC4 inflammasome receptors for bacterial flagellin and type III secretion apparatus. *Nature*. 2011;477(1476-4687 (Electronic)). doi:10.1038/nature10510
38. Lightfield KL, Persson J, Brubaker SW, et al. Critical function for Naip5 in inflammasome activation by a conserved carboxy-terminal domain of flagellin. *Nat Immunol*. 2008;9(1529-2916 (Electronic)). doi:10.1038/ni.1646
39. Vance RE. The NAIP/NLRC4 inflammasomes. *Curr Opin Immunol*. 2015;32(1879-0372 (Electronic)):84-89. doi:10.1016/j.coi.2015.01.010
40. Park YH, Wood G, Kastner DL, Chae JJ. Pyrin inflammasome activation and RhoA signaling in the autoinflammatory diseases FMF and HIDS. *Nat Immunol*. 2016;17(8):914-921. doi:10.1038/ni.3457
41. Gros Lambert M, Py B. Spotlight on the NLRP3 inflammasome pathway. *J Inflamm Res*. 2018;Volume 11:359-374. doi:10.2147/JIR.S141220
42. Mariathasan S, Weiss DS, Newton K, et al. Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature*. 2006;440(7081):228-232. doi:10.1038/nature04515
43. Muñoz-Planillo R, Kuffa P, Martínez-Colón G, Smith B, Rajendiran T, Núñez G. K⁺ Efflux Is the Common Trigger of NLRP3 Inflammasome Activation by Bacterial Toxins and Particulate Matter. *Immunity*. 2013;38(6):1142-1153. <http://linkinghub.elsevier.com/retrieve/pii/S1074761313002434>.
44. Harder J, Franchi L, Munoz-Planillo R, Park JH, Reimer T, Nunez G. Activation of the Nlrp3 inflammasome by *Streptococcus pyogenes* requires streptolysin O and NF-kappa B activation but proceeds independently of TLR signaling and P2X7 receptor. *J Immunol*. 2009;183(9):5823-5829. doi:10.4049/jimmunol.0900444
45. Wang X, Jiang W, Yan Y, et al. RNA viruses promote activation of the NLRP3 inflammasome through a RIP1-RIP3-DRP1 signaling pathway. *Nat Immunol*. 2014;15(1529-2916 (Electronic)). doi:10.1038/ni.3015
46. Kayagaki N, Warming S, Lamkanfi M, et al. Non-canonical inflammasome activation targets caspase-11. *Nature*. 2011;479(7371):117-121. doi:10.1038/nature10558

47. Kayagaki N, Gonzalez LC, Akashi-takamura S, et al. Noncanonical Inflammasome Activation by Intracellular LPS Independent of TLR4. *Science* (80-). 2013;1246(September):1246-1250. doi:10.5061/dryad.bt51g
48. Shi J, Zhao Y, Wang Y, et al. Inflammatory caspases are innate immune receptors for intracellular LPS. *Nature*. 2014;514(7521):187-192. doi:10.1038/nature13683
49. Gaidt MM, Ebert TS, Chauhan D, et al. Human Monocytes Engage an Alternative Inflammasome Pathway. *Immunity*. 2016;44(4):833-846. doi:10.1016/j.immuni.2016.01.012
50. Gurung P, Anand PK, Malireddi RKS, et al. FADD and Caspase-8 Mediate Priming and Activation of the Canonical and Noncanonical Nlrp3 Inflammasomes. *J Immunol*. 2014;192(4):1835-1846. doi:10.4049/jimmunol.1302839
51. Bauernfeind FG, Horvath G, Stutz A, et al. Cutting Edge: NF- B Activating Pattern Recognition and Cytokine Receptors License NLRP3 Inflammasome Activation by Regulating NLRP3 Expression. *J Immunol*. 2009;183(2):787-791. doi:10.4049/jimmunol.0901363
52. Ghonime MG, Shamaa OR, Das S, et al. Inflammasome priming by lipopolysaccharide is dependent upon ERK signaling and proteasome function. *J Immunol*. 2014;192(8):3881-3888. doi:10.4049/jimmunol.1301974
53. Kang TB, S-H Y, Toth B, Kovalenko A, Wallach D. Caspase-8 blocks kinase RIPK3-mediated activation of the NLRP3 inflammasome. *Immunity*. 2013;38(1097-4180 (Electronic)):27-40.
54. Lin KM, Hu W, Troutman TD, et al. IRAK-1 bypasses priming and directly links TLRs to rapid NLRP3 inflammasome activation. *Proc Natl Acad Sci U S A*. 2014;111(1091-6490 (Electronic)).
55. Kang S, Fernandes-Alnemri T, Rogers C, et al. Caspase-8 scaffolding function and MLKL regulate NLRP3 inflammasome activation downstream of TLR3. *Nat Commun*. 2015;6(May):1-15. doi:10.1038/ncomms8515
56. Juliana C, Fernandes-Alnemri T, Kang S, Farias A, Qin F, Alnemri ES. Non-transcriptional priming and deubiquitination regulate NLRP3 inflammasome activation. *J Biol Chem*. 2012;287(43):36617-36622. doi:10.1074/jbc.M112.407130
57. Bauernfeind F, Bartok E, Rieger A, Franchi L, Nunez G, Hornung V. Cutting Edge: Reactive Oxygen Species Inhibitors Block Priming, but Not Activation, of the NLRP3 Inflammasome. *J Immunol*. 2011;187(2):613-617. doi:10.4049/jimmunol.1100613
58. Palazón-Riquelme P, Worboys JD, Green J, et al. USP7 and USP47 deubiquitinases regulate NLRP3 inflammasome activation. *EMBO Rep*. 2018:e44766. doi:10.15252/embr.201744766
59. Kawashima A, Karasawa T, Tago K, et al. ARIH2 Ubiquitinates NLRP3 and Negatively Regulates NLRP3 Inflammasome Activation in Macrophages. *J Immunol*. 2017;199(10):3614-3622. doi:10.4049/jimmunol.1700184
60. Py BF, Kim MS, Vakifahmetoglu-Norberg H, Yuan J. Deubiquitination of NLRP3 by BRCC3 Critically Regulates Inflammasome Activity. *Mol Cell*. 2013;49(2):331-338. doi:10.1016/j.molcel.2012.11.009
61. Han SH, Lear TB, Jerome JA, et al. Lipopolysaccharide primes the NALP3 inflammasome by inhibiting its ubiquitination and degradation mediated by the SC^{FBXL2} E3 ligase. *J Biol Chem*. 2015;290(29):18124-

18133. doi:10.1074/jbc.M115.645549

62. Song H, Liu B, Huai W, et al. The E3 ubiquitin ligase TRIM31 attenuates NLRP3 inflammasome activation by promoting proteasomal degradation of NLRP3. *Nat Commun.* 2016;7:1-11. doi:10.1038/ncomms13727
63. Ren G, Zhang X, Xiao Y, et al. ABRO1 promotes NLRP3 inflammasome activation through regulation of NLRP3 deubiquitination. *EMBO J.* 2019:e100376. doi:10.15252/embj.2018100376
64. Humphries F, Bergin R, Jackson R, et al. The E3 ubiquitin ligase Pellino2 mediates priming of the NLRP3 inflammasome. *Nat Commun.* 2018;9(1). doi:10.1038/s41467-018-03669-z
65. Song N, Liu ZS, Xue W, et al. NLRP3 Phosphorylation Is an Essential Priming Event for Inflammasome Activation. *Mol Cell.* 2017;68(1):185-197.e6. doi:10.1016/j.molcel.2017.08.017
66. Stutz A, Kolbe C-C, Stahl R, et al. NLRP3 inflammasome assembly is regulated by phosphorylation of the pyrin domain. *J Exp Med.* 2017;214(6):1725-1736. doi:10.1084/jem.20160933
67. Spalinger MR, Lang S, Gottier C, et al. PTPN22 regulates NLRP3-mediated IL1B secretion in an autophagy-dependent manner. *Autophagy.* 2017;13(9):1590-1601. doi:10.1080/15548627.2017.1341453
68. Guan K, Wei C, Zheng Z, et al. MAVS Promotes Inflammasome Activation by Targeting ASC for K63-Linked Ubiquitination via the E3 Ligase TRAF3. *J Immunol.* 2015;194(10):4880-4890. doi:10.4049/jimmunol.1402851
69. Rodgers MA, Bowman JW, Fujita H, et al. The linear ubiquitin assembly complex (LUBAC) is essential for NLRP3 inflammasome activation. *J Exp Med.* 2014;211(7):1333-1347. doi:10.1084/jem.20132486
70. Shi CS, Shenderov K, Huang NN, et al. Activation of autophagy by inflammatory signals limits IL-1 β production by targeting ubiquitinated inflammasomes for destruction. *Nat Immunol.* 2012;13(3):255-263. doi:10.1038/ni.2215
71. Lee JY, Seo D, You J, et al. The deubiquitinating enzyme, ubiquitin-specific peptidase 50, regulates inflammasome activation by targeting the ASC adaptor protein. *FEBS Lett.* 2017;591(3):479-490. doi:10.1002/1873-3468.12558
72. Hara H, Tsuchiya K, Kawamura I, et al. Phosphorylation of the adaptor ASC acts as a molecular switch that controls the formation of speck-like aggregates and inflammasome activity. *Nat Immunol.* 2013;14(12):1247-1255. doi:10.1038/ni.2749
73. Labbé K, McIntire CR, Doiron K, Leblanc PM, Saleh M. Cellular Inhibitors of Apoptosis Proteins cIAP1 and cIAP2 Are Required for Efficient Caspase-1 Activation by the Inflammasome. *Immunity.* 2011;35(6):897-907. doi:10.1016/j.immuni.2011.10.016
74. Compan V, Baroja-Mazo A, Lopez-Castejon G, et al. Cell volume regulation modulates NLRP3 inflammasome activation. *Immunity.* 2012;37(10):97-4180 (Electronic). doi:10.1016/j.immuni.2012.06.013
75. He Y, Zeng MY, Yang D, Motro B, Nunez G. NEK7 is an essential mediator of NLRP3 activation downstream of potassium efflux. *Nature.* 2016;530(7590):354-357. doi:10.1038/nature16959
76. Shi H, Wang Y, Li X, et al. NLRP3 activation and mitosis are mutually exclusive events coordinated by NEK7, a new inflammasome component. *Nat Immunol.* 2016;17(3):250-258. doi:10.1038/ni.3333

77. Schmid-Burgk JL, Chauhan D, Schmidt T, et al. A Genome-wide CRISPR (Clustered Regularly Interspaced Short Palindromic Repeats) Screen Identifies NEK7 as an Essential Component of NLRP3 Inflammasome Activation. *J Biol Chem*. 2016;291(1):103-109. doi:10.1074/jbc.C115.700492
78. Sharif H, Wang L, Wang WL, et al. Structural mechanism for NEK7-licensed activation of NLRP3 inflammasome. *Nature*. 2019. doi:10.1038/s41586-019-1295-z
79. Indramohan M, Stehlik C, Dorfleutner A. COPs and POPs Patrol Inflammasome Activation. *J Mol Biol*. 2018;430(2):153-173. doi:10.1016/j.jmb.2017.10.004
80. Gurung P, Li B, Subbarao Malireddi RK, Lamkanfi M, Geiger TL, Kanneganti TD. Chronic TLR Stimulation Controls NLRP3 Inflammasome Activation through IL-10 Mediated Regulation of NLRP3 Expression and Caspase-8 Activation. *Sci Rep*. 2015;5:14488. doi:10.1038/srep14488
81. Boucher D, Monteleone M, Coll RC, et al. Caspase-1 self-cleavage is an intrinsic mechanism to terminate inflammasome activity. *J Exp Med*. 2018. doi:10.1084/jem.20172222
82. Aachoui Y, Leaf IA, Hagar JA, et al. Caspase-11 protects against bacteria that escape the vacuole. *Science (80-)*. 2013;339(6122):975-978. doi:science.1230751 [pii] 10.1126/science.1230751 [doi]
83. Kayagaki N, Stowe IB, Lee BL, et al. Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. *Nature*. 2015;526(7575):666-671. doi:10.1038/nature15541
84. Ruhl S, Broz P. Caspase-11 activates a canonical NLRP3 inflammasome by promoting K(+) efflux. *Eur J Immunol*. 2015;45(10):2927-2936. doi:10.1002/eji.201545772
85. Yang D, He Y, Munoz-Planillo R, Liu Q, Nunez G. Caspase-11 Requires the Pannexin-1 Channel and the Purinergic P2X7 Pore to Mediate Pyroptosis and Endotoxic Shock. *Immunity*. 2015;43(5):923-932. doi:10.1016/j.immuni.2015.10.009
86. Zanoni I, Tan Y, Gioia M Di, et al. An endogenous caspase-11 ligand elicits IL-1 release from living DCs. *Science (80-)*. 2016;3036(April):1-9.
87. Zanoni I, Tan Y, Di Gioia M, Springstead JR, Kagan JC. By Capturing Inflammatory Lipids Released from Dying Cells, the Receptor CD14 Induces Inflammasome-Dependent Phagocyte Hyperactivation. *Immunity*. 2017;47(4):697-709.e3. doi:10.1016/j.immuni.2017.09.010
88. Evavold CL, Ruan J, Tan Y, Xia S, Wu H, Kagan JC. The Pore-Forming Protein Gasdermin D Regulates Interleukin-1 Secretion from Living Macrophages. *Immunity*. 2017;48(1):35-44.e6. doi:10.1016/j.immuni.2017.11.013
89. LaRock CN, Nizet V. Inflammasome/IL-1beta Responses to Streptococcal Pathogens. *Front Immunol*. 2015;6:518. doi:10.3389/fimmu.2015.00518
90. Shi J, Zhao Y, Wang K, et al. Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature*. 2015;526(7575):660-665. doi:10.1038/nature15514
91. Liu X, Zhang Z, Ruan J, et al. Inflammasome-activated gasdermin D causes pyroptosis by forming membrane pores. *Nature*. 2016;535(7610):153-158. doi:10.1038/nature18629
92. Sborgi L, Ruhl S, Mulvihill E, et al. GSDMD membrane pore formation constitutes the mechanism of pyroptotic cell death. *EMBO J*. 2016;35(16):1766-1778. doi:10.15252/embj.201694696

93. Liu T, Yamaguchi Y, Shirasaki Y, et al. Single-cell imaging of caspase-1 dynamics reveals an all-or-none inflammasome signaling response. *Cell Rep.* 2014;8(4):974-982. doi:10.1016/j.celrep.2014.07.012
94. Brough D, Rothwell NJ. Caspase-1-dependent processing of pro-interleukin-1beta is cytosolic and precedes cell death. *J Cell Sci.* 2007;120(Pt 5):772-781. doi:10.1242/jcs.03377
95. Chen KW, Groß CJ, Sotomayor FV, et al. The Neutrophil NLRC4 Inflammasome Selectively Promotes IL-1 β Maturation without Pyroptosis during Acute Salmonella Challenge. *Cell Rep.* 2014;8(2):570-582. doi:10.1016/j.celrep.2014.06.028
96. Karmakar M, Katsnelson MA, Dubyak GR, Pearlman E. Neutrophil P2X7 receptors mediate NLRP3 inflammasome-dependent IL-1beta secretion in response to ATP. *Nat Commun.* 2016;7:10555. doi:10.1038/ncomms10555
97. Karmakar M, Katsnelson M, Malak HA, et al. Neutrophil IL-1beta processing induced by pneumolysin is mediated by the NLRP3/ASC inflammasome and caspase-1 activation and is dependent on K⁺ efflux. *J Immunol.* 2015;(1550-6606 (Electronic)). doi:10.4049/jimmunol.1401624
98. Broz P, Von Moltke J, Jones JW, Vance RE, Monack DM. Differential requirement for caspase-1 autoproteolysis in pathogen-induced cell death and cytokine processing. *Cell Host Microbe.* 2010;8(6):471-483. doi:10.1016/j.chom.2010.11.007
99. Rühl S, Shkarina K, Demarco B, Heilig R, Santos JC, Broz P. ESCRT-dependent membrane repair negatively regulates pyroptosis downstream of GSDMD activation. *Science (80-).* 2018;362(6417):956-960. doi:10.1126/science.aar7607
100. DiPeso L, Ji DX, Vance RE, Price J V. Cell death and cell lysis are separable events during pyroptosis. *Cell Death Discov.* 2017;3(June):17070. doi:10.1038/cddiscovery.2017.70
101. Suzuki S, Mimuro H, Kim M, et al. *Shigella* IpaH7.8 E3 ubiquitin ligase targets glomulin and activates inflammasomes to demolish macrophages. *Proc Natl Acad Sci.* 2014;111(40):E4254-E4263. doi:10.1073/pnas.1324021111
102. Shin S, Brodsky IE. The inflammasome: Learning from bacterial evasion strategies. *Semin Immunol.* 2015;27(2):102-110. doi:10.1016/j.smim.2015.03.006
103. Yen H, Sugimoto N, Tobe T. Enteropathogenic *Escherichia coli* Uses NleA to Inhibit NLRP3 Inflammasome Activation. *PLoS Pathog.* 2015;11(9):1-23. doi:10.1371/journal.ppat.1005121
104. Niebler M, Qian X, Höfler D, et al. Post-Translational Control of IL-1 β via the Human Papillomavirus Type 16 E6 Oncoprotein: A Novel Mechanism of Innate Immune Escape Mediated by the E3-Ubiquitin Ligase E6-AP and p53. *PLoS Pathog.* 2013;9(8). doi:10.1371/journal.ppat.1003536
105. Larock CN, Cookson BT. The yersinia virulence effector YopM binds caspase-1 to arrest inflammasome assembly and processing. *Cell Host Microbe.* 2012;12(6):799-805. doi:10.1016/j.chom.2012.10.020
106. Johnston JB, Barrett JW, Nazarian SH, et al. A poxvirus-encoded pyrin domain protein interacts with ASC-1 to inhibit host inflammatory and apoptotic responses to infection. *Immunity.* 2005;23(6):587-598. doi:10.1016/j.immuni.2005.10.003
107. Schotte P, Denecker G, Van Den Broeke A, Vandenabeele P, Cornelis GR, Beyaert R. Targeting Rac1 by the Yersinia effector protein YopE inhibits caspase-1-mediated maturation and release of interleukin-

- 1 β . *J Biol Chem*. 2004;279(24):25134-25142. doi:10.1074/jbc.M401245200
108. Galle M, Schotte P, Haegman M, et al. The *Pseudomonas aeruginosa* Type III secretion system plays a dual role in the regulation of caspase-1 mediated IL-1 β maturation. *J Cell Mol Med*. 2008;12(5A):1767-1776. doi:10.1111/j.1582-4934.2007.00190.x
109. Perez-Lopez A, Rosales-Reyes R, Alpuche-Aranda CM, Ortiz-Navarrete V. Salmonella Downregulates Nod-like Receptor Family CARD Domain Containing Protein 4 Expression To Promote Its Survival in B Cells by Preventing Inflammasome Activation and Cell Death. *J Immunol*. 2013;190(3):1201-1209. doi:10.4049/jimmunol.1200415
110. Shimada T, Park BG, Wolf AJ, et al. Staphylococcus aureus Evades Lysozyme-Based Peptidoglycan Digestion that Links Phagocytosis, Inflammasome Activation, and IL-1 β Secretion. *Cell Host Microbe*. 2010;7(1):38-49. doi:10.1016/j.chom.2009.12.008
111. Malynn BA, Ma A. Ubiquitin Makes Its Mark on Immune Regulation. *Immunity*. 2010;33(6):843-852. doi:10.1016/j.immuni.2010.12.007
112. Komander D, Rape M. The Ubiquitin Code. *Annu Rev Biochem*. 2012;81(1):203-229. doi:10.1146/annurev-biochem-060310-170328
113. Swatek KN, Komander D. Ubiquitin modifications. *Cell Res*. 2016;26(4):399-422. doi:10.1038/cr.2016.39
114. Kwon YT, Ciechanover A. The Ubiquitin Code in the Ubiquitin-Proteasome System and Autophagy. *Trends Biochem Sci*. 2017;42(11):873-886. doi:10.1016/j.tibs.2017.09.002
115. Dikic I, Wakatsuki S, Walters KJ. Ubiquitin-binding domains from structures to functions. *Nat Rev Mol Cell Biol*. 2009;10(10):659-671. doi:10.1038/nrm2767
116. Kimura Y, Tanaka K. Regulatory mechanisms involved in the control of ubiquitin homeostasis. *J Biochem*. 2010;147(6):793-798. doi:10.1093/jb/mvq044
117. Eldridge MJG, Sanchez-Garrido J, Hoben GF, Goddard PJ, Shenoy AR. The Atypical Ubiquitin E2 Conjugase UBE2L3 Is an Indirect Caspase-1 Target and Controls IL-1 β Secretion by Inflammasomes. *Cell Rep*. 2017;18(5):1285-1297. doi:10.1016/j.celrep.2017.01.015
118. Zhang L, Liu Y, Wang B, et al. POH1 deubiquitinates pro-interleukin-1 β and restricts inflammasome activity. *Nat Commun*. 2018;9(1). doi:10.1038/s41467-018-06455-z
119. Duong BH, Onizawa M, Oses-Prieto JA, et al. A20 restricts ubiquitination of pro-interleukin-1beta protein complexes and suppresses NLRP3 inflammasome activity. *Immunity*. 2015;42(1):55-67. doi:10.1016/j.immuni.2014.12.031
120. Jacobson KA, Muller CE. Medicinal chemistry of adenosine, P2Y and P2X receptors. *Neuropharmacology*. 2016;104:31-49. doi:10.1016/j.neuropharm.2015.12.001
121. Di Virgilio F, Dal Ben D, Sarti AC, Giuliani AL, Falzoni S. The P2X7 Receptor in Infection and Inflammation. *Immunity*. 2017;47(1):15-31. doi:10.1016/j.immuni.2017.06.020
122. Burnstock G, Boeynaems JM. Purinergic signalling and immune cells. *Purinergic Signal*. 2014;10(4):529-564. doi:10.1007/s11302-014-9427-2

123. Adinolfi E, Lisa Giuliani A, De Marchi E, Pegoraro A, Orioli E, Di Virgilio F. The P2X7 receptor: a main player in inflammation. *Biochem Pharmacol.* 2017. doi:10.1016/j.bcp.2017.12.021
124. Di Virgilio F, Schmalzing G, Markwardt F. The Elusive P2X7 Macropore. *Trends Cell Biol.* 2018. doi:10.1016/j.tcb.2018.01.005
125. Hattori M, Gouaux E. Molecular mechanism of ATP binding and ion channel activation in P2X receptors. *Nature.* 2012;485(1476-4687 (Electronic)):207-213. doi:10.1038/nature11010
126. Di Virgilio F, Giuliani ALL, Vultaggio-Poma V, Falzoni S, Sarti ACC. Non-nucleotide Agonists Triggering P2X7 Receptor Activation and Pore Formation. *Front Pharmacol.* 2018;9(FEB):39. doi:10.3389/fphar.2018.00039
127. Surprenant A, Rassendren F, Kawashima E, North RA, Buell G. The cytolytic P2Z receptor for extracellular ATP identified as a P2X receptor (P2X7). *Science.* 1996;272(5262):1-9.
128. Morciano G, Sarti AC, Marchi S, et al. Use of luciferase probes to measure ATP in living cells and animals. *Nat Protoc.* 2017;12(8):1542-1562. doi:10.1038/nprot.2017.052
129. Denlinger LC, Fisetle PL, Sommer JA, et al. Cutting Edge: The Nucleotide Receptor P2X7 Contains Multiple Protein- and Lipid-Interaction Motifs Including a Potential Binding Site for Bacterial Lipopolysaccharide. *J Immunol.* 2001;167(4):1871-1876. doi:10.4049/jimmunol.167.4.1871
130. Wickert LE, Blanchette JB, Waldschmidt N V, et al. The C-terminus of human nucleotide receptor P2X7 is critical for receptor oligomerization and N-linked glycosylation. *PLoS One.* 2013;8(5):e63789. doi:10.1371/journal.pone.0063789
131. Bartlett R, Stokes L, Sluyter R. The P2X7 receptor channel: recent developments and the use of P2X7 antagonists in models of disease. *Pharmacol Rev.* 2014;66(3):638-675. doi:10.1124/pr.113.008003
132. Adinolfi E, Cirillo M, Woltersdorf R, et al. Trophic activity of a naturally occurring truncated isoform of the P2X7 receptor. *FASEB J.* 2010;24(1530-6860 (Electronic)):3393-3404.
133. Masin M, Young C, Lim K, et al. Expression, assembly and function of novel C-terminal truncated variants of the mouse P2X7 receptor: re-evaluation of P2X7 knockouts. *Br J Pharmacol.* 2012;165(4):978-993. doi:10.1111/j.1476-5381.2011.01624.x
134. Nicke A, Kuan YH, Masin M, et al. A functional P2X7 splice variant with an alternative transmembrane domain I escapes gene inactivation in P2X7 knock-out mice. *J Biol Chem.* 2009;284(38):25813-25822. doi:10.1074/jbc.M109.033134
135. Hong S, Schwarz N, Brass A, et al. Differential regulation of P2X7 receptor activation by extracellular nicotinamide adenine dinucleotide and ecto-ADP-ribosyltransferases in murine macrophages and T cells. *J Immunol.* 2009;183(1):578-592. doi:183/1/578 [pii] 10.4049/jimmunol.0900120 [doi]
136. Schwarz N, Drouot L, Nicke A, et al. Alternative splicing of the N-terminal cytosolic and transmembrane domains of P2X7 controls gating of the ion channel by ADP-ribosylation. *PLoS One.* 2012;(1932-6203 (Electronic)). doi:10.1371/journal.pone.0041269
137. Adriouch S, Bannas P, Schwarz N, et al. ADP-ribosylation at R125 gates the P2X7 ion channel by presenting a covalent ligand to its nucleotide binding site. *FASEB J.* 2008;22(3):861-869. doi:10.1096/fj.07-9294com

138. Seman M, Adriouch S, Scheuplein F, et al. NAD-induced T cell death: ADP-ribosylation of cell surface proteins by ART2 activates the cytolytic P2X7 purinoceptor. *Immunity*. 2003;19(1074-7613 (Print)).
139. Chessell IP, Hatcher JP, Bountra C, et al. Disruption of the P2X7 purinoceptor gene abolishes chronic inflammatory and neuropathic pain. *Pain*. 2005;114(3):386-396. doi:10.1016/j.pain.2005.01.002
140. Solle M, Labasi J, Perregaux DG, et al. Altered cytokine production in mice lacking P2X(7) receptors. *J Biol Chem*. 2001;276(1):125-132. doi:10.1074/jbc.M006781200
141. Taylor SR, Gonzalez-Begne M, Sojka DK, et al. Lymphocytes from P2X7-deficient mice exhibit enhanced P2X7 responses. *J Leukoc Biol*. 2009;85(1938-3673 (Electronic)):978-986. doi:10.1189/jlb.0408251
142. Adriouch S, Dox C, Welge V, Seman M, Koch-Nolte F, Haag F. Cutting edge: a natural P451L mutation in the cytoplasmic domain impairs the function of the mouse P2X7 receptor. *J Immunol*. 2002;169(0022-1767 (Print)).
143. Le Stunff H, Auger R, Kanellopoulos J, Raymond M-N. The Pro-451 to Leu polymorphism within the C-terminal tail of P2X7 receptor impairs cell death but not phospholipase D activation in murine thymocytes. *J Biol Chem*. 2004;279(0021-9258 (Print)).
144. Steinberg TH, Newman AS, Swansonq JA, Silverstein SC. ATP4- Permeabilizes the Plasma Membrane of Mouse Macrophages to Fluorescent Dyes. 1987;262(18):8884-8888.
145. Browne LE, Compan V, Bragg L, North RA. P2X7 receptor channels allow direct permeation of nanometer-sized dyes. *J Neurosci*. 2013;33(8):3557-3566. doi:10.1523/JNEUROSCI.2235-12.2013
146. Pelegrin P, Surprenant A. Pannexin-1 mediates large pore formation and interleukin-1beta release by the ATP-gated P2X7 receptor. *EMBO J*. 2006;25(0261-4189 (Print)):5071-5082. doi:10.1038/sj.emboj.7601378
147. Qu Y, Misaghi S, Newton K, et al. Pannexin-1 is required for ATP release during apoptosis but not for inflammasome activation. *J Immunol*. 2011;186(11):6553-6561. doi:10.4049/jimmunol.1100478
148. Harkat M, Peverini L, Cerdan AH, et al. On the permeation of large organic cations through the pore of ATP-gated P2X receptors. *Proc Natl Acad Sci*. 2017;114(19):E3786-E3795. doi:10.1073/pnas.1701379114
149. Pippel A, Stolz M, Woltersdorf R, Kless A, Schmalzing G, Markwardt F. Localization of the gate and selectivity filter of the full-length P2X7 receptor. *Proc Natl Acad Sci*. 2017;114(11):E2156-E2165. doi:10.1073/pnas.1610414114
150. Smart ML, Gu B, Panchal RG, et al. P2X7 receptor cell surface expression and cytolytic pore formation are regulated by a distal C-terminal region. *J Biol Chem*. 2003;278(10).
151. Becker D, Woltersdorf R, Boldt W, et al. The P2X7 carboxyl tail is a regulatory module of P2X7 receptor channel activity. *J Biol Chem*. 2008;283(37):25725-25734. doi:10.1074/jbc.M803855200
152. Karasawa A, Michalski K, Mikhelzon P, Kawate T. The P2X7 receptor forms a dye-permeable pore independent of its intracellular domain but dependent on membrane lipid composition. *Elife*. 2017;6. doi:10.7554/eLife.31186
153. Robinson LE, Shridar M, Smith P, Murrell-Lagnado RD. Plasma membrane cholesterol as a regulator of

- human and rodent P2X7 receptor activation and sensitization. *J Biol Chem.* 2014;289(46):31983-31994. doi:10.1074/jbc.M114.574699
154. Gonnord P, Delarasse C, Auger R, et al. Palmitoylation of the P2X7 receptor, an ATP-gated channel, controls its expression and association with lipid rafts. *FASEB J.* 2009;23(3):795-805. doi:10.1096/fj.08-114637
155. Elssner A, Duncan M, Gavrilin M, Wewers MD. A novel P2X7 receptor activator, the human cathelicidin-derived peptide LL37, induces IL-1 beta processing and release. *J Immunol.* 2004;172(0022-1767 (Print)):4987-4994.
156. Sanz JM, Chiozzi P, Ferrari D, et al. Activation of Microglia by Amyloid Requires P2X7 Receptor Expression. *J Immunol.* 2009;182(7):4378-4385. doi:10.4049/jimmunol.0803612
157. Ferrari D, Pizzirani C, Adinolfi E, et al. The Antibiotic Polymyxin B Modulates P2X7 Receptor Function. *J Immunol.* 2014;173(7):4652-4660. doi:10.4049/jimmunol.173.7.4652
158. Nörenberg W, Hempel C, Urban N, Sobottka H, Illes P, Schaefer M. Clemastine potentiates the human P2X7 receptor by sensitizing it to lower ATP concentrations. *J Biol Chem.* 2011;286(13):11067-11081. doi:10.1074/jbc.M110.198879
159. Guo C, Masin M, Qureshi OS, Murrell-Lagnado RD. Evidence for functional P2X4/P2X7 heteromeric receptors. *Mol Pharmacol.* 2007;72(6):1447-1456. doi:10.1124/mol.107.035980
160. Kawano A, Tsukimoto M, Mori D, et al. Regulation of P2X7-dependent inflammatory functions by P2X4 receptor in mouse macrophages. *Biochem Biophys Res Commun.* 2012;420(1):102-107. doi:10.1016/j.bbrc.2012.02.122
161. Perez-Flores G, Levesque SA, Pacheco J, et al. The P2X7/P2X4 interaction shapes the purinergic response in murine macrophages. *Biochem Biophys Res Commun.* 2015;467(3):484-490. doi:10.1016/j.bbrc.2015.10.025
162. Boumechache M, Masin M, Edwardson JM, Gorecki DC, Murrell-Lagnado R. Analysis of assembly and trafficking of native P2X4 and P2X7 receptor complexes in rodent immune cells. *J Biol Chem.* 2009;284(20):13446-13454. doi:10.1074/jbc.M901255200
163. Schneider M, Prudic K, Pippel A, et al. Interaction of purinergic P2X4 and P2X7 receptor subunits. *Front Pharmacol.* 2017;8(NOV):1-14. doi:10.3389/fphar.2017.00860
164. Young CNJ, Chira N, Rog J, et al. Sustained activation of P2X7 induces MMP-2-evoked cleavage and functional purinoceptor inhibition. *J Mol Cell Biol.* 2017. doi:10.1093/jmcb/mjx030
165. Jiang LH. Inhibition of P2X7 receptors by divalent cations: Old action and new insight. *Eur Biophys J.* 2009;38(3):339-346. doi:10.1007/s00249-008-0315-y
166. Karasawa A, Kawate T. Structural basis for subtype-specific inhibition of the P2X7 receptor. *Elife.* 2016;5(DECEMBER2016):1-17. doi:10.7554/eLife.22153
167. Pelegrin P, Surprenant A. The P2X(7) receptor-pannexin connection to dye uptake and IL-1beta release. *Purinergic Signal.* 2009;5(2):129-137. doi:10.1007/s11302-009-9141-7
168. Di A, Xiong S, Ye Z, et al. The TWIK2 Potassium Efflux Channel in Macrophages Mediates NLRP3 Inflammasome-Induced Inflammation. *Immunity.* 2018:1-10. doi:10.1016/j.immuni.2018.04.032

169. Franceschini A, Capece M, Chiozzi P, et al. The P2X7 receptor directly interacts with the NLRP3 inflammasome scaffold protein. (1530-6860 (Electronic)).
170. Dubyak GR. P2X7 receptor regulation of non-classical secretion from immune effector cells. *Cell Microbiol.* 2012;14(11):1697-1706. doi:10.1111/cmi.12001
171. Adamczyk M, Griffiths R, Dewitt S, Knauper V, Aeschlimann D. P2X7 receptor activation regulates rapid unconventional export of transglutaminase-2. *J Cell Sci.* 2015;128(24):4615-4628. doi:10.1242/jcs.175968
172. Clark AK, Wodarski R, Guida F, Sasso O, Malcangio M. Cathepsin S release from primary cultured microglia is regulated by the P2X7 receptor. *Glia.* 2010;58(14):1710-1726. doi:10.1002/glia.21042
173. Lopez-Castejon G, Theaker J, Pelegrin P, Clifton AD, Braddock M, Surprenant A. P2X(7) receptor-mediated release of cathepsins from macrophages is a cytokine-independent mechanism potentially involved in joint diseases. *J Immunol.* 2010;185(4):2611-2619. doi:10.4049/jimmunol.1000436
174. Gu BJ, Wiley JS. Rapid ATP-induced release of matrix metalloproteinase 9 is mediated by the P2X7 receptor. *Blood.* 2006;107(12):4946-4953. doi:10.1182/blood-2005-07-2994
175. Pfeiffer ZA, Aga M, Prabhu U, Watters JJ, Hall DJ, Bertics PJ. The nucleotide receptor P2X7 mediates actin reorganization and membrane blebbing in RAW 264.7 macrophages via p38 MAP kinase and Rho. *J Leukoc Biol.* 2004;75(6):1173-1182. doi:10.1189/jlb.1203648
176. Verhoef PA, Estacion M, Schilling W, Dubyak GR. P2X7 receptor-dependent blebbing and the activation of Rho-effector kinases, caspases, and IL-1 beta release. *J Immunol.* 2003;170(0022-1767 (Print)):5728-5738.
177. Valimaki E, Cypriak W, Virkanen J, et al. Calpain Activity Is Essential for ATP-Driven Unconventional Vesicle-Mediated Protein Secretion and Inflammasome Activation in Human Macrophages. *J Immunol.* 2016;197(8):3315-3325. doi:10.4049/jimmunol.1501840
178. Qu Y, Franchi L, Nunez G, Dubyak GR. Nonclassical IL-1 Secretion Stimulated by P2X7 Receptors Is Dependent on Inflammasome Activation and Correlated with Exosome Release in Murine Macrophages. *J Immunol.* 2007;179(3):1913-1925. doi:10.4049/jimmunol.179.3.1913
179. Andrei, C, Dazzi, C, Lotti, L, Torrisi, M R, Chimini, G, Rubartelli A. The Secretory Route of the Leaderless Protein Interleukin 1B Involves Exocytosis of Endolysosome-related Vesicles. *Mol Biol Cell.* 1999;10:1463-1475.
180. Andrei C, Margiocco P, Poggi A, Lotti L V, Torrisi MR, Rubartelli A. Phospholipases C and A2 control lysosome-mediated IL-1 beta secretion: Implications for inflammatory processes. *Proc Natl Acad Sci U S A.* 2004;101(26):9745-9750. doi:10.1073/pnas.0308558101
181. Ferrari D, McNamee EN, Idzko M, Gambari R, Eltzschig HK. Purinergic Signaling During Immune Cell Trafficking. *Trends Immunol.* 2016;37(6):399-411. doi:10.1016/j.it.2016.04.004
182. Qu Y, Dubyak GR. P2X7 receptors regulate multiple types of membrane trafficking responses and non-classical secretion pathways. *Purinergic Signal.* 2009;5(2):163-173. doi:10.1007/s11302-009-9132-8
183. Wiley JS, Gu BJ. A new role for the P2X7 receptor: a scavenger receptor for bacteria and apoptotic cells in the absence of serum and extracellular ATP. *Purinergic Signal.* 2012;8:579-586. doi:10.1007/s11302-

184. Janks L, Sprague RS, Egan TM. ATP-Gated P2X7 Receptors Require Chloride Channels To Promote Inflammation in Human Macrophages. *J Immunol*. 2018. doi:10.4049/jimmunol.1801101
185. Verhoef PA, Kertesz SB, Lundberg K, Kahlenberg JM, Dubyak GR. Inhibitory Effects of Chloride on the Activation of Caspase-1, IL-1 Secretion, and Cytolysis by the P2X7 Receptor. *J Immunol*. 2005;175(11):7623-7634. doi:10.4049/jimmunol.175.11.7623
186. Savio LEB, Mello P de A, da Silva CG, Coutinho-Silva R. The P2X7 receptor in inflammatory diseases: Angel or demon? *Front Pharmacol*. 2018;9(FEB). doi:10.3389/fphar.2018.00052
187. Sims JE, Smith DE. The IL-1 family: regulators of immunity. *Nat Rev Immunol*. 2010;10(2):89-102. doi:10.1038/nri2691
188. Auron PE, Webb AC, Rosenwasser LJ, et al. Nucleotide sequence of human monocyte interleukin 1 precursor cDNA. *Proc Natl Acad Sci*. 1984;81(24):7907-7911. doi:10.1073/pnas.81.24.7907
189. Lomedico PT, et al. Cloning and expression of murine interleukin-1 cDNA. *Nature*. 1984;312:458-461.
190. Dinarello CA, Renfer L, Wolff SM. Human leukocytic pyrogen: Purification and development of a radioimmunoassay. *Proc Natl Acad Sci U S A*. 1977;74(10):4624-4627.
191. Dinarello CA. A clinical perspective of IL-1 β as the gatekeeper of inflammation. *Eur J Immunol*. 2011;41(5):1203-1217. doi:10.1002/eji.201141550
192. Dinarello CA. Immunological and inflammatory functions of the interleukin-1 family. *Annu Rev Immunol*. 2009;27:519-550. doi:10.1146/annurev.immunol.021908.132612
193. Dinarello CA. Overview of the IL- - 1 family in innate inflammation and acquired immunity. *Immunol Rev*. 2018;281:8-27. doi:10.1111/imr.12621
194. Shouval DS, Biswas A, Kang YH, et al. Interleukin 1 β Mediates Intestinal Inflammation in Mice and Patients With Interleukin 10 Receptor Deficiency. *Gastroenterology*. 2016;151(6):1100-1104. doi:10.1053/j.gastro.2016.08.055
195. Guarda G, Braun M, Staehli F, et al. Type I interferon inhibits interleukin-1 production and inflammasome activation. *Immunity*. 2011;34(2):213-223. doi:10.1016/j.immuni.2011.02.006
196. Carta S, Lavieri R, Rubartelli A. Different Members of the IL-1 Family Come Out in Different Ways: DAMPs vs. Cytokines? *Front Immunol*. 2013;4:123. doi:10.3389/fimmu.2013.00123
197. Cerretti DP, Kozlosky CJ, Mosley B, et al. Molecular Cloning of the Interleukin- 1 β Converting Enzyme. *Science (80-)*. 1992;256(5053):97-100.
198. Fantuzzi G, Ku G, Harding MW, et al. Response to local inflammation of IL-1 beta-converting enzyme-deficient mice. *J Immunol*. 1997;158(4):1818-1824. <http://www.ncbi.nlm.nih.gov/pubmed/9029121>.
199. Coeshott C, Ohnemus C, Pilyavskaya A, et al. Converting enzyme-independent release of tumor necrosis factor α and IL-1 β from a stimulated human monocytic cell line in the presence of activated neutrophils or purified proteinase 3. *Proc Natl Acad Sci U S A*. 1999;96(11):6261-6266. doi:10.1073/pnas.96.11.6261
200. LaRock CN, Todd J, LaRock DL, et al. IL-1 β is an innate immune sensor of microbial proteolysis. *Sci Immunol*. 2016;1(2):eah3539. doi:10.1126/sciimmunol.aah3539

201. Beauséjour A, Goulet JP, Grenier D, DN. Proteolytic activation of the IL-1 β precursor by *Candida albicans*. *Infect Immun*. 1998;66(2):676-681.
202. Harris J, Hartman M, Roche C, et al. Autophagy controls IL-1 β secretion by targeting Pro-IL-1 β for degradation. *J Biol Chem*. 2011;286(11):9587-9597. doi:10.1074/jbc.M110.202911
203. Ainscough JS, Frank Gerberick G, Zahedi-Nejad M, et al. Dendritic cell IL-1 α and IL-1 β are polyubiquitinated and degraded by the proteasome. *J Biol Chem*. 2014;289(51):35582-35592. doi:10.1074/jbc.M114.595686
204. Lopez-Castejon G, Brough D. Understanding the mechanism of IL-1 β secretion. *Cytokine Growth Factor Rev*. 2011;22(4):189-195. doi:10.1016/j.cytogfr.2011.10.001
205. Rabouille C. Pathways of Unconventional Protein Secretion. *Trends Cell Biol*. 2017;27(3):230-240. doi:10.1016/j.tcb.2016.11.007
206. Sitia R, Rubartelli A. The unconventional secretion of IL-1 β : Handling a dangerous weapon to optimize inflammatory responses. *Semin Cell Dev Biol*. 2018;83:12-21. doi:10.1016/j.semcdb.2018.03.011
207. van Oostrum J, Priestle JP, Grütter MG, Schmitz A. The Structure of Murine Interleukin-1 β at 2.8 Å Resolution. *J Struct*. 1991;107:189-195.
208. Heilig R, Dick MS, Sborgi L, Meunier E, Hiller S, Broz P. The Gasdermin-D pore acts as a conduit for IL-1 β secretion in mice. *Eur J Immunol*. 2018;0:1-9. doi:10.1002/eji.201747404
209. Schneider KS, Gross CJ, Dreier RF, et al. The Inflammasome Drives GSDMD-Independent Secondary Pyroptosis and IL-1 Release in the Absence of Caspase-1 Protease Activity. *Cell Rep*. 2017;21(13):3846-3859. doi:10.1016/j.celrep.2017.12.018
210. Monteleone M, Stanley AC, Chen KW, et al. Interleukin-1 β Maturation Triggers Its Relocation to the Plasma Membrane for Gasdermin-D-Dependent and -Independent Secretion. *Cell Rep*. 2018;24(6):1425-1433. doi:10.1016/j.celrep.2018.07.027
211. Russo HM, Rathkey J, Boyd-Tressler A, Katsnelson MA, Abbott DW, Dubyak GR. Active Caspase-1 Induces Plasma Membrane Pores That Precede Pyroptotic Lysis and Are Blocked by Lanthanides. *J Immunol*. 2016;197(4):1353-1367. doi:10.4049/jimmunol.1600699
212. Gutierrez KD, Davis MA, Daniels BP, et al. MLKL Activation Triggers NLRP3-Mediated Processing and Release of IL-1 β Independently of Gasdermin-D. *J Immunol*. 2017;198(5):2156-2164. doi:10.4049/jimmunol.1601757
213. Verhoef, P A, Kertesz, S B, Estacion, M, Schilling W P DGR. Maitotoxin Induces Biphasic Interleukin-1 β Secretion and Membrane Blebbing in Murine Macrophages. *Mol Pharmacol*. 2004;66(4):909-9020.
214. Martín-Sánchez F, Martínez-García JJ, Muñoz-García M, et al. Lytic cell death induced by melittin bypasses pyroptosis but induces NLRP3 inflammasome activation and IL-1 β release. *Cell Death Dis*. 2017;8(8):e2984. doi:10.1038/cddis.2017.390
215. Martin-Sanchez F, Diamond C, Zeitler M, et al. Inflammasome-dependent IL-1 β release depends upon membrane permeabilisation. *Cell Death Differ*. 2016;23(7):1219-1231. doi:10.1038/cdd.2015.176
216. Hamon Y, Luciani MF, Becq F, Verrier B, Rubartelli A, Chimini G. Interleukin-1 β secretion is impaired by inhibitors of the Atp binding cassette transporter, ABC1. *Blood*. 1997;90(8):2911-2915.

217. Marty V, Medina C, Combe C, Parnet P, Amedee T. ATP binding cassette transporter ABC1 is required for the release of interleukin-1beta by P2X7-stimulated and lipopolysaccharide-primed mouse Schwann cells. *Glia*. 2005;49(4):511-519. doi:10.1002/glia.20138
218. Zhou X, Engel T, Goepfert C, Erren M, Assmann G, von Eckardstein A. The ATP binding cassette transporter A1 contributes to the secretion of interleukin 1 β from macrophages but not from monocytes. *Biochem Biophys Res Commun*. 2002;291(3):598-604. doi:10.1006/bbrc.2002.6473
219. Lamkanfi M, Mueller JL, Vitari AC, et al. Glyburide inhibits the Cryopyrin/Nalp3 inflammasome. *J Cell Biol*. 2009;187(1):61-70. doi:10.1083/jcb.200903124
220. Dahl G, Qiu F, Wang J. The bizarre pharmacology of the ATP release channel pannexin1. *Neuropharmacology*. 2013;75:583-593. doi:10.1016/j.neuropharm.2013.02.019
221. Pelegrin P, Barroso-Gutierrez C, Surprenant A. P2X7 receptor differentially couples to distinct release pathways for IL-1beta in mouse macrophage. *J Immunol*. 2008;180(0022-1767 (Print)):7147-7157.
222. Parzych K, Zetterqvist A V, Wright WR, Kirkby NS, Mitchell JA, Paul-Clark MJ. Differential role of pannexin-1/ATP/P2X7 axis in IL-1beta release by human monocytes. *FASEB J*. 2017;31(6):2439-2445. doi:10.1096/fj.201600256
223. Chen KW, Demarco B, Broz P. Pannexin-1 promotes NLRP3 activation during apoptosis but is dispensable for canonical or Non-canonical inflammasome activation. *Eur J Immunol*. 2019;eji.201948254. doi:10.1002/eji.201948254
224. Temmerman K, Ebert AD, Müller HM, Sinning I, Tews I, Nickel W. A direct role for phosphatidylinositol-4,5-bisphosphate in unconventional secretion of fibroblast growth factor 2. *Traffic*. 2008;9(7):1204-1217. doi:10.1111/j.1600-0854.2008.00749.x
225. Brough D, Pelegrin P, Nickel W. An emerging case for membrane pore formation as a common mechanism for the unconventional secretion of FGF2 and IL-1beta. *J Cell Sci*. 2017;130(19):3197-3202. doi:10.1242/jcs.204206
226. Steringer JP, Lange S, Čujová S, et al. Key steps in unconventional secretion of fibroblast growth factor 2 reconstituted with purified components. *Elife*. 2017;6:1-36. doi:10.7554/elife.28985
227. Blott EJ, Griffiths GM. Secretory lysosomes. *Nat Rev Mol Cell Biol*. 2002;3(2):122-131. doi:10.1038/nrm732
228. Carta S, Tassi S, Semino C, et al. Histone deacetylase inhibitors prevent exocytosis of interleukin-1beta-containing secretory lysosomes: role of microtubules. *Blood*. 2006;108(5):1618-1626. doi:10.1182/blood-2006-03-014126
229. Rubartelli A, Cozzolinol F, Talio M, Sitia R. A novel secretory pathway for interleukin-1beta, a protein lacking a signal sequence. *EMBO J*. 1990;9(5):1503-1510.
230. Johnson DE, Ostrowski P, Jaumouillé V, Grinstein S. The position of lysosomes within the cell determines their luminal pH. *J Cell Biol*. 2016;212(6):677-692. doi:10.1083/jcb.201507112
231. Man SM, Hopkins LJ, Nugent E, et al. Inflammasome activation causes dual recruitment of NLRC4 and NLRP3 to the same macromolecular complex. *Proc Natl Acad Sci U S A*. 2014;111(1091-6490 (Electronic)).

232. Monteleone M, Stow JL, Schroder K. Mechanisms of unconventional secretion of IL-1 family cytokines. *Cytokine*. 2015;74(2):213-218. doi:10.1016/j.cyto.2015.03.022
233. MacKenzie A, Wilson HL, Kiss-Toth E, Dower SK, North RA, Surprenant A. Rapid Secretion of Interleukin-1b by Microvesicle Shedding. *Immunity*. 2001;8.
234. Pizzirani C, Ferrari D, Chiozzi P, et al. Stimulation of P2 receptors causes release of IL-1b-loaded microvesicles from human dendritic cells. *Blood*. 2007;109(9):3856-3864. doi:10.1182/blood-2005-06-
235. Gardella S, Andrei C, Ferrera D, et al. The nuclear protein HMGB1 is secreted by monocytes via a non-classical, vesicle-mediated secretory pathway. *EMBO Rep*. 2002;3(10):995-1001. doi:10.1093/embo-reports/kvf198
236. Brough D, Le Feuvre RA, Wheeler RD, et al. Ca²⁺ Stores and Ca²⁺ Entry Differentially Contribute to the Release of IL-1 and IL-1 from Murine Macrophages. *J Immunol*. 2003;170(6):3029-3036. doi:10.4049/jimmunol.170.6.3029
237. van Niel G, D'Angelo G, Raposo G. Shedding light on the cell biology of extracellular vesicles. *Nat Rev Mol Cell Biol*. 2018;19(4):213-228. doi:10.1038/nrm.2017.125
238. Qu Y, Ramachandra L, Mohr S, et al. P2X7 receptor-stimulated secretion of MHC class II-containing exosomes requires the ASC/NLRP3 inflammasome but is independent of caspase-1. *J Immunol*. 2009;182(8):5052-5062. doi:10.4049/jimmunol.0802968
239. Giuliani AL, Sarti AC, Falzoni S, Di Virgilio F. The P2X7 Receptor-Interleukin-1 Liaison. *Front Pharmacol*. 2017;8:123. doi:10.3389/fphar.2017.00123
240. Bhattacharya A, Prakash YS, Eissa NT. Secretory function of autophagy in innate immune cells. *Cell Microbiol*. 2014;16(11):1637-1645. doi:10.1111/cmi.12365
241. Dupont N, Jiang S, Pilli M, Ornatowski W, Bhattacharya D, Deretic V. Autophagy-based unconventional secretory pathway for extracellular delivery of IL-1beta. *EMBO J*. 2011;30(23):4701-4711. doi:10.1038/emboj.2011.398
242. Zhang M, Kenny SJ, Ge L, Xu K, Schekman R. Translocation of interleukin-1beta into a vesicle intermediate in autophagy-mediated secretion. *Elife*. 2015;4. doi:10.7554/eLife.11205
243. Carta S, Tassi S, Pettinati I, Delfino L, Dinarello CA, Rubartelli A. The rate of interleukin-1 β secretion in different myeloid cells varies with the extent of redox response to Toll-like receptor triggering. *J Biol Chem*. 2011;286(31):27069-27080. doi:10.1074/jbc.M110.203398
244. Bessen DE. Population biology of the human restricted pathogen, *Streptococcus pyogenes*. *Infect Genet Evol*. 2009;9(4):581-593. doi:10.1016/j.meegid.2009.03.002
245. Brown SP, Cornforth DM, Mideo N. Evolution of virulence in opportunistic pathogens: Generalism, plasticity, and control. *Trends Microbiol*. 2012;20(7):336-342. doi:10.1016/j.tim.2012.04.005
246. Carapetis JR, Steer AC, Mulholland EK, Weber M. The global burden of group A streptococcal diseases. *Lancet Infect Dis*. 2005;5(11):685-694. doi:10.1016/s1473-3099(05)70267-x
247. Barnett TC, Bowen AC, Carapetis JR. The fall and rise of Group A *Streptococcus* diseases. *Epidemiol Infect*. 2018;9-14. doi:10.1017/S0950268818002285

248. Cunningham MW. Pathogenesis of group A streptococcal infections. *Clin Microbiol Rev.* 2000;13(3):470-511. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC88944/pdf/cm000470.pdf>.
249. Snider LA, Swedo SE. PANDAS: current status and directions for research. *Mol Psychiatry.* 2004;9(10):900-907. doi:10.1038/sj.mp.4001542
250. Pinho-Ribeiro FA, Baddal B, Haarsma R, et al. Blocking Neuronal Signaling to Immune Cells Treats Streptococcal Invasive Infection. *Cell.* 2018;173(5):1083-1097.e22. doi:10.1016/J.CELL.2018.04.006
251. Wilkening R V, Federle MJ. Evolutionary Constraints Shaping Streptococcus pyogenes-Host Interactions. *Trends Microbiol.* 2017;25(7):562-572. doi:10.1016/j.tim.2017.01.007
252. Lamagni TL, Darenberg J, Luca-Harari B, et al. Epidemiology of severe Streptococcus pyogenes disease in Europe. *J Clin Microbiol.* 2008;46(7):2359-2367. doi:10.1128/JCM.00422-08
253. de Gier B, Vlamincxx BJM, Woudt SHS, van Sorge NM, van Asten L. Associations between common respiratory viruses and invasive group A streptococcal infection: A time-series analysis. *Influenza Other Respi Viruses.* 2019;(February):1-6. doi:10.1111/irv.12658
254. Lancefield R. A Serological Differentiation of Human and Other Groups of Hemolytic Streptococci. *J Exp Med.* 1933;57(4):571-595.
255. Walker MJ, Barnett TC, McArthur JD, et al. Disease manifestations and pathogenic mechanisms of Group A Streptococcus. *Clin Microbiol Rev.* 2014;27(2):264-301. doi:10.1128/CMR.00101-13
256. Dale JB, Batzloff MR, Cleary PP, Courtney HS, Good MF, Grandi G, Halperin S, Margarit IY, McNeil S, Pandey M, Smeesters PR, Steer AC . Current Approaches to Group A Streptococcal Vaccine Development. *Streptococcus pyogenes Basic Biol to Clin Manifestations.* 2016:993-1023.
257. Cunningham MW, McCormack JM, Fenderson PG, Ho MK, Beachey EH, Dale JB. Human and murine antibodies cross-reactive with streptococcal M protein and myosin recognize the sequence GLN-LYS-SER-LYS-GLN in M protein. *J Immunol.* 1989;143(8):2677-2683.
258. Nobbs AH, Lamont RJ, Jenkinson HF. Streptococcus adherence and colonization. *Microbiol Mol Biol Rev.* 2009;73(3):407-450, Table of Contents. doi:10.1128/MMBR.00014-09
259. Okada N, Liszewski MK, Atkinson JP, Caparon M. Membrane cofactor protein (CD46) is a keratinocyte receptor for the M protein of the group A streptococcus. *Proc Natl Acad Sci.* 2006;92(7):2489-2493. doi:10.1073/pnas.92.7.2489
260. Cywes C, Stamenkovic I, Wessels MR. CD44 as a receptor for colonization of the pharynx by group A streptococcus [In Process Citation]. *JClinInvest.* 2000;106(8):995-1002.
261. Brouwer S, Barnett TC, Rivera-Hernandez T, Rohde M, Walker MJ. Streptococcus pyogenes adhesion and colonization. *FEBS Lett.* 2016. doi:10.1002/1873-3468.12254
262. Siemens N, Chakrakodi B, Shambat SM, et al. Biofilm in group A streptococcal necrotizing soft tissue infections. *JCI Insight.* 2016;1(10). doi:10.1172/jci.insight.87882
263. Lapenta D, Rubens C, Chi E, Cleary PP. Group A Streptococci efficiently invade human respiratory epithelial cells. *Proc Natl Acad Sci U S A.* 1994;91(December):12115-12119.
264. Barnett TC, Liebl D, Seymour LM, et al. The globally disseminated MIT1 clone of group A

- Streptococcus evades autophagy for intracellular replication. *Cell Host Microbe*. 2013;14(6):675-682. doi:10.1016/j.chom.2013.11.003
265. Roberts AL, Connolly KL, Kirse DJ, et al. Detection of group A Streptococcus in tonsils from pediatric patients reveals high rate of asymptomatic streptococcal carriage. *BMC Pediatr*. 2012;12. doi:10.1186/1471-2431-12-3
266. Thulin P, Johansson L, Low DE, et al. Viable group A streptococci in macrophages during acute soft tissue infection. *PLoS Med*. 2006;3(3):371-379. doi:10.1371/journal.pmed.0030053
267. O'Neill AM, Thurston TL, Holden DW. Cytosolic Replication of Group A Streptococcus in Human Macrophages. *MBio*. 2016;7(2). doi:10.1128/mBio.00020-16
268. Herten E, Johansson L, Wallin R, et al. M1 protein-dependent intracellular trafficking promotes persistence and replication of Streptococcus pyogenes in macrophages. *J Innate Immun*. 2010;2(6):534-545. doi:000317635 [pii] 10.1159/000317635 [doi]
269. Bastiat-Sempe B, Love JF, Lomayeva N, Wessels MR, Mbio. Streptolysin O and NAD-glycohydrolase prevent phagolysosome acidification and promote group a streptococcus survival in macrophages. *MBio*. 2014;5(5):e01690-14. doi:10.1128/mBio.01690-14
270. Carlsson F, Berggård K, Stålhammar-Carlemalm M, Lindahl G. Evasion of Phagocytosis through Cooperation between Two Ligand-binding Regions in Streptococcus pyogenes M Protein. *J Exp Med*. 2003;198(7):1057-1068. doi:10.1084/jem.20030543
271. Dale JB, Washburn RG, Marques MB, Wessels MR. Hyaluronate capsule and surface M protein in resistance to opsonization of group A streptococci. *Infect Immun*. 1996;64(5):1495-1501.
272. Naegeli A, Bratanis E, Karlsson C, et al. Streptococcus pyogenes evades adaptive immunity through specific IgG glycan hydrolysis. *J Exp Med*. 2019;jem.20190293. doi:10.1084/jem.20190293
273. Edwards RJ, Taylor GW, Ferguson M, et al. Specific C-Terminal Cleavage and Inactivation of Interleukin-8 by Invasive Disease Isolates of Streptococcus pyogenes. *J Infect Dis*. 2005;192(5):783-790. doi:10.1086/432485
274. Goldblatt J, Lawrenson RA, Muir L, et al. A Requirement for Neutrophil Glycosaminoglycans in Chemokine:Receptor Interactions Is Revealed by the Streptococcal Protease SpyCEP. *J Immunol*. 2019;202. doi:10.4049/jimmunol.1801688
275. Flaherty RA, Puricelli JM, Higashi DL, Park CJ, Lee SW. Streptolysin S Promotes Programmed Cell Death and Enhances Inflammatory Signaling in Epithelial Keratinocytes during Group A Streptococcus Infection. *Infect Immun*. 2015;83(10):4118-4133. doi:10.1128/IAI.00611-15
276. Sierig G, Cywes C, Wessels MR, Ashbaugh CD. Cytotoxic effects of streptolysin o and streptolysin s enhance the virulence of poorly encapsulated group a streptococci. *Infect Immun*. 2003;71(1):446-455. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC143243/pdf/0789.pdf>.
277. Barnett TC, Cole JN, Rivera-Hernandez T, et al. Streptococcal toxins: role in pathogenesis and disease. *Cell Microbiol*. 2015;17(12):1721-1741. doi:10.1111/cmi.12531
278. Kasper KJ, Zeppa JJ, Wakabayashi AT, et al. Bacterial superantigens promote acute nasopharyngeal infection by Streptococcus pyogenes in a human MHC Class II-dependent manner. *PLoS Pathog*.

- 2014;10(5):e1004155. doi:10.1371/journal.ppat.1004155
279. Zeppa JJ, Kasper KJ, Mohorovic I, Mazzuca DM, Haeryfar SMM, McCormick JK. Nasopharyngeal infection by *Streptococcus pyogenes* requires superantigen-responsive V β -specific T cells. *Proc Natl Acad Sci*. 2017;201700858. doi:10.1073/pnas.1700858114
280. Gryllos I, Levin JC, Wessels MR. The CsrR/CsrS two-component system of group A *Streptococcus* responds to environmental Mg²⁺. *Proc Natl Acad Sci*. 2003;100(7):4227-4232. doi:10.1073/pnas.0636231100
281. Gryllos I, Tran-Winkler HJ, Cheng M-F, et al. Induction of group A *Streptococcus* virulence by a human antimicrobial peptide. *Proc Natl Acad Sci U S A*. 2008;105(1091-6490 (Electronic)). doi:10.1073/pnas.0803815105
282. Federle MJ, McIver KS, Scott JR. A response regulator that represses transcription of several virulence operons in the group A streptococcus. *J Bacteriol*. 1999;181(12):3649-3657.
283. Sumby P, Whitney AR, Graviss EA, DeLeo FR, Musser JM. Genome-wide analysis of group A streptococci reveals a mutation that modulates global phenotype and disease specificity. *PLoS Pathog*. 2006;2(1):0041-0049. doi:10.1371/journal.ppat.0020005
284. Engleberg NC, Heath A, Miller A, Rivera C, DiRita VJ. Spontaneous Mutations in the CsrRS Two-Component Regulatory System of *Streptococcus pyogenes* Result in Enhanced Virulence in a Murine Model of Skin and Soft Tissue Infection. *J Infect Dis*. 2002;183(7):1043-1054. doi:10.1086/319291
285. Hollands A, Pence MA, Timmer AM, et al. A Genetic Switch to Hypervirulence Reduces Colonization Phenotypes of the Globally Disseminated Group A *Streptococcus* MIT1 Clone. *J Infect Dis*. 2010;202(1):11-19. doi:10.1086/653124.A
286. Alam FM, Turner CE, Smith K, Wiles S, Sriskandan S. Inactivation of the CovR/S virulence regulator impairs infection in an improved murine model of *Streptococcus pyogenes* naso-pharyngeal infection. *PLoS One*. 2013;8(4):e61655. doi:10.1371/journal.pone.0061655
287. Wollein Waldetoft K, Raberg L. To harm or not to harm? On the evolution and expression of virulence in group A streptococci. *Trends Microbiol*. 2014;22(1):7-13. doi:10.1016/j.tim.2013.10.006
288. Nelson GE, Pondo T, Toews KA, et al. Epidemiology of invasive group A streptococcal infections in the United States, 2005-2012. *Clin Infect Dis*. 2016;63(4):478-486. doi:10.1093/cid/ciw248
289. Chatellier S, Ihendyane N, Kansal RG, et al. Genetic relatedness and superantigen expression in group A streptococcus serotype M1 isolates from patients with severe and nonsevere invasive diseases. *Infect Immun*. 2000;68(6):3523-3534. doi:10.1128/IAI.68.6.3523-3534.2000
290. Rogers S, Commons R, Danchin MH, et al. Strain Prevalence, Rather than Innate Virulence Potential, Is the Major Factor Responsible for an Increase in Serious Group A *Streptococcus* Infections. *J Infect Dis*. 2007;195(11):1625-1633. doi:10.1086/513875
291. Nasser W, Beres SB, Olsen RJ, et al. Evolutionary pathway to increased virulence and epidemic group A *Streptococcus* disease derived from 3,615 genome sequences. *Proc Natl Acad Sci U S A*. 2014;111(17):E1768-76. doi:10.1073/pnas.1403138111
292. Sumby P, Porcella SF, Madrigal AG, et al. Evolutionary origin and emergence of a highly successful

- clone of serotype M1 group A Streptococcus involved multiple horizontal gene transfer events. *J Infect Dis.* 2005;192(5):771-782. doi:10.1086/432514
293. Bradford Kline J, Collins CM. Analysis of the superantigenic activity of mutant and allelic forms of streptococcal pyrogenic exotoxin A. *Infect Immun.* 1996;64(3):861-869.
294. Zhu L, Olsen RJ, Nasser W, et al. A molecular trigger for intercontinental epidemics of group A Streptococcus. *J Clin Invest.* 2015;125(9):3545-3559. doi:10.1172/JCI82478
295. Zhu L, Olsen RJ, Nasser W, De La Riva Morales I, Mussera JM. Trading capsule for increased cytotoxin production: Contribution to virulence of a newly emerged clade of emm89 streptococcus pyogenes. *MBio.* 2015;6(5):1-10. doi:10.1128/mBio.01378-15
296. Zhu L, Olsen RJ, Lee JD, Porter AR, DeLeo FR, Musser JM. Contribution of Secreted NADase and Streptolysin O to the Pathogenesis of Epidemic Serotype M1 Streptococcus pyogenes Infections. *Am J Pathol.* 2017;187(3):605-613. doi:10.1016/j.ajpath.2016.11.003
297. Bricker AL, Carey VJ, Wessels MR. Role of NADase in virulence in experimental invasive group A streptococcal infection. *Infect Immun.* 2005;73(10):6562-6566. doi:10.1128/IAI.73.10.6562-6566.2005
298. Tatsuno I, Isaka M, Minami M, Hasegawa T. NADase as a target molecule of in vivo suppression of the toxicity in the invasive M-1 group A Streptococcal isolates. *BMC Microbiol.* 2010;10:144. doi:10.1186/1471-2180-10-144
299. Stevens DL, Salmi DB, McIndoo ER, Bryant AE. Molecular epidemiology of nga and NAD glycohydrolase/ADP-ribosyltransferase activity among Streptococcus pyogenes causing streptococcal toxic shock syndrome. *J Infect Dis.* 2000;182(4):1117-1128. doi:10.1086/315850 [pii] 10.1086/315850 [doi]
300. Tatsuno I, Sawai J, Okamoto A, et al. Characterization of the NAD-glycohydrolase in streptococcal strains. *Microbiology.* 2007;153(Pt 12):4253-4260. doi:10.1099/mic.0.2007/009555-0
301. Riddle DJ, Bessen DE, Caparon MG. Variation in Streptococcus pyogenes NAD+ glycohydrolase is associated with tissue tropism. *J Bacteriol.* 2010;192(14):3735-3746. doi:10.1128/jb.00234-10
302. Chandrasekaran S, Ghosh J, Port GC, Koh EI, Caparon MG. Analysis of polymorphic residues reveals distinct enzymatic and cytotoxic activities of the Streptococcus pyogenes NAD+ glycohydrolase. *J Biol Chem.* 2013;288(27):20064-20075.
303. Kimoto H, Fujii Y, Yokota Y, Taketo A. Molecular characterization of NADase-streptolysin O operon of hemolytic streptococci. *Biochim Biophys Acta.* 2005;1681(2-3):134-149. doi:10.1016/j.bbaexp.2004.10.011
304. Madden JC, Ruiz N, Caparon M. Cytolysin-mediated translocation (CMT): a functional equivalent of type III secretion in gram-positive bacteria. *Cell.* 2001;104(1):143-152.
305. Meehl MA, Pinkner JS, Anderson PJ, Hultgren SJ, Caparon MG. A novel endogenous inhibitor of the secreted streptococcal NAD-glycohydrolase. *PLoS Pathog.* 2005;1(4):e35. doi:10.1371/journal.ppat.0010035
306. Savic DJ, McShan WM, Ferretti JJ. Autonomous expression of the slo gene of the bicistronic nga-slo operon of Streptococcus pyogenes. *Infect Immun.* 2002;70(5):2730-2733.
307. Velarde JJ, O'Seaghdha M, Baddal B, Bastiat-Sempe B, Wessels MR. Binding of NAD(+)-

- Glycohydrolase to Streptolysin O Stabilizes Both Toxins and Promotes Virulence of Group A Streptococcus. *MBio*. 2017;8(5). doi:10.1128/mBio.01382-17
308. Michos A, Gryllos I, Hakansson A, Srivastava A, Kokkotou E, Wessels MR. Enhancement of streptolysin O activity and intrinsic cytotoxic effects of the group A streptococcal toxin, NAD-glycohydrolase. *J Biol Chem*. 2006;281(12):8216-8223. doi:10.1074/jbc.M511674200
309. Tweten RK. Cholesterol-dependent cytolysins, a family of versatile pore-forming toxins. *Infect Immun*. 2005;73(10):6199-6209. doi:10.1128/IAI.73.10.6199-6209.2005
310. Giddings KS, Johnson AE, Tweten RK. Redefining cholesterol's role in the mechanism of the cholesterol-dependent cytolysins. *Proc Natl Acad Sci*. 2003;100(20):11315-11320. doi:10.1073/pnas.2033520100
311. Shewell LK, Harvey RM, Higgins MA, et al. The cholesterol-dependent cytolysins pneumolysin and streptolysin O require binding to red blood cell glycans for hemolytic activity. *Proc Natl Acad Sci*. 2014;111(49):E5312-E5320. doi:10.1073/pnas.1412703111
312. Mozola CC, Caparon MG. Dual modes of membrane binding direct pore formation by Streptolysin O. *Mol Microbiol*. 2015. doi:10.1111/mmi.13085
313. Mozola CC, Magassa N, Caparon MG. A novel cholesterol-insensitive mode of membrane binding promotes cytolysin-mediated translocation by Streptolysin O. *Mol Microbiol*. 2014;94(3):675-687. doi:10.1111/mmi.12786
314. Cywes Bentley C, Hakansson A, Christianson J, Wessels MR. Extracellular group A Streptococcus induces keratinocyte apoptosis by dysregulating calcium signalling. *Cell Microbiol*. 2005;7(7):945-955. doi:10.1111/j.1462-5822.2005.00525.x
315. Timmer AM, Timmer JC, Pence MA, et al. Streptolysin O promotes group A Streptococcus immune evasion by accelerated macrophage apoptosis. *J Biol Chem*. 2009;284(2):862-871. doi:10.1074/jbc.M804632200
316. Goldmann O, Sastalla I, Wos-Oxley M, Rohde M, Medina E. Streptococcus pyogenes induces oncosis in macrophages through the activation of an inflammatory programmed cell death pathway. *Cell Microbiol*. 2009;11(1462-5822 (Electronic)):138-155.
317. Limbago B, Penumalli V, Weinrick B, Scott JR. Role of streptolysin O in a mouse model of invasive group A streptococcal disease. *Infect Immun*. 2000;68(11):6384-6390. doi:10.1128/IAI.68.11.6384-6390.2000
318. Bhattacharjee P, Keyel PA. Cholesterol-dependent cytolysins impair pro-inflammatory macrophage responses. *Sci Rep*. 2018;79430. doi:10.1038/s41598-018-24955-2
319. Ruiz N, Wang B, Pentland A, Caparon M. Streptolysin O and adherence synergistically modulate proinflammatory responses of keratinocytes to group A streptococci. *Mol Microbiol*. 1998;27(2):337-346.
320. Hakansson A, Bentley CC, Shakhnovic EA, Wessels MR. Cytolysin-dependent evasion of lysosomal killing. *Proc Natl Acad Sci U S A*. 2005;102(14):5192-5197. doi:10.1073/pnas.0408721102
321. Uchiyama S, Dohrmann S, Timmer AM, et al. Streptolysin O Rapidly Impairs Neutrophil Oxidative Burst

- and Antibacterial Responses to Group A Streptococcus. *Front Immunol.* 2015;6:581. doi:10.3389/fimmu.2015.00581
322. Meehl MA, Caparon MG. Specificity of streptolysin O in cytolysin-mediated translocation. *Mol Microbiol.* 2004;52(6):1665-1676. doi:10.1111/j.1365-2958.2004.04082.x
 323. Ghosh J, Caparon MG. Specificity of Streptococcus pyogenes NAD(+) glycohydrolase in cytolysin-mediated translocation. *Mol Microbiol.* 2006;62(4):1203-1214. doi:10.1111/j.1365-2958.2006.05430.x
 324. Magassa N, Chandrasekaran S, Caparon MG. Streptococcus pyogenes cytolysin-mediated translocation does not require pore formation by streptolysin O. *EMBO Rep.* 2010;(1469-3178 (Electronic)). doi:10.1038/embor.2010.37
 325. Carlson BYAS, Kellner A, Bernheimer, Alan W Ph D, Freeman EB. A Streptococcal enzyme that acts specifically upon diphosphopyridine nucleotide: characterization of the enzyme and its separation from Streptolysin O. *J Exp Med.* 1957;106(1):15-26.
 326. Smith CL, Ghosh J, Elam JS, et al. Structural basis of Streptococcus pyogenes immunity to its NAD+ glycohydrolase toxin. *Structure.* 2011;19(2):192-202. doi:10.1016/j.str.2010.12.013
 327. Kimoto H, Fujii Y, Hirano S, Yokota Y, Taketo A. Genetic and biochemical properties of streptococcal NAD-glycohydrolase inhibitor. *J Biol Chem.* 2006;281(14):9181-9189. doi:10.1074/jbc.M506879200
 328. Karasawa T, Takasawa S, Yamakawa K, Yonekura H, Okamoto H, Nakamura S. NAD(+)-glycohydrolase from Streptococcus pyogenes shows cyclic ADP-ribose forming activity. *FEMS Microbiol Lett.* 1995;130(2-3):201-204.
 329. Ghosh J, Anderson PJ, Chandrasekaran S, Caparon MG. Characterization of Streptococcus pyogenes beta-NAD+ glycohydrolase: re-evaluation of enzymatic properties associated with pathogenesis. *J Biol Chem.* 2010;285(8):5683-5694. doi:10.1074/jbc.M109.070300
 330. Grange PA, Raingeaud J, Calvez V, Dupin N. Nicotinamide inhibits Propionibacterium acnes-induced IL-8 production in keratinocytes through the NF-kappaB and MAPK pathways. *J Dermatol Sci.* 2009;56(2):106-112. doi:10.1016/j.jdermsci.2009.08.001
 331. Ungerstedt JS, Blomback M, Soderstrom T. Nicotinamide is a potent inhibitor of proinflammatory cytokines. *Clin Exp Immunol.* 2003;131(1):48-52. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1808598/pdf/cei0131-0048.pdf>.
 332. Putney JW. Calcium signaling. *Calcium Signaling, Second Ed.* 2005:1-540. doi:10.1201/9781420038231
 333. Gasser A, Glassmeier G, Fliegert R, et al. Activation of T cell calcium influx by the second messenger ADP-ribose. *J Biol Chem.* 2006;281(5):2489-2496. doi:10.1074/jbc.M506525200
 334. Bricker AL, Cywes C, Ashbaugh CD, Wessels MR. NAD+-glycohydrolase acts as an intracellular toxin to enhance the extracellular survival of group A streptococci. *Mol Microbiol.* 2002;44(1):257-269.
 335. Chandrasekaran S, Caparon MG. The Streptococcus pyogenes NAD glycohydrolase modulates epithelial cell PARylation and HMGB1 release. *Cell Microbiol.* 2015. doi:10.1111/cmi.12442
 336. Chandrasekaran S, Caparon MG. The NADase-Negative Variant of the Streptococcus pyogenes Toxin NAD(+) Glycohydrolase Induces JNK1-Mediated Programmed Cellular Necrosis. *MBio.* 2016;7(1):e02215-15. doi:10.1128/mBio.02215-15

337. Sharma O, O'Seaghda M, Velarde JJ, Wessels MR. NAD⁺-Glycohydrolase Promotes Intracellular Survival of Group A Streptococcus. *PLoS Pathog.* 2016;12(3):e1005468. doi:10.1371/journal.ppat.1005468
338. Gibson BA, Kraus WL, Nat Rev Mol Cell B. New insights into the molecular and cellular functions of poly(ADP-ribose) and PARPs. (1471-0080 (Electronic)).
339. Kim S, Kim SY. Signaling of High Mobility Group Box 1 (HMGB1) through Toll-like Receptor 4 in Macrophages Requires CD14. *Mol Med.* 2013;19(1):1. doi:10.2119/molmed.2012.00306
340. O'Seaghda M, Wessels MR. Streptolysin O and its co-toxin NAD-glycohydrolase protect group A Streptococcus from Xenophagic killing. *PLoS Pathog.* 2013;9(6):e1003394. doi:10.1371/journal.ppat.1003394
341. Ratner AJ, Hippe KR, Aguilar JL, Bender MH, Nelson AL, Weiser JN. Epithelial cells are sensitive detectors of bacterial pore-forming toxins. *J Biol Chem.* 2006;281(18):12994-12998. doi:10.1074/jbc.M511431200
342. Kloft N, Busch T, Neukirch C, et al. Pore-forming toxins activate MAPK p38 by causing loss of cellular potassium. *Biochem Biophys Res Commun.* 2009;385(4):503-506. doi:10.1016/j.bbrc.2009.05.121
343. Navarini AA, Lang KS, Verschoor A, et al. Innate immune-induced depletion of bone marrow neutrophils aggravates systemic bacterial infections. *Proc Natl Acad Sci.* 2009;106(17):7107-7112. doi:10.1073/pnas.0901162106
344. Goldmann O, Rohde M, Chhatwal GS, Medina E. Role of Macrophages in Host Resistance to Group A Streptococci. *Infect Immun.* 2004;72(5):2956-2963. doi:10.1128/iai.72.5.2956-2963.2004
345. Loof TG, Rohde M, Chhatwal GS, Jung S, Medina E. The Contribution of Dendritic Cells to Host Defenses against Streptococcus pyogenes. *J Infect Dis.* 2007;196(12):1794-1803. doi:10.1086/523647
346. Dileepan T, Linehan JL, Moon JJ, Pepper M, Jenkins MK, Cleary PP. Robust antigen specific th17 T cell response to group A Streptococcus is dependent on IL-6 and intranasal route of infection. *PLoS Pathog.* 2011;7(9):e1002252. doi:10.1371/journal.ppat.1002252
347. Mishalian I, Ordan M, Peled A, et al. Recruited macrophages control dissemination of group A Streptococcus from infected soft tissues. *J Immunol.* 2011;187(11):6022-6031. doi:10.4049/jimmunol.1101385
348. Metzger DW, Raeder R, Cleave VH Van, Boyle MDP. Protection of Mice from Group A Streptococcal Skin Infection by Interleukin-12. *J Infect Dis.* 1995;171:1643-1645. doi:10.1093/infdis/171.6.1643
349. Gratz N, Hartweger H, Matt U, et al. Type I Interferon Production Induced By Streptococcus Pyogenes-Derived Nucleic Acids is Required for Host Protection. *PLoS Pathog.* 2011;7(5):1-16. doi:10.1371/journal.ppat.1001345
350. Castiglia V, Piersigilli A, Ebner F, et al. Type I Interferon Signaling Prevents IL-1 β -Driven Lethal Systemic Hyperinflammation during Invasive Bacterial Infection of Soft Tissue. *Cell Host Microbe.* 2016;19(3):375-387. doi:10.1016/j.chom.2016.02.003
351. Gratz N, Siller M, Schaljo B, et al. Group A streptococcus activates type I interferon production and MyD88-dependent signaling without involvement of TLR2, TLR4, and TLR9. *J Biol Chem.*

- 2008;283(29):19879-19887. doi:10.1074/jbc.M802848200
352. Loof TG, Goldmann O, Gessner A, Herwald H, Medina E. Aberrant inflammatory response to *Streptococcus pyogenes* in mice lacking myeloid differentiation factor 88. *Am J Pathol.* 2010;176(1525-2191 (Electronic)). doi:10.2353/ajpath.2010.090422
353. Loof TG, Goldmann O, Medina E. Immune recognition of *Streptococcus pyogenes* by dendritic cells. *Infect Immun.* 2008;76(6):2785-2792. doi:10.1128/IAI.01680-07
354. Von Bernuth H, Picard C, Jin Z, et al. Pyogenic bacterial infections in humans with MyD88 deficiency. *Science (80-).* 2008;321(5889):691-696. doi:10.1126/science.1158298
355. Park JM, Ng VH, Maeda S, Rest RF, Karin M. Anthrolysin O and Other Gram-positive Cytolysins Are Toll-like Receptor 4 Agonists. *J Exp Med.* 2004;200(12):1647-1655. doi:10.1084/jem.20041215
356. Wang X, Fan X, Bi S, Li N, Wang B. Toll-like Receptors 2 and 4-Mediated Reciprocal Th17 and Antibody Responses to Group A *Streptococcus* Infection. *J Infect Dis.* 2017;215(4):644-652. doi:10.1093/infdis/jiw598
357. Liadaki K, Petinaki E, Skoulakis C, et al. Toll-like receptor 4 gene (TLR4), but not TLR2, polymorphisms modify the risk of tonsillar disease due to *Streptococcus pyogenes* and *Haemophilus influenzae*. *Clin Vaccine Immunol.* 2011;18(2):217-222. doi:10.1128/CVI.00460-10
358. Fieber C, Janos M, Koestler T, et al. Innate immune response to streptococcus pyogenes depends on the combined activation of TLR13 and TLR2. *PLoS One.* 2015;10(3):1-20. doi:10.1371/journal.pone.0119727
359. Hafner A, Kolbe U, Freund I, et al. Crucial Role of Nucleic Acid Sensing via Endosomal Toll-Like Receptors for the Defense of *Streptococcus pyogenes* in vitro and in vivo. *Front Immunol.* 2019;10(FEB):1-13. doi:10.3389/fimmu.2019.00198
360. Moverf E, Lienard J, Valfridsson C, Nordström T, Johansson-Lindbom B, Carlsson F. Streptococcal M protein promotes IL-10 production by cGAS-independent activation of the STING signaling pathway. *PLoS Pathog.* 2018;14(3). doi:10.1371/journal.ppat.1006969
361. Goldmann O, Lengeling A, Böse J, et al. The Role of the MHC on Resistance to Group A Streptococci in Mice. *J Immunol.* 2005;175(6):3862-3872. doi:10.4049/jimmunol.175.6.3862
362. Medina E, Goldmann O, Rohde M, Lengeling A, Chhatwals GS. Genetic Control of Susceptibility to Group A Streptococcal Infection in Mice. *J Infect Dis.* 2001;184(7):846-852. doi:10.1086/323292
363. Bessen D, Fischetti VA. PASSIVE ACQUIRED MUCOSAL IMMUNITY TO GROUP A STREPTOCOCCI BY SECRETORY IMMUNOGLOBULIN A. *J Exp Med.* 1988;167(June).
364. Shaikh N, Leonard E, Martin JM. Prevalence of Streptococcal Pharyngitis and Streptococcal Carriage in Children: A Meta-analysis. *Pediatrics.* 2010;126(3):e557-e564. doi:10.1542/peds.2009-2648
365. Mortensen R, Nissen TN, Blauenfeldt T, Christensen JP, Andersen P, Dietrich J. Adaptive Immunity against *Streptococcus pyogenes* in Adults Involves Increased IFN- γ and IgG3 Responses Compared with Children. *J Immunol.* 2015;195(4):1657-1664. doi:10.4049/jimmunol.1500804
366. Costalonga M, Cleary PP, Fischer LA, Zhao Z. Intranasal bacteria induce Th1 but not Treg or Th2. *Mucosal Immunol.* 2009;2(1):85-95. doi:10.1038/mi.2008.67

367. Wang B, Dileepan T, Briscoe S, et al. Induction of TGF-beta1 and TGF-beta1-dependent predominant Th17 differentiation by group A streptococcal infection. *Proc Natl Acad Sci U S A*. 2010;107(13):5937-5942. doi:10.1073/pnas.0904831107
368. Linehan JL, Dileepan T, Kashem SW, Kaplan DH, Cleary P, Jenkins MK. Generation of Th17 cells in response to intranasal infection requires TGF-beta1 from dendritic cells and IL-6 from CD301b+ dendritic cells. *Proc Natl Acad Sci U S A*. 2015;112(41):12782-12787. doi:10.1073/pnas.1513532112
369. Dileepan T, Smith ED, Knowland D, et al. Group A Streptococcus intranasal infection promotes CNS infiltration by streptococcal-specific Th17 cells. *J Clin Invest*. 2016;126(1):303-317. doi:10.1172/JCI80792
370. Chella Krishnan K, Mukundan S, Alagarsamy J, Laturnus D, Kotb M. Host Genetic Variations and Sex Differences Potentiate Predisposition, Severity, and Outcomes of Group A Streptococcus-Mediated Necrotizing Soft Tissue Infections. *Infect Immun*. 2016;84(2):416-424. doi:10.1128/IAI.01191-15
371. Kotb M, Norrby-Teglund A, McGeer A, et al. An immunogenetic and molecular basis for differences in outcomes of invasive group A streptococcal infections. *Nat Med*. 2002;8(12):1398-1404. doi:10.1038/nm800
372. Norrby-Teglund A, Chatellier S, Low DE, McGeer A, Green K, Kotb M. Host variation in cytokine responses to superantigens determine the severity of invasive group A streptococcal infection. *Eur J Immunol*. 2000;30(11):3247-3255.
373. Nooh MM, Nookala S, Kansal R, Kotb M. Individual Genetic Variations Directly Effect Polarization of Cytokine Responses to Superantigens Associated with Streptococcal Sepsis: Implications for Customized Patient Care. *J Immunol*. 2011;186(5):3156-3163. doi:10.4049/jimmunol.1002057
374. Nooh MM, El-Gengehi N, Kansal R, David CS, Kotb M. HLA Transgenic Mice Provide Evidence for a Direct and Dominant Role of HLA Class II Variation in Modulating the Severity of Streptococcal Sepsis. *J Immunol*. 2007;178(5):3076-3083. doi:10.4049/jimmunol.178.5.3076
375. Goldmann O, Chhatwal GS, Medina E. Immune mechanisms underlying host susceptibility to infection with group A streptococci. *J Infect Dis*. 2003;187(5):854-861. doi:10.1086/368390
376. Chella Krishnan K, Mukundan S, Alagarsamy J, et al. Genetic Architecture of Group A Streptococcal Necrotizing Soft Tissue Infections in the Mouse. *PLoS Pathog*. 2016;12(7):e1005732. doi:10.1371/journal.ppat.1005732
377. Hsu LC, Enzler T, Seita J, et al. IL-1 β -driven neutrophilia preserves antibacterial defense in the absence of the kinase IKK β . *Nat Immunol*. 2011;12(2):144-150. doi:10.1038/ni.1976
378. Flaherty RA, Donahue DL, Carothers KE, et al. Neutralization of Streptolysin S-Dependent and Independent Inflammatory Cytokine IL-1 β Activity Reduces Pathology During Early Group A Streptococcal Skin Infection. *Front Cell Infect Microbiol*. 2018;8(July):1-14. doi:10.3389/fcimb.2018.00211
379. Lin AE, Beasley FC, Keller N, et al. A group A Streptococcus ADP-ribosyltransferase toxin stimulates a protective interleukin 1beta-dependent macrophage immune response. *MBio*. 2015;6(2):e00133. doi:10.1128/mBio.00133-15
380. Valderrama JA, Riestra AM, Gao NJ, et al. Group A streptococcal M protein activates the NLRP3

- inflammasome. *Nat Microbiol.* 2017;2(10):1425-1434. doi:10.1038/s41564-017-0005-6
381. Hancz D, Westerlund E, Valfridsson C, et al. Streptolysin O induces ubiquitination and degradation of pro-IL-1 β . *J Innate Immun.* 2019;in press. doi:10.1159/000496403
382. Keyel PA, Roth R, Yokoyama WM, Heuser JE, Salter RD. Reduction of streptolysin O (SLO) pore-forming activity enhances inflammasome activation. *Toxins (Basel).* 2013;5(6):1105-1118. doi:10.3390/toxins5061105
383. Hancz D, Westerlund E, Bastiat-Sempe B, et al. Inhibition of Inflammasome-Dependent Interleukin 1beta Production by Streptococcal NAD⁺-Glycohydrolase: Evidence for Extracellular Activity. *MBio.* 2017;8(4). doi:10.1128/mBio.00756-17
384. Westerlund E, Valfridsson C, Yi DX, Persson JJ. The Secreted Virulence Factor NADase of Group A Streptococcus Inhibits P2X7 Receptor-Mediated Release of IL-1 β . *Front Immunol.* 2019;10(June). doi:10.3389/fimmu.2019.01385
385. Sun J, Siroy A, Lokareddy RK, et al. The tuberculosis necrotizing toxin kills macrophages by hydrolyzing NAD. *Nat Struct Mol Biol.* 2015;22(9):672-678. doi:10.1038/nsmb.3064
386. Tak U, Vlach J, Garza-Garcia A, et al. The tuberculosis necrotizing toxin is an NAD⁺ and NADP⁺ glycohydrolase with distinct enzymatic properties. *J Biol Chem.* 2018;294(9):3024-3036. doi:10.1074/jbc.ra118.005832
387. de Torre-Minguela C, Barbera-Cremades M, Gomez AI, Martin-Sanchez F, Pelegrin P. Macrophage activation and polarization modify P2X7 receptor secretome influencing the inflammatory process. *Sci Rep.* 2016;6:22586. doi:10.1038/srep22586
388. Gu BJ, Zhang W, Worthington RA, et al. A Glu-496 to Ala Polymorphism Leads to Loss of Function of the Human P2X7 Receptor. *J Biol Chem.* 2001;276(14):11135-11142. doi:10.1074/jbc.M010353200
389. Sokolovska A, Becker CE, Ip WKE, et al. Activation of caspase-1 by the NLRP3 inflammasome regulates the NADPH oxidase NOX2 to control phagosome function. *Nat Immunol.* 2013;14(6):543-553. doi:10.1038/ni.2595