



LUND UNIVERSITY

Convergence analysis of the nonoverlapping Robin-Robin method for nonlinear elliptic equations

Engström, Emil; Hansen, Eskil

Published in:
SIAM Journal on Numerical Analysis

DOI:
[10.1137/21M1414942](https://doi.org/10.1137/21M1414942)

2022

Document Version:
Early version, also known as pre-print

[Link to publication](#)

Citation for published version (APA):
Engström, E., & Hansen, E. (2022). Convergence analysis of the nonoverlapping Robin-Robin method for nonlinear elliptic equations. *SIAM Journal on Numerical Analysis*, 60(2), 585-605.
<https://doi.org/10.1137/21M1414942>

Total number of authors:
2

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

CONVERGENCE ANALYSIS OF THE NONOVERLAPPING ROBIN–ROBIN METHOD FOR NONLINEAR ELLIPTIC EQUATIONS*

EMIL ENGSTRÖM[†] AND ESKIL HANSEN[‡]

Abstract. We prove convergence for the nonoverlapping Robin–Robin method applied to nonlinear elliptic equations with a p -structure, including degenerate diffusion equations governed by the p -Laplacian. This nonoverlapping domain decomposition is commonly encountered when discretizing elliptic equations, as it enables the usage of parallel and distributed hardware. Convergence has been derived in various linear contexts, but little has been proven for nonlinear equations. Hence, we develop a new theory for nonlinear Steklov–Poincaré operators based on the p -structure and the L^p -generalization of the Lions–Magenes spaces. This framework allows the reformulation of the Robin–Robin method into a Peaceman–Rachford splitting on the interfaces of the subdomains, and the convergence analysis then follows by employing elements of the abstract theory for monotone operators. The analysis is performed on Lipschitz domains and without restrictive regularity assumptions on the solutions.

Key words. Robin–Robin method, Nonoverlapping domain decomposition, Nonlinear elliptic equation, Convergence, Steklov–Poincaré operator

AMS subject classifications. 65N55, 65J15, 35J70, 47N20

1. Introduction. Approximating the solution of an elliptic partial differential equation (PDE) typically demands large-scale computations requiring the usage of parallel and distributed hardware. In this context, a nonoverlapping domain decomposition method is a suitable choice, as it can be implemented in parallel with local communication. After decomposing the equation’s spatial domain into nonoverlapping subdomains, the method consists of an iterative procedure that solves the equation on each subdomain and thereafter communicates the results via the boundaries to the adjacent subdomains. For a general introduction we refer to [18, 23].

There is a vast amount of methods in the literature, employing different transmission conditions between the subdomains. The standard examples are based on the alternate use of Dirichlet and Neumann boundary conditions, but a competitive alternative is the Robin–Robin method, where the same type of Robin boundary condition is used for all subdomains. The Robin–Robin method was introduced in [13] together with a convergence proof when applied to linear elliptic equations. After applying a finite element discretization, convergence rates of the form $1 - \mathcal{O}(\sqrt{h})$, with h denoting the mesh width, have been derived in various linear contexts [9, 15, 24]; also see [7]. For generalizations and further applications of the Robin–Robin method applied to linear PDEs we refer to [3, 9] and references therein.

When considering nonlinear elliptic PDEs the literature is more limited. Convergence studies relating to overlapping Schwarz methods are given in [6, 21, 22]. However, there are hardly any results dealing with nonoverlapping domain decomposition schemes. One exception is [2], where the convergence of the Dirichlet–Neumann and Robin–Robin methods are analyzed for a family of one-dimensional elliptic equations.

*Version of April 26, 2021.

Funding: This work was supported by the Swedish Research Council under the grant 2019–05396.

[†]Centre for Mathematical Sciences, Lund University, P.O. Box 118, SE-22100 Lund, Sweden, (emil.engstrom@math.lth.se).

[‡]Centre for Mathematical Sciences, Lund University, P.O. Box 118, SE-22100 Lund, Sweden, (eskil.hansen@math.lth.se).

A related study is [20], where the equivalence between a class of nonlinear elliptic equations and the corresponding transmission problems are proven for nonoverlapping decompositions with cross points, but no numerical scheme is considered. Apart from [22], all these nonlinear studies rely on frameworks similar to the linear case, e.g., assuming that the diffusion is uniformly positive.

Hence, the aim of this paper is to derive a genuinely nonlinear extension of the linear convergence result given in [13] for the nonoverlapping Robin–Robin method. We will focus on nonlinear elliptic equations of the form

$$(1.1) \quad \begin{cases} -\nabla \cdot \alpha(\nabla u) + g(u) = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded domain in $\mathbb{R}^d$, $d = 1, 2, \dots$, with boundary $\partial\Omega$. The functions α and g are assumed to have a p -structure; defined in section 2. This structure enables a clear-cut convergence analysis for a broad family of degenerate elliptic equations, i.e., $\alpha(\nabla u)$ may vanish for nonzero values of u . The latter typically prevents the existence of a strong solution in $W^{2,p}(\Omega)$.

The archetypical examples of nonlinear elliptic equations with a p -structure are those governed by the p -Laplacian, where $\alpha(z) = |z|^{p-2}z$. Examples include the computation of the nonlinear resolvent

$$(1.2) \quad -\nabla \cdot (|\nabla u|^{p-2} \nabla u) + \lambda u = f,$$

arising in the context of an implicit Euler discretization of the parabolic p -Laplace equation, and the nonlinear reaction-diffusion problem

$$(1.3) \quad -\nabla \cdot (|\nabla u|^{p-2} \nabla u) + \lambda |u|^{p-2} u = f.$$

For sake of simplicity, we decompose the original domain Ω into two nonoverlapping subdomains $\{\Omega_i, \text{ for } i = 1, 2\}$, with boundaries denoted by $\partial\Omega_i$, and separated by the interface Γ , i.e.,

$$\bar{\Omega} = \bar{\Omega}_1 \cup \bar{\Omega}_2, \quad \Omega_1 \cap \Omega_2 = \emptyset \quad \text{and} \quad \Gamma = (\partial\Omega_1 \cap \partial\Omega_2) \setminus \partial\Omega.$$

Two examples of such decompositions are illustrated in Figures 1a and 1b, respectively. The analysis presented here can also, in a trivial fashion, be extended to the case when Ω_i is a family of nonadjacent subdomains, e.g., the stripwise domain decomposition illustrated in Figure 1c. The strong form of the Robin–Robin method applied to (1.1) is then to find (u_1^n, u_2^n) , for $n = 1, 2, \dots$, such that

$$(1.4) \quad \begin{cases} -\nabla \cdot \alpha(\nabla u_1^{n+1}) + g(u_1^{n+1}) = f & \text{in } \Omega_1, \\ u_1^{n+1} = 0 & \text{on } \partial\Omega_1 \setminus \Gamma, \\ \alpha(\nabla u_1^{n+1}) \cdot \nu_1 + s u_1^{n+1} = \alpha(\nabla u_2^n) \cdot \nu_1 + s u_2^n & \text{on } \Gamma, \\ -\nabla \cdot \alpha(\nabla u_2^{n+1}) + g(u_2^{n+1}) = f & \text{in } \Omega_2, \\ u_2^{n+1} = 0 & \text{on } \partial\Omega_2 \setminus \Gamma, \\ \alpha(\nabla u_2^{n+1}) \cdot \nu_2 + s u_2^{n+1} = \alpha(\nabla u_1^{n+1}) \cdot \nu_2 + s u_1^{n+1} & \text{on } \Gamma, \end{cases}$$

where ν_i denotes the unit outward normal vector of $\partial\Omega_i$, u_2^0 is a given initial guess and $s > 0$ is a method parameter. Here, u_i^n and $u_i^n|_\Gamma$ approximate the solution u restricted to Ω_i and Γ , respectively.




Fig. 1: Examples of different domain decompositions: (a) illustrates a domain decomposition with two intersection points; (b) a decomposition without intersection points; (c) a stripwise decomposition without cross points; (d) a full decomposition with cross points.

The convergence analysis is organized as follows. For linear elliptic equations, i.e., equations with a 2-structure, the analysis relies on the trace operator from $H^1(\Omega_i)$ onto $H^{1/2}(\partial\Omega_i)$, and the Lions–Magenes spaces $H_{00}^{1/2}(\Gamma)$. We therefore start by introducing the generalized p -version of the trace operator, now given from $W^{1,p}(\Omega_i)$ onto $W^{1-1/p,p}(\partial\Omega_i)$, and the corresponding Lions–Magenes spaces Λ_i ; see [sections 3](#) and [4](#). There is a surprising lack of proofs in the literature dealing with this generalized p -setting. We therefore make an effort to give precise definitions and proof references.

With the correct function spaces in place, we prove that the nonlinear transmission problem is equivalent to the weak form of [\(1.1\)](#) in [Theorem 5.2](#), and introduce the new nonlinear Steklov–Poincaré operators, as maps from Λ_i to Λ_i^* , in [section 6](#). The latter yields that the transmission problem can be stated as a problem on Γ and the Robin–Robin method reduces to the Peaceman–Rachford splitting. The main challenge is then to derive the fundamental properties of the nonlinear Steklov–Poincaré operators from the p -structure, which is achieved in [section 7](#). By interpreting the nonlinear Steklov–Poincaré operators as unbounded, monotone maps on $L^2(\Gamma)$ we finally prove that the Robin–Robin method is well defined on $W^{1,p}(\Omega_1) \times W^{1,p}(\Omega_2)$; see [Corollary 8.6](#), and convergent in the same space; see [Theorem 8.9](#). The latter relies on the abstract theory of the Peaceman–Rachford splittings [\[14\]](#).

The continuous convergence analysis presented here also holds in the finite dimensional case obtained after a suitable spatial discretization, e.g., by employing finite elements. However, we will limit ourselves to the continuous case in this paper. Hence, important issues including convergence rates for the finite-dimensional case and the influence of the mesh width on the optimal choice of the method parameter s will be explored elsewhere.

Finally, c_i and C_i will denote generic positive constants that assume different values at different occurrences.

2. Nonlinear elliptic equations with p -structure. Throughout the paper, we will consider the nonlinear elliptic equation (1.1) with $f \in L^2(\Omega)$ and Ω being a bounded Lipschitz domain. The equation is assumed to have a p -structure of the form:

Assumption 2.1. The parameters (p, r) and the functions $\alpha : \mathbb{R}^d \rightarrow \mathbb{R}^d, g : \mathbb{R} \rightarrow \mathbb{R}$ satisfy the properties below.

- Let $p \in [2, \infty)$ and $r \in (1, \infty)$. If $p < d$ then $r \leq dp/(2(d-p)) + 1$.
- The functions α and g are continuous and satisfy the growth conditions

$$|\alpha(z)| \leq C_1 |z|^{p-1} \quad \text{and} \quad |g(x)| \leq C_2 |x|^{r-1}, \quad \text{for all } z \in \mathbb{R}^d, x \in \mathbb{R}.$$

- The function α is strictly monotone with the bound

$$(\alpha(z) - \alpha(\tilde{z})) \cdot (z - \tilde{z}) \geq c_1 |z - \tilde{z}|^p, \quad \text{for all } z, \tilde{z} \in \mathbb{R}^d.$$

- The function α is coercive with the bound

$$\alpha(z) \cdot z \geq c_2 |z|^p, \quad \text{for all } z \in \mathbb{R}^d.$$

- The function g is strictly monotone and coercive with the bounds

$$(g(x) - g(\tilde{x}))(x - \tilde{x}) \geq c_3 |x - \tilde{x}|^r \quad \text{and} \quad g(x)x \geq c_4 |x|^r, \quad \text{for all } x, \tilde{x} \in \mathbb{R}.$$

Example 2.2. The equation (1.2) satisfies Assumption 2.1 with $\alpha(z) = |z|^{p-2}z$, $\lambda > 0$, $g(x) = \lambda x$ and $r=2$. The same holds for equation (1.3) with $g(x) = \lambda |x|^{p-2}x$ and $r = p$.

Remark 2.3. The last assertion of Assumption 2.1 is made in order to ensure that the convergence analysis of the domain decomposition is valid without employing the Poincaré inequality, which allows decompositions where $\partial\Omega \setminus \partial\Omega_i = \emptyset$; see Figure 1b. If the latter setting is excluded, then the analysis is valid for a broader class of functions g , especially $g = 0$.

Remark 2.4. Possible generalizations of Assumption 2.1, which we omit for sake of notational simplicity, include the cases: dependence on the spatial variable and first-order terms, e.g., $\alpha(\nabla u) = \alpha(x, u, \nabla u)$; the parameter choice $p \in (2d/(d+1), 2)$, which requires an additional set of embedding results for trace spaces; other variations similar to the p -Laplacian, e.g.,

$$\alpha(z) = (|z_1|^{p-2}z_1, \dots, |z_d|^{p-2}z_d),$$

which only require slight reformulations of the monotonicity and coercivity conditions.

Let $V = W_0^{1,p}(\Omega)$ and define the form $a : V \times V \rightarrow \mathbb{R}$ by

$$a(u, v) = \int_{\Omega} \alpha(\nabla u) \cdot \nabla v + g(u)v \, dx.$$

The weak form of (1.1) is to find $u \in V$ such that

$$(2.1) \quad a(u, v) = (f, v)_{L^2(\Omega)}, \quad \text{for all } v \in V.$$

The p -structure implies that there exist a unique weak solution of (2.1); see, e.g., [19, Theorem 2.36]. A central part of the existence proof, and our convergence analysis as well, is to observe that the p -structure directly implies that the form a is bounded, strictly monotone and coercive.

LEMMA 2.5. *If [Assumption 2.1](#) holds, then $a : V \times V \rightarrow \mathbb{R}$ is well defined and satisfies the upper bound*

$$|a(u, v)| \leq C_1 (\|\nabla u\|_{L^p(\Omega)^d}^{p-1} \|\nabla v\|_{L^p(\Omega)^d} + \|u\|_{L^r(\Omega)}^{r-1} \|v\|_{L^r(\Omega)}),$$

the strict monotonicity bound

$$a(u, u - v) - a(v, u - v) \geq c_1 (\|\nabla(u - v)\|_{L^p(\Omega)^d}^p + \|u - v\|_{L^r(\Omega)}^r)$$

and the coercivity bound

$$a(u, u) \geq c_2 (\|\nabla u\|_{L^p(\Omega)^d}^p + \|u\|_{L^r(\Omega)}^r),$$

for all $u, v \in V$.

In order to conduct the convergence analysis, we also make the following additional regularity assumption on the weak solution.

Assumption 2.6. The weak solution $u \in V$ of [\(2.1\)](#) satisfies $\alpha(\nabla u) \in C(\overline{\Omega})^d$.

Note that the above regularity assumption does *not* imply that u is a strong solution in $W^{2,p}(\Omega)$. A possible generalization of [Assumption 2.6](#) is discussed in [Remark 8.3](#).

Example 2.7. Consider the equations given by the p -Laplacian in [Example 2.2](#). If $p \geq d$ then the weak solution $u \in V$ is also in $C(\overline{\Omega})$. If in addition $f \in L^\infty(\Omega)$ and the boundary $\partial\Omega$ is $C^{1,\beta}$, then [\[12, Theorem 1\]](#) yields that $u \in C^{1,\beta}(\overline{\Omega})$. The latter implies that [Assumption 2.6](#) is valid in this context.

Finally, we will make frequent use of the fact that, under [Assumption 2.1](#), the standard $W^{1,p}(\Omega)$ -norm is equivalent to the norm

$$(2.2) \quad u \mapsto \|\nabla u\|_{L^p(\Omega)^d} + \|u\|_{L^r(\Omega)}.$$

For $r \geq p$ this follows directly by the Sobolev embedding theorem together with the assumed restrictions on (p, r) . For $r < p$ the equivalence follows by an additional bootstrap argument.

3. Function spaces and trace operators on Ω_i . We start by considering a manifold $\mathcal{M}$ in $\mathbb{R}^d$, which will play the role of $\partial\Omega_i$ or Γ . The manifold $\mathcal{M}$ is said to be Lipschitz if there exist open, overlapping sets Θ_r such that

$$\mathcal{M} = \bigcup_{r=1}^m \Theta_r,$$

where each Θ_r can be described as the graph of a Lipschitz continuous function b_r . More precisely, there exists $(d-1)$ -dimensional cubes θ_r and local charts $\psi_r : \Theta_r \rightarrow \theta_r$ that are bijective and Lipschitz continuous. The charts have the structure $\psi_r^{-1} = A_r^{-1} \circ Q_r$, where $A_r : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a coordinate transformation and

$$Q_r : \theta_r \rightarrow \mathbb{R}^d : x_r \mapsto (x_r, b_r(x_r))$$

for the Lipschitz continuous map $b_r : \theta_r \rightarrow \mathbb{R}$. A function $\mu : \mathcal{M} \rightarrow \mathbb{R}$ now has the local components $\mu \circ \psi_r^{-1}$. We refer to [\[10, Section 6.2\]](#) for further details.

On a Lipschitz manifold we may introduce a measure [\[16, Chapter 3\]](#) and thus define the integral and the space $L^p(\mathcal{M})$; see, e.g., [\[4\]](#). From [\[4, Chapters 3.4–3.5\]](#) it

follows that $L^p(\mathcal{M})$ is a Banach space and that $L^2(\mathcal{M})$ is a Hilbert space with the inner product

$$(\eta, \mu)_{L^2(\mathcal{M})} = \int_{\mathcal{M}} \eta \mu \, dS.$$

Let $\{\varphi_r\}$ be a partition of unity of $\mathcal{M}$. The integral then satisfies

$$(3.1) \quad \int_{\mathcal{M}} \mu \, dS = \sum_{r=1}^m \int_{\theta_r} (\mu \varphi_r) \circ \psi_r^{-1} |n_r| \, dx,$$

where $n_r = (\partial_1 b_r, \partial_2 b_r, \dots, \partial_{d-1} b_r, -1)$; see [16, Theorem 3.9]. Seemingly obvious properties of the integral, including

$$\int_{\mathcal{M}} \mu \, dS = \int_{\mathcal{M}_0} \mu \, dS + \int_{\mathcal{M} \setminus \mathcal{M}_0} \mu \, dS,$$

relies heavily on the observation that the integral is independent of the representation (Θ_r, A_r, b_r) and the choice of partition of unity $\{\varphi_r\}$; see [16, Theorems 3.5 and 3.7] and the comments thereafter.

The equality (3.1) also shows that our integral and L^p -spaces are equivalent to the ones used in [10]. Moreover, by [10, Lemma 6.3.5], the L^p -norm used here is equivalent to the norm

$$\mu \mapsto \left(\sum_{r=1}^m \|\mu \circ \psi_r^{-1}\|_{L^p(\theta_r)}^p \right)^{1/p}.$$

Finally, recall that for a Lipschitz manifold $\mathcal{M}$ the unit outward normal vector $\nu = (\nu^1, \dots, \nu^d)$ is defined almost everywhere; see [10, Section 6.10.1]. The normal vector is given locally by $\nu \circ \psi_r^{-1} = n_r/|n_r|$ and the Lipschitz continuity of b_r yields that $\nu^\ell \in L^\infty(\mathcal{M})$.

Assumption 3.1. The boundaries $\partial\Omega_i$ and the interface Γ are all $(d-1)$ -dimensional Lipschitz manifolds.

We use the notation $(\Theta_r^i, \theta_r^i, m_i, \psi_r^i, b_r^i, \phi_r^i, \nu_i)$ for the quantities related to the local representations of $\partial\Omega_i$. For later use, we note that

$$\nu_1 = -\nu_2 \quad \text{on } \Gamma.$$

Next, we define the fractional Sobolev spaces on the $(d-1)$ -dimensional cubes θ_r . Let $0 < s < 1$, then $W^{s,p}(\theta_r)$ is defined as all $u \in L^p(\theta_r)$ such that

$$|u|_{s,\theta_r} = \left(\int_{\theta_r} \int_{\theta_r} \frac{|u(x) - u(y)|^p}{|x - y|^{d-1+sp}} \, dx \, dy \right)^{1/p} < \infty.$$

The corresponding norm is given by

$$\|u\|_{W^{s,p}(\theta_r)} = \|u\|_{L^p(\theta_r)} + |u|_{s,\theta_r}.$$

Having defined the fractional Sobolev space on θ_r^i we can also define them on $\partial\Omega_i$. For $0 < s < 1$, introduce

$$W^{s,p}(\partial\Omega_i) = \{\mu \in L^p(\partial\Omega_i) : \mu \circ (\psi_r^i)^{-1} \in W^{s,p}(\theta_r^i), \text{ for } r = 1, \dots, m_i\},$$

equipped with the norm

$$\|\mu\|_{W^{s,p}(\partial\Omega_i)} = \left(\sum_{r=1}^{m_i} \|\mu \circ (\psi_r^i)^{-1}\|_{W^{s,p}(\theta_r^i)}^p \right)^{1/p}.$$

By the definitions of the norms, it follows directly that

$$\|\mu\|_{L^p(\partial\Omega_i)} \leq C \|\mu\|_{W^{s,p}(\partial\Omega_i)}.$$

Furthermore, the space $W^{s,p}(\partial\Omega_i)$ is complete and reflexive; see [10, Definition 6.8.6] and the comment thereafter. Next, we recapitulate the trace theorem for $W^{1,p}$ -functions on Lipschitz domains; see, e.g., [10, Theorems 6.8.13 and 6.9.2].

LEMMA 3.2. *If the [Assumptions 2.1](#) and [3.1](#) are valid, then there exists a surjective bounded linear operator $T_{\partial\Omega_i} : W^{1,p}(\Omega_i) \rightarrow W^{1-1/p,p}(\partial\Omega_i)$ such that $T_{\partial\Omega_i}u = u|_{\partial\Omega_i}$ when $u \in C^\infty(\Omega_i)$. The operator $T_{\partial\Omega_i}$ has a bounded linear right inverse $R_{\partial\Omega_i} : W^{1-1/p,p}(\partial\Omega_i) \rightarrow W^{1,p}(\Omega_i)$.*

We can then define the Sobolev spaces on Ω_i required for the domain decomposition, namely

$$V_i^0 = W_0^{1,p}(\Omega_i) \quad \text{and} \quad V_i = \{v \in W^{1,p}(\Omega_i) : (T_{\partial\Omega_i}v)|_{\partial\Omega_i \setminus \Gamma} = 0\}.$$

The spaces are equipped with the norm

$$\|v\|_{V_i} = \|\nabla v\|_{L^p(\Omega_i)^d} + \|v\|_{L^p(\Omega_i)}.$$

As for (2.2), this norm is equivalent to the standard $W^{1,p}(\Omega_i)$ -norm under [Assumption 2.1](#). Furthermore, the spaces V_i^0 and V_i are reflexive Banach spaces.

4. Function spaces and trace operators on Γ . The L^p -form of the Lions–Magenes spaces can be defined as

$$\Lambda_i = \{\mu \in L^p(\Gamma) : E_i\mu \in W^{1-1/p,p}(\partial\Omega_i)\}, \quad \text{with} \quad \|\mu\|_{\Lambda_i} = \|E_i\mu\|_{W^{1-1/p,p}(\partial\Omega_i)}.$$

Here, $E_i\mu$ denotes the extension by zero of μ to $\partial\Omega_i$. We also define the trace space

$$\Lambda = \{\mu \in L^p(\Gamma) : \mu \in \Lambda_i, \text{ for } i = 1, 2\}, \quad \text{with} \quad \|\mu\|_{\Lambda} = \|\mu\|_{\Lambda_1} + \|\mu\|_{\Lambda_2}.$$

LEMMA 4.1. *If the [Assumptions 2.1](#) and [3.1](#) hold, then Λ_i and Λ are reflexive Banach spaces.*

Proof. Observe that E_i is a linear isometry from Λ_i onto

$$(4.1) \quad R(E_i) = \{\mu \in W^{1-1/p,p}(\partial\Omega_i) : \mu|_{\partial\Omega \setminus \Gamma} = 0\}.$$

Next, consider a sequence $\{\mu^k\} \subset R(E_i)$ such that $\mu^k \rightarrow \mu$ in $W^{1-1/p,p}(\partial\Omega_i)$. Then $\mu^k|_{\Omega_i \setminus \Gamma} = 0$ and $\mu^k \rightarrow \mu$ in $L^p(\partial\Omega_i)$, which implies that

$$(4.2) \quad \int_{\partial\Omega_i \setminus \Gamma} |\mu|^p dS = \int_{\partial\Omega_i \setminus \Gamma} |\mu^k - \mu|^p dS \leq \int_{\partial\Omega_i} |\mu^k - \mu|^p dS \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

Hence, $\mu \in R(E_i)$ and consequently $R(E_i)$ is a closed subset of $W^{1-1/p,p}(\partial\Omega_i)$. The space Λ_i is therefore isomorphic to a closed subset of the reflexive Banach space $W^{1-1/p,p}(\partial\Omega_i)$, i.e., Λ_i is complete and reflexive [11, Chapter 8, Theorem 15].

To prove that the same holds true for Λ introduce the reflexive Banach space $X = W^{1-1/p,p}(\partial\Omega_1) \times W^{1-1/p,p}(\partial\Omega_2)$, with the norm

$$\|(\mu_1, \mu_2)\|_X = \|\mu_1\|_{W^{1-1/p,p}(\partial\Omega_1)} + \|\mu_2\|_{W^{1-1/p,p}(\partial\Omega_2)},$$

and the operator $E : \Lambda \rightarrow X$ defined by $E\mu = (E_1\mu, E_2\mu)$. As E is a linear isometry from Λ onto

$$R(E) = \{(\mu_1, \mu_2) \in X : \mu_1|_{\partial\Omega_1 \setminus \Gamma} = 0, \mu_2|_{\partial\Omega_2 \setminus \Gamma} = 0, \mu_1|_{\Gamma} = \mu_2|_{\Gamma}\},$$

it is again sufficient to prove that $R(E)$ is a closed subset of X . Let $\{(\mu_1^k, \mu_2^k)\} \subset R(E)$ be a convergent sequence in X with the limit (μ_1, μ_2) . By the same argument as (4.2), we obtain that $\mu_i|_{\Omega_i \setminus \Gamma} = 0$. As $\mu_1^k|_{\Gamma} = \mu_2^k|_{\Gamma}$, we also have the limit

$$\int_{\Gamma} |\mu_1 - \mu_2|^p dS \leq 2^{p-1} \left(\int_{\Gamma} |\mu_1^k - \mu_1|^p dS + \int_{\Gamma} |\mu_2^k - \mu_2|^p dS \right) \rightarrow 0, \quad \text{as } k \rightarrow \infty,$$

i.e., $\mu_1|_{\Gamma} = \mu_2|_{\Gamma}$ in $L^p(\Gamma)$ and we obtain that $(\mu_1, \mu_2) \in R(E)$. Thus, $R(E)$ is closed and Λ is therefore a reflexive Banach space. $\square$

LEMMA 4.2. *If the Assumptions 2.1 and 3.1 hold, then Λ_i and Λ are dense in $L^2(\Gamma)$.*

The proof of the lemma is almost identical to the proof of [10, Theorem 6.6.3] and is therefore left out.

Remark 4.3. We conjecture that $\Lambda_1 = \Lambda_2$. However, we will move on to a $L^2(\Gamma)$ -framework for which it is not necessary to make this identification.

Collecting these results yield the Gelfand triplets

$$\Lambda_i \xhookrightarrow{d} L^2(\Gamma) \cong L^2(\Gamma)^* \xhookrightarrow{d} \Lambda_i^* \quad \text{and} \quad \Lambda \xhookrightarrow{d} L^2(\Gamma) \cong L^2(\Gamma)^* \xhookrightarrow{d} \Lambda^*.$$

For future use, we introduce the Riesz isomorphism on $L^2(\Gamma)$ given by

$$J : L^2(\Gamma) \rightarrow L^2(\Gamma)^* : \mu \mapsto (\mu, \cdot)_{L^2(\Gamma)},$$

which satisfies the relations

$$\langle J\eta, \mu_i \rangle_{\Lambda_i^* \times \Lambda_i} = (\eta, \mu_i)_{L^2(\Gamma)} \quad \text{and} \quad \langle J\eta, \mu \rangle_{\Lambda^* \times \Lambda} = (\eta, \mu)_{L^2(\Gamma)},$$

for all $\eta \in L^2(\Gamma)$, $\mu_i \in \Lambda_i$ and $\mu \in \Lambda$. Here, $\langle \cdot, \cdot \rangle_{X^* \times X}$ denotes the dual pairing between a Banach space X and its dual X^* . In the following we will drop the subscripts on the dual pairings.

In order to relate the spaces V_i and Λ_i , we observe that for $v \in V_i$ one has $T_{\partial\Omega_i}v \in R(E_i)$; see (4.1). Hence, the trace operator

$$T_i : V_i \rightarrow \Lambda_i : v \mapsto (T_{\partial\Omega_i}v)|_{\Gamma}$$

is well defined. We also introduce the linear operator

$$R_i : \Lambda_i \rightarrow V_i : \mu \mapsto R_{\partial\Omega_i}E_i\mu.$$

LEMMA 4.4. *If the Assumptions 2.1 and 3.1 hold, then T_i and R_i are bounded, and R_i is a right inverse of T_i .*

Proof. For $v \in V_i$ and $\mu \in \Lambda_i$ we have, by Lemma 3.2, that

$$\|T_i v\|_{\Lambda_i} = \|E_i((T_{\partial\Omega_i} v)|_\Gamma)\|_{W^{1-1/p,p}(\partial\Omega_i)} = \|T_{\partial\Omega_i} v\|_{W^{1-1/p,p}(\partial\Omega_i)} \leq C_i \|v\|_{V_i}$$

$$\text{and } \|R_i \mu\|_{V_i} = \|R_{\partial\Omega_i} E_i \mu\|_{V_i} \leq C_i \|E_i \mu\|_{W^{1-1/p,p}(\partial\Omega_i)} = C_i \|\mu\|_{\Lambda_i}.$$

Hence, the linear operators T_i and R_i are bounded. Furthermore, for every $\mu \in \Lambda_i$ we have

$$T_i R_i \mu = (T_{\partial\Omega_i} R_{\partial\Omega_i} E_i \mu)|_\Gamma = (E_i \mu)|_\Gamma = \mu,$$

i.e., R_i is a right inverse of T_i . $\square$

We continue by deriving a few useful properties related to the operator T_i .

LEMMA 4.5. *If the Assumptions 2.1 and 3.1 hold and $v \in V$, then $\mu = T_1 v|_{\Omega_1} = T_2 v|_{\Omega_2}$ is an element in Λ .*

Proof. Let $v \in V$. As $C_0^\infty(\Omega)$ is dense in V , there exists a sequence $\{v^k\} \subset C_0^\infty(\Omega)$ such that $v^k \rightarrow v$ in V . Set $v_i = v|_{\Omega_i}$ and $v_i^k = v^k|_{\Omega_i}$. Clearly, $T_1 v_1^k = T_2 v_2^k$. Since $v^k \rightarrow v$ in V , we also have that $v_i^k \rightarrow v_i$ in V_i . The continuity of T_i then implies that $T_i v_i^k \rightarrow T_i v_i$ in Λ_i . Putting this together gives us

$$\Lambda_1 \ni T_1 v_1 = \lim_{k \rightarrow \infty} T_1 v_1^k = \lim_{k \rightarrow \infty} T_2 v_2^k = T_2 v_2 \in \Lambda_2 \quad \text{in } L^p(\Gamma).$$

If we now define $\mu = T_1 v_1 = T_2 v_2$, then μ is an element in $\Lambda = \Lambda_1 \cap \Lambda_2$. $\square$

LEMMA 4.6. *Let the Assumptions 2.1 and 3.1 hold. If two elements $v_1 \in V_1$ and $v_2 \in V_2$ satisfies $T_1 v_1 = T_2 v_2$, then $v = \{v_1 \text{ on } \Omega_1; v_2 \text{ on } \Omega_2\}$ is an element in V .*

Proof. It is clear that $v \in L^p(\Omega)$. For each component $1 \leq \ell \leq d$, there exists a weak derivative $\partial_\ell v_i \in L^p(\Omega_i)$ of $v_i \in V_i \subset W^{1,p}(\Omega_i)$. If we define

$$z_\ell = \{\partial_\ell v_1 \text{ on } \Omega_1; \partial_\ell v_2 \text{ on } \Omega_2\},$$

then $z_\ell \in L^p(\Omega)$. Let $w \in C_0^\infty(\Omega)$ and set $w_i = w|_{\Omega_i} \in C^\infty(\Omega_i)$. The $W^{1,p}(\Omega_i)$ -version of Green's formula [17, Section 3.1.2] yields that

$$\begin{aligned} \int_\Omega z_\ell w \, dx &= \sum_{i=1}^2 \int_{\Omega_i} \partial_\ell v_i w_i \, dx = \sum_{i=1}^2 - \int_{\Omega_i} v_i \partial_\ell w_i \, dx + \int_{\partial\Omega_i} (T_{\partial\Omega_i} v_i) w_i \nu_i^\ell \, dS \\ &= - \int_\Omega v \partial_\ell w \, dx + \sum_{i=1}^2 \int_\Gamma (T_i v_i) w \nu_i^\ell \, dS = - \int_\Omega v \partial_\ell w \, dx, \end{aligned}$$

i.e., z_ℓ is the ℓ th weak partial derivative of v . By construction $T_{\partial\Omega} v = 0$, and v is therefore an element in V . $\square$

5. Transmission problem and the Robin–Robin method. In order to state the Robin–Robin method we reformulate the nonlinear elliptic equation (2.1) on Ω into two equations on Ω_i connected via Γ , i.e., we consider a nonlinear transmission problem. To this end, on each V_i we define $a_i : V_i \times V_i \rightarrow \mathbb{R}$ by

$$a_i(u_i, v_i) = \int_{\Omega_i} \alpha(\nabla u_i) \cdot \nabla v_i + g(u_i) v_i \, dx.$$

We also define $f_i = f|_{\Omega_i} \in L^2(\Omega_i)$.

LEMMA 5.1. *If the [Assumptions 2.1](#) and [3.1](#) hold, then $a_i : V_i \times V_i \rightarrow \mathbb{R}$ is well defined and satisfies the growth, strict monotonicity and coercivity bounds stated in [Lemma 2.5](#), with the terms (a, V, Ω) replaced by (a_i, V_i, Ω_i) .*

The weak form of the nonlinear transmission problem is then to find $(u_1, u_2) \in V_1 \times V_2$ such that

$$(5.1) \quad \begin{cases} a_i(u_i, v_i) = (f_i, v_i)_{L^2(\Omega_i)}, & \text{for all } v_i \in V_i^0, i = 1, 2, \\ T_1 u_1 = T_2 u_2, \\ \sum_{i=1}^2 a_i(u_i, R_i \mu) - (f_i, R_i \mu)_{L^2(\Omega_i)} = 0, & \text{for all } \mu \in \Lambda. \end{cases}$$

The framework given in [section 4](#) enables us to prove equivalence between the nonlinear elliptic equation and the nonlinear transmission problem along the same lines as done for linear equations [[18](#), [Lemma 1.2.1](#)].

THEOREM 5.2. *Let [Assumptions 2.1](#) and [3.1](#) hold. If $u \in V$ solves [\(2.1\)](#), then $(u_1, u_2) = (u|_{\Omega_1}, u|_{\Omega_2})$ solves [\(5.1\)](#). Conversely, if (u_1, u_2) solves [\(5.1\)](#), then $u = \{u_1 \text{ on } \Omega_1; u_2 \text{ on } \Omega_2\}$ solves [\(2.1\)](#).*

Proof. Assume that $u \in V$ solves [\(2.1\)](#) and define $(u_1, u_2) = (u|_{\Omega_1}, u|_{\Omega_2}) \in V_1 \times V_2$. For $v_i \in V_i^0$ we can extend by zero to $w_i \in V$, by using [Lemma 4.6](#). Then

$$a_i(u_i, v_i) = a(u, w_i) = (f, w_i)_{L^2(\Omega)} = (f_i, v_i)_{L^2(\Omega_i)}.$$

Moreover, $T_1 u_1 = T_2 u_2$ follows immediately from [Lemma 4.5](#). For an arbitrary $\mu \in \Lambda$ let $v_i = R_i \mu$. Then, by [Lemma 4.6](#), $v = \{v_1 \text{ on } \Omega_1; v_2 \text{ on } \Omega_2\}$ is an element in V and

$$a_1(u_1, R_1 \mu) + a_2(u_2, R_2 \mu) = a(u, v) = (f, v)_{L^2(\Omega)} = (f_1, R_1 \mu)_{L^2(\Omega_1)} + (f_2, R_2 \mu)_{L^2(\Omega_2)}.$$

This proves that (u_1, u_2) solves [\(5.1\)](#). Conversely, let $(u_1, u_2) \in V_1 \times V_2$ be a solution to [\(5.1\)](#) and define $u = \{u_1 \text{ on } \Omega_1; u_2 \text{ on } \Omega_2\}$. By [Lemma 4.6](#), we have that $u \in V$. Next, consider $v \in V$ and let $v_i = v|_{\Omega_i} \in V_i$. From [Lemma 4.5](#) we have that $\mu = T_i v_i$ is well defined and $\mu \in \Lambda$. The observation that $v_i - R_i \mu \in V_i^0$, for $i = 1, 2$, then implies the equality

$$\begin{aligned} a(u, v) &= \sum_{i=1}^2 a_i(u_i, v_i - R_i \mu) + a_i(u_i, R_i \mu) = \sum_{i=1}^2 (f_i, v_i - R_i \mu)_{L^2(\Omega_i)} + a_i(u_i, R_i \mu) \\ &= \sum_{i=1}^2 (f_i, v_i)_{L^2(\Omega_i)} + a_i(u_i, R_i \mu) - (f_i, R_i \mu)_{L^2(\Omega_i)} = (f, v)_{L^2(\Omega)}. \end{aligned}$$

As v can be chosen arbitrarily, u solves [\(2.1\)](#). □

REMARK 5.3. As the nonlinear elliptic equation [\(2.1\)](#) has a unique weak solution, [Theorem 5.2](#) implies that the same holds true for the nonlinear transmission problem [\(5.1\)](#).

REMARK 5.4. The equivalence can easily be generalized to more than two subdomains without cross points as in [Figure 1c](#). However, for domain decompositions with cross points such as in [Figure 1d](#), the situation is more delicate. A proof of the equivalence for quasilinear equations in $H^1(\Omega)$ can be found in [[20](#)]. This result can most likely be generalized to our $W^{1,p}(\Omega)$ -setting, but we will study this aspects elsewhere.

In order to approximate the weak solution $(u_1^n, u_2^n) \in V_1 \times V_2$ of the nonlinear transmission problem in a parallel fashion, we consider the Robin–Robin method. Multiplying by test functions and formally applying Green’s formula to the equations (1.4) yield that the weak form of the method is given by finding $(u_1^n, u_2^n) \in V_1 \times V_2$, for $n = 1, 2, \dots$, such that

$$(5.2) \quad \begin{cases} a_1(u_1^{n+1}, v_1) = (f_1, v_1)_{L^2(\Omega_1)}, & \text{for all } v_1 \in V_1^0 \\ a_1(u_1^{n+1}, R_1\mu) - (f_1, R_1\mu)_{L^2(\Omega_1)} + a_2(u_2^n, R_2\mu) \\ \quad - (f_2, R_2\mu)_{L^2(\Omega_2)} = s(T_2 u_2^n - T_1 u_1^{n+1}, \mu)_{L^2(\Gamma)}, & \text{for all } \mu \in \Lambda, \\ a_2(u_2^{n+1}, v_2) = (f_2, v_2)_{L^2(\Omega_2)}, & \text{for all } v_2 \in V_2^0 \\ a_2(u_2^{n+1}, R_2\mu) - (f_2, R_2\mu)_{L^2(\Omega_2)} + a_1(u_1^{n+1}, R_1\mu) \\ \quad - (f_1, R_1\mu)_{L^2(\Omega_1)} = s(T_1 u_1^{n+1} - T_2 u_2^{n+1}, \mu)_{L^2(\Gamma)}, & \text{for all } \mu \in \Lambda, \end{cases}$$

where $u_2^0 \in V_2$ is an initial guess and $s > 0$ is the given method parameter.

6. Interface formulations. The ambition is now to reformulate the nonlinear transmission problem and the Robin–Robin method, which are all given on the domains Ω_i , into problems and methods only stated on the interface Γ . As a preparation, we observe that nonlinear elliptic equations on Ω_i with inhomogeneous Dirichlet conditions have unique weak solutions.

LEMMA 6.1. *If the Assumptions 2.1 and 3.1 hold, then for each $\eta \in \Lambda_i$ there exists a unique $u_i \in W^{1,p}(\Omega_i)$ such that*

$$(6.1) \quad a_i(u_i, v_i) = (f_i, v_i)_{L^2(\Omega_i)}, \quad \text{for all } v_i \in V_i^0,$$

and $T_{\partial\Omega_i} u_i = E_i \eta$ in $W^{1-1/p,p}(\partial\Omega_i)$.

The proof can, e.g., be found in [19, Theorem 2.36]. With the notation of Lemma 6.1, consider the operator

$$F_i : \eta \mapsto u_i,$$

i.e., the map from a given boundary value on Γ to the corresponding weak solution of the nonlinear elliptic problem (6.1) on Ω_i . From the statement of Lemma 6.1 we see that

$$F_i : \Lambda_i \rightarrow V_i \quad \text{and} \quad T_i F_i \eta = \eta \text{ for } \eta \in \Lambda_i.$$

In other words, the operator F_i is a nonlinear right inverse to T_i . This property will be frequently used as it, together with the boundedness and linearity of T_i , gives rise to bounds of the forms

$$\|\eta\|_{\Lambda_i} \leq C_i \|F_i \eta\|_{V_i} \quad \text{and} \quad \|\eta - \mu\|_{\Lambda_i} \leq C_i \|F_i \eta - F_i \mu\|_{V_i}.$$

We can now define the nonlinear Steklov–Poincaré operators as

$$\langle S_i \eta, \mu \rangle = a_i(F_i \eta, R_i \mu) - (f_i, R_i \mu)_{L^2(\Omega_i)}, \quad \text{for all } \eta, \mu \in \Lambda_i, \quad \text{and}$$

$$\langle S \eta, \mu \rangle = \sum_{i=1}^2 \langle S_i \eta, \mu \rangle = \sum_{i=1}^2 a_i(F_i \eta, R_i \mu) - (f_i, R_i \mu)_{L^2(\Omega_i)}, \quad \text{for all } \eta, \mu \in \Lambda.$$

Thus, we may restate the nonlinear transmission problem (5.1) as the Steklov–Poincaré equation, i.e., finding $\eta \in \Lambda$ such that

$$(6.2) \quad \langle S \eta, \mu \rangle = 0, \quad \text{for all } \mu \in \Lambda.$$

That the reformulation is possible follows directly from the definitions of the operators F_i and S , but we state this as a lemma for future reference.

LEMMA 6.2. *Let the [Assumptions 2.1](#) and [3.1](#) hold. If (u_1, u_2) solves [\(5.1\)](#), then $\eta = T_1 u_1 = T_2 u_2$ solves [\(6.2\)](#). Conversely, if η solves [\(6.2\)](#), then $(u_1, u_2) = (F_1 \eta, F_2 \eta)$ solves [\(5.1\)](#).*

We next turn to the Robin–Robin method, which is equivalent to the Peaceman–Rachford splitting on the interface Γ . The weak form of the splitting is given by finding $(\eta_1^n, \eta_2^n) \in \Lambda_1 \times \Lambda_2$, for $n = 1, 2, \dots$, such that

$$(6.3) \quad \begin{cases} \langle (sJ + S_1)\eta_1^{n+1}, \mu \rangle = \langle (sJ - S_2)\eta_2^n, \mu \rangle, \\ \langle (sJ + S_2)\eta_2^{n+1}, \mu \rangle = \langle (sJ - S_1)\eta_1^{n+1}, \mu \rangle, \end{cases}$$

for all $\mu \in \Lambda$, where $\eta_2^0 \in \Lambda_2$ is an initial guess.

The observation that the Robin–Robin method and the Peaceman–Rachford splitting are equivalent for linear elliptic equations was made in [\[1\]](#). The equivalence was also utilized in [\[5, Section 4.4.1\]](#) for the linear setting of the Stokes–Darcy coupling.

LEMMA 6.3. *Let the [Assumptions 2.1](#) and [3.1](#) be valid. If $(u_1^n, u_2^n)_{n \geq 1}$ is a weak Robin–Robin approximation [\(5.2\)](#), then $(\eta_1^n, \eta_2^n)_{n \geq 1} = (T_1 u_1^n, T_2 u_2^n)_{n \geq 1}$ is a weak Peaceman–Rachford approximation [\(6.3\)](#), with $\eta_2^0 = T_2 u_2^0$. Conversely, if $(\eta_1^n, \eta_2^n)_{n \geq 1}$ is given by [\(6.3\)](#), then $(u_1^n, u_2^n)_{n \geq 1} = (F_1 \eta_1^n, F_2 \eta_2^n)_{n \geq 1}$ fulfils [\(5.2\)](#), with $u_2^0 = F_2 \eta_2^0$.*

Proof. First assume that $(u_1^n, u_2^n)_{n \geq 1} \subset V_1 \times V_2$ is a weak Robin–Robin approximation and define $\eta_i^n = T_i u_i^n \in \Lambda_i$. This definition, the existence of a unique solution of [\(6.1\)](#) and the first and third assertions of [\(5.2\)](#) yield the identification $u_i^n = F_i \eta_i^n$. Inserting this into the second and fourth assertion of [\(5.2\)](#) gives us

$$\begin{aligned} & s(T_1 F_1 \eta_1^{n+1}, \mu)_{L^2(\Gamma)} + a_1(F_1 \eta_1^{n+1}, R_1 \mu) - (f, R_1 \mu)_{L^2(\Omega_1)} \\ & = s(T_2 F_2 \eta_2^n, \mu)_{L^2(\Gamma)} - a_2(F_2 \eta_2^n, R_2 \mu) + (f, R_2 \mu)_{L^2(\Omega_2)}, \quad \text{and} \\ & s(T_2 F_2 \eta_2^{n+1}, \mu)_{L^2(\Gamma)} + a_2(F_2 \eta_2^{n+1}, R_2 \mu) - (f, R_2 \mu)_{L^2(\Omega_2)} \\ & = s(T_1 F_1 \eta_1^{n+1}, \mu)_{L^2(\Gamma)} - a_1(F_1 \eta_1^{n+1}, R_1 \mu) + (f, R_1 \mu)_{L^2(\Omega_1)}, \end{aligned}$$

for all $\mu \in \Lambda$, which is precisely the weak form of the Peaceman–Rachford splitting [\(6.3\)](#), with $\eta_2^0 = T_2 u_2^0$. Conversely, suppose that $(\eta_1^n, \eta_2^n)_{n \geq 1} \subset \Lambda_1 \times \Lambda_2$ is a weak Peaceman–Rachford approximation and define $u_i^n = F_i \eta_i^n \in V_i$. Inserting this into [\(6.3\)](#) directly gives that $(u_1^n, u_2^n)_{n \geq 1}$, with $u_2^0 = F_2 \eta_2^0$, is a weak Robin–Robin approximation [\(5.2\)](#). $\square$

Remark 6.4. For now we do not know if the weak Robin–Robin and Peaceman–Rachford approximations actually exist, but we will return to this issue shortly.

7. Properties of the nonlinear Steklov–Poincaré operators. We proceed by deriving the central properties of the Steklov–Poincaré operators S_i, S when interpreted as maps from Λ_i, Λ into the corresponding dual spaces.

LEMMA 7.1. *If the [Assumptions 2.1](#) and [3.1](#) hold, then $S_i : \Lambda_i \rightarrow \Lambda_i^*$ and $S : \Lambda \rightarrow \Lambda^*$ are well defined.*

Proof. The linearity of the functionals $S_i \eta$ and $S \eta$ follow by definition. As $F_i \eta \in$

V_i we have, by [Lemma 5.1](#), that

$$\begin{aligned} |\langle S_i \eta, \mu \rangle| &\leq |a_i(F_i \eta, R_i \mu)| + |(f_i, R_i \mu)_{L^2(\Omega_i)}| \\ &\leq c_i (\|\nabla F_i \eta\|_{L^p(\Omega_i)^d}^{p-1} \|\nabla R_i \mu\|_{L^p(\Omega_i)^d} + \|F_i \eta\|_{L^r(\Omega_i)}^{r-1} \|R_i \mu\|_{L^r(\Omega_i)}) \\ &\quad + \|f_i\|_{L^2(\Omega_i)} \|R_i \mu\|_{L^2(\Omega_i)} \\ &\leq C_i (\|\nabla F_i \eta\|_{V_i}, \|f_i\|_{L^2(\Omega_i)}) \|R_i \mu\|_{V_i} \leq C_i \|\mu\|_{\Lambda_i}, \end{aligned}$$

for all $\mu \in \Lambda_i$. Thus, $S_i \eta$ is a bounded functional on Λ_i . The boundedness of $S \eta$ follows directly by summing up the bounds for S_i . $\square$

LEMMA 7.2. *If the [Assumptions 2.1](#) and [3.1](#) hold, then the operators $S_i : \Lambda_i \rightarrow \Lambda_i^*$ and $S : \Lambda \rightarrow \Lambda^*$ are strictly monotone with*

$$\langle S_i \eta - S_i \mu, \eta - \mu \rangle \geq c_i (\|\nabla(F_i \eta - F_i \mu)\|_{L^p(\Omega_i)^d}^p + \|F_i \eta - F_i \mu\|_{L^r(\Omega_i)}^r),$$

for all $\eta, \mu \in \Lambda_i$, and

$$\langle S \eta - S \mu, \eta - \mu \rangle \geq c \sum_{i=1}^2 (\|\nabla(F_i \eta - F_i \mu)\|_{L^p(\Omega_i)^d}^p + \|F_i \eta - F_i \mu\|_{L^r(\Omega_i)}^r),$$

for all $\eta, \mu \in \Lambda$, respectively.

Proof. Since, $w_i = R_i(\eta - \mu) - (F_i \eta - F_i \mu) \in V_i^0$, for all $\eta, \mu \in \Lambda_i$, we have according to the definition of F_i that

$$(7.1) \quad a_i(F_i \eta, w_i) - a_i(F_i \mu, w_i) = 0.$$

By this equality and [Lemma 5.1](#), it follows that

$$\begin{aligned} \langle S_i \eta - S_i \mu, \eta - \mu \rangle &= a_i(F_i \eta, R_i(\eta - \mu)) - a_i(F_i \mu, R_i(\eta - \mu)) \\ &= a_i(F_i \eta, w_i) + a_i(F_i \eta, F_i \eta - F_i \mu) - a_i(F_i \mu, w_i) - a_i(F_i \mu, F_i \eta - F_i \mu) \\ &\geq c_i (\|\nabla(F_i \eta - F_i \mu)\|_{L^p(\Omega_i)^d}^p + \|F_i \eta - F_i \mu\|_{L^r(\Omega_i)}^r), \end{aligned}$$

for all $\eta, \mu \in \Lambda_i$, which proves the monotonicity bound for S_i . The bound for S follows directly by summing up the bounds for S_i . $\square$

LEMMA 7.3. *If the [Assumptions 2.1](#) and [3.1](#) hold, then the operators $S_i : \Lambda_i \rightarrow \Lambda_i^*$ and $S : \Lambda \rightarrow \Lambda^*$ are coercive, i.e.,*

$$\lim_{\|\eta\|_{\Lambda_i} \rightarrow \infty} \frac{\langle S_i \eta, \eta \rangle}{\|\eta\|_{\Lambda_i}} = \infty \quad \text{and} \quad \lim_{\|\eta\|_{\Lambda} \rightarrow \infty} \frac{\langle S \eta, \eta \rangle}{\|\eta\|_{\Lambda}} = \infty.$$

Proof. As $\|F_i \eta\|_{V_i} = \|\nabla F_i \eta\|_{L^p(\Omega_i)^d} + \|F_i \eta\|_{L^r(\Omega_i)} \geq c_i \|\eta\|_{\Lambda_i}$, we have that

$$P(\|\nabla F_i \eta\|_{L^p(\Omega_i)^d}, \|F_i \eta\|_{L^r(\Omega_i)}) \rightarrow \infty, \quad \text{as } \|\eta\|_{\Lambda_i} \rightarrow \infty,$$

where $P(x, y) = (x^p + y^r)/(x + y)$. In particular, we assume from now on that

$$P(\|\nabla F_i \eta\|_{L^p(\Omega_i)^d}, \|F_i \eta\|_{L^r(\Omega_i)}) \geq \|f_i\|_{L^2(\Omega_i)}.$$

By observing that $R_i\eta - F_i\eta \in V_i^0$, [Lemma 5.1](#) yields the lower bound

$$\begin{aligned}
\langle S_i\eta, \eta \rangle &= a_i(F_i\eta, F_i\eta) + a_i(F_i\eta, R_i\eta - F_i\eta) - (f_i, R_i\eta)_{L^2(\Omega_i)} \\
&= a_i(F_i\eta, F_i\eta) + (f_i, R_i\eta - F_i\eta)_{L^2(\Omega_i)} - (f_i, R_i\eta)_{L^2(\Omega_i)} \\
&\geq c_i(\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}^p + \|F_i\eta\|_{L^r(\Omega_i)}^r) - (f_i, F_i\eta)_{L^2(\Omega_i)} \\
&\geq c_i P(\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}, \|F_i\eta\|_{L^r(\Omega_i)}) \|F_i\eta\|_{V_i} - \|f_i\|_{L^2(\Omega_i)} \|F_i\eta\|_{V_i} \\
&\geq c_i(P(\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}, \|F_i\eta\|_{L^r(\Omega_i)}) - \|f_i\|_{L^2(\Omega_i)}) \|\eta\|_{\Lambda_i},
\end{aligned}$$

which implies that S_i is coercive. For S we obtain that

$$\frac{\langle S\eta, \eta \rangle}{\|\eta\|_{\Lambda}} \geq \frac{\sum_{i=1}^2 (c_i P(\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}, \|F_i\eta\|_{L^r(\Omega_i)}) - \|f_i\|_{L^2}) \|\eta\|_{\Lambda_i}}{\|\eta\|_{\Lambda_1} + \|\eta\|_{\Lambda_2}},$$

which tends to infinity as $\|\eta\|_{\Lambda}$ tends to infinity. Thus, S is also coercive. $\square$

In order to prove that the operators S_i, S are demicontinuous, we first consider the continuity of the operators F_i .

LEMMA 7.4. *If the [Assumptions 2.1](#) and [3.1](#) hold, then the nonlinear operators $F_i : \Lambda_i \rightarrow V_i$ are continuous.*

Proof. Let η, μ be elements in Λ_i . Using the equality [\(7.1\)](#) together with [Lemma 5.1](#) gives us the bound

$$\begin{aligned}
(7.2) \quad &c_i(\|\nabla(F_i\eta - F_i\mu)\|_{L^p(\Omega_i)^d}^p + \|F_i\eta - F_i\mu\|_{L^r(\Omega_i)}^r) \\
&\leq a_i(F_i\eta, F_i\eta - F_i\mu) - a_i(F_i\mu, F_i\eta - F_i\mu) \\
&= a_i(F_i\eta, R_i(\eta - \mu)) - a_i(F_i\mu, w_i) - a_i(F_i\mu, R_i(\eta - \mu)) + a_i(F_i\mu, w_i) \\
&\leq C_i(\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}^{p-1} \|\nabla R_i(\eta - \mu)\|_{L^p(\Omega_i)^d} + \|F_i\eta\|_{L^r(\Omega_i)}^{r-1} \|R_i(\eta - \mu)\|_{L^r(\Omega_i)} \\
&\quad + \|\nabla F_i\mu\|_{L^p(\Omega_i)^d}^{p-1} \|\nabla R_i(\eta - \mu)\|_{L^p(\Omega_i)^d} + \|F_i\mu\|_{L^r(\Omega_i)}^{r-1} \|R_i(\eta - \mu)\|_{L^r(\Omega_i)}) \\
&\leq C_i(\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}^{p-1} + \|F_i\eta\|_{L^r(\Omega_i)}^{r-1} \\
&\quad + \|\nabla F_i\mu\|_{L^p(\Omega_i)^d}^{p-1} + \|F_i\mu\|_{L^r(\Omega_i)}^{r-1}) \|\eta - \mu\|_{\Lambda_i}.
\end{aligned}$$

Letting $\mu = 0$ in [\(7.2\)](#) and employing the inequality $|x|^p - 2^{p-1}|y|^p \leq 2^{p-1}|x - y|^p$, twice, yield that

$$\begin{aligned}
&c_i(\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}^p + \|F_i\eta\|_{L^r(\Omega_i)}^r - 2^{p-1}\|\nabla F_i 0\|_{L^p(\Omega_i)^d}^p - 2^{r-1}\|F_i 0\|_{L^r(\Omega_i)}^r) \\
&\leq c_i(2^{p-1}\|\nabla(F_i\eta - F_i 0)\|_{L^p(\Omega_i)^d}^p + 2^{r-1}\|F_i\eta - F_i 0\|_{L^r(\Omega_i)}^r) \\
&\leq C_i(\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}^{p-1} + \|F_i\eta\|_{L^r(\Omega_i)}^{r-1} + \|\nabla F_i 0\|_{L^p(\Omega_i)^d}^{p-1} + \|F_i 0\|_{L^r(\Omega_i)}^{r-1}) \|\eta\|_{\Lambda_i}.
\end{aligned}$$

Thus, we have a bound of the form

$$(7.3) \quad \frac{\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}^p + \|F_i\eta\|_{L^r(\Omega_i)}^r - c_1}{\|\nabla F_i\eta\|_{L^p(\Omega_i)^d}^{p-1} + \|F_i\eta\|_{L^r(\Omega_i)}^{r-1} + c_2} \leq C_i \|\eta\|_{\Lambda_i},$$

for every $\eta \in \Lambda_i$, where $c_\ell = c_\ell(\|\nabla F_i 0\|_{L^p(\Omega_i)^d}, \|F_i 0\|_{L^r(\Omega_i)}) \geq 0$.

Assume that $\eta^k \rightarrow \eta$ in Λ_i . As η^k is bounded in Λ_i , the bound [\(7.3\)](#) implies that $\nabla F_i\eta^k$ and $F_i\eta^k$ are bounded in $L^p(\Omega_i)^d$ and $L^r(\Omega_i)$, respectively. By setting $\mu = \eta^k$ in [\(7.2\)](#), we finally obtain that $\nabla F_i\eta^k \rightarrow \nabla F_i\eta$ in $L^p(\Omega_i)^d$ and $F_i\eta^k \rightarrow F_i\eta$ in $L^r(\Omega_i)$, i.e., $F_i\eta^k \rightarrow F_i\eta$ in V_i . $\square$

LEMMA 7.5. *If the [Assumptions 2.1](#) and [3.1](#) hold, then the operators $S_i : \Lambda_i \rightarrow \Lambda_i^*$ and $S : \Lambda \rightarrow \Lambda^*$ are demicontinuous, i.e., if $\eta_i^k \rightarrow \eta_i$ in Λ_i and $\eta^k \rightarrow \eta$ in Λ then*

$$\langle S_i \eta_i^k - S_i \eta_i, \mu_i \rangle \rightarrow 0 \quad \text{and} \quad \langle S \eta^k - S \eta, \mu \rangle \rightarrow 0,$$

for all $\mu_i \in \Lambda_i$ and $\mu \in \Lambda$.

Proof. Assume that $\eta_i^k \rightarrow \eta_i$ in Λ_i . [Lemma 7.4](#) then implies that $\nabla F_i \eta_i^k \rightarrow \nabla F_i \eta_i$ in $L^p(\Omega_i)^d$ and $F_i \eta_i^k \rightarrow F_i \eta_i$ in $L^r(\Omega_i)$. By the assumed continuity and boundedness of the functions $\alpha : z \mapsto (\alpha_1(z), \dots, \alpha_d(z))$ and g , we also have that the corresponding Nemyckii operators $\alpha_\ell : L^p(\Omega_i)^d \rightarrow L^{p/(p-1)}(\Omega_i)$ and $g : L^r(\Omega_i) \rightarrow L^{r/(r-1)}(\Omega_i)$ are continuous [[25](#), Proposition 26.6]. The demicontinuity of S_i then holds, as

$$\begin{aligned} |\langle S_i \eta_i^k - S_i \eta_i, \mu_i \rangle| &\leq \left(\sum_{\ell=1}^d \|\alpha_\ell(\nabla F_i \eta_i^k) - \alpha_\ell(\nabla F_i \eta_i)\|_{L^{p/(p-1)}(\Omega_i)} \right. \\ &\quad \left. + \|g(F_i \eta_i^k) - g(F_i \eta_i)\|_{L^{r/(r-1)}(\Omega_i)} \right) \|R_i \mu_i\|_{V_i}, \end{aligned}$$

for every $\mu_i \in \Lambda_i$. The demicontinuity of S_i directly implies the same property for S , as a convergent sequence $\{\eta^k\}$ in Λ is also convergent in Λ_1 and Λ_2 . $\square$

THEOREM 7.6. *If the [Assumptions 2.1](#) and [3.1](#) hold, then the nonlinear Steklov–Poincaré operators $S_i : \Lambda_i \rightarrow \Lambda_i^*$ and $S : \Lambda \rightarrow \Lambda^*$ are bijective.*

Proof. The spaces Λ_i and Λ are real, reflexive Banach spaces and, by [Lemmas 7.2](#), [7.3](#) and [7.5](#), the operators $S_i : \Lambda_i \rightarrow \Lambda_i^*$ and $S : \Lambda \rightarrow \Lambda^*$ are all strictly monotone, coercive and demicontinuous. With these properties, the Browder–Minty theorem; see, e.g., [[25](#), Theorem 26.A(a,c,f)], implies that the operators are bijective. $\square$

The next corollary follows by the same argumentation as for the bijectivity of S_i .

COROLLARY 7.7. *If the [Assumptions 2.1](#) and [3.1](#) hold, then the operators $sJ + S_i : \Lambda_i \rightarrow \Lambda_i^*$ are bijective, for every $s > 0$.*

8. Existence and convergence of the Robin–Robin method. The weak form of the Peaceman–Rachford splitting ([6.3](#)) seems to be too general for a convergence analysis. To remedy this, we will restrict the domains of the operators S_i, S such that the Steklov–Poincaré equation ([6.2](#)) and the Peaceman–Rachford splitting can be interpreted on $L^2(\Gamma)$ instead of on the dual spaces Λ_i^*, Λ^* . This comes at the cost of requiring more regularity of the weak solution and of the initial guess η_2^0 .

More precisely, we define the operators $\mathcal{S}_i : D(\mathcal{S}_i) \subseteq L^2(\Gamma) \rightarrow L^2(\Gamma)$ as

$$D(\mathcal{S}_i) = \{\mu \in \Lambda_i : S_i \mu \in L^2(\Gamma)^*\} \quad \text{and} \quad \mathcal{S}_i \mu = J^{-1} S_i \mu \text{ for } \mu \in D(\mathcal{S}_i).$$

Analogously, we introduce $\mathcal{S} : D(\mathcal{S}) \subseteq L^2(\Gamma) \rightarrow L^2(\Gamma)$ given by

$$D(\mathcal{S}) = \{\mu \in \Lambda : S \mu \in L^2(\Gamma)^*\} \quad \text{and} \quad \mathcal{S} \mu = J^{-1} S \mu \text{ for } \mu \in D(\mathcal{S}).$$

As the zero functional obviously is an element in $L^2(\Gamma)^*$, the unique solution $\eta \in \Lambda$ of the Steklov–Poincaré equation is in $D(\mathcal{S})$ and

$$S \eta = 0.$$

Remark 8.1. By the above construction, one obtains that $D(\mathcal{S}_1) \cap D(\mathcal{S}_2) \subseteq D(\mathcal{S})$ and

$$\mathcal{S} \mu = \mathcal{S}_1 \mu + \mathcal{S}_2 \mu, \quad \text{for all } \mu \in D(\mathcal{S}_1) \cap D(\mathcal{S}_2).$$

However, the definition of the domains do not ensure that $D(\mathcal{S})$ is equal to $D(\mathcal{S}_1) \cap D(\mathcal{S}_2)$, as $(S_1 + S_2) \mu \in L^2(\Gamma)^*$ does not necessarily imply that $S_i \mu \in L^2(\Gamma)^*$.

If the weak solution of the nonlinear elliptic equation (2.1) satisfies the additional regularity property stated in Assumption 2.6, then the corresponding solution of $\mathcal{S}\eta = 0$ is in fact an element in $D(\mathcal{S}_1) \cap D(\mathcal{S}_2)$. This propagation of regularity will be crucial when proving convergence of the Peaceman–Rachford splitting.

LEMMA 8.2. *If the Assumptions 2.1, 2.6 and 3.1 hold and $\mathcal{S}\eta = 0$, then $\eta \in D(\mathcal{S}_1) \cap D(\mathcal{S}_2)$.*

Proof. As $u = \{F_1\eta \text{ on } \Omega_1; F_2\eta \text{ on } \Omega_2\}$ is the weak solution of (2.1), we have that

$$\int_{\Omega} \alpha(\nabla u) \cdot \nabla v \, dx = - \int_{\Omega} (g(u) - f) v \, dx, \quad \text{for all } v \in C_0^\infty(\Omega).$$

The restrictions on r in Assumption 2.1 yield that $u \in W^{1,p}(\Omega) \hookrightarrow L^{2(r-1)}(\Omega)$. This together with the observation $|g(u)|^2 \leq C|u|^{2(r-1)}$ implies that $g(u) - f \in L^2(\Omega)$, i.e., the distributional divergence of $\alpha(\nabla u)$ is in $L^2(\Omega)^d$. By Assumption 2.6 and restricting to Ω_i , we arrive at

$$\begin{aligned} \alpha(\nabla F_i \eta) &\in H(\operatorname{div}, \Omega_i) \cap C(\overline{\Omega}_i)^d, \quad \alpha(\nabla F_i \eta) \cdot \nu_i \in L^\infty(\partial\Omega_i) \quad \text{and} \\ \nabla \cdot \alpha(\nabla F_i \eta) &= g(F_i \eta) - f_i \in L^2(\Omega_i). \end{aligned}$$

The $H(\operatorname{div}, \Omega_i)$ -version of Green's formula [8, Chapter 1, Corollary 2.1] then gives us

$$\int_{\Omega_i} \alpha(\nabla F_i \eta) \cdot \nabla v \, dx = - \int_{\Omega_i} \nabla \cdot \alpha(\nabla F_i \eta) v \, dx + \int_{\partial\Omega_i} \alpha(\nabla F_i \eta) \cdot \nu_i T_{\partial\Omega_i} v \, dS,$$

for all $v \in H^1(\Omega_i)$. Hence,

$$\begin{aligned} \langle \mathcal{S}_i \eta, \mu \rangle &= \int_{\Omega_i} \alpha(\nabla F_i \eta) \cdot \nabla R_i \mu \, dx + \int_{\Omega_i} (g(F_i \eta) - f_i) R_i \mu \, dx \\ &= \int_{\partial\Omega_i} \alpha(\nabla F_i \eta) \cdot \nu_i T_{\partial\Omega_i} R_i \mu \, dS = (\alpha(\nabla F_i \eta) \cdot \nu_i, \mu)_{L^2(\Gamma)}, \quad \text{for all } \mu \in \Lambda_i, \end{aligned}$$

which implies that $\eta \in D(\mathcal{S}_i)$, for $i = 1, 2$. $\square$

Remark 8.3. From the proof it is clear that the regularity assumption $\alpha(\nabla u) \in C(\overline{\Omega})^d$ is stricter than necessary, and could be replaced by assuming that the normal component of $\alpha(\nabla u)$ on Γ can be interpreted as an element in $L^2(\Gamma)^*$. However, characterizing the spatial regularity of u required to satisfy this weaker assumption demands a more elaborate trace theory than the one considered in section 3.

LEMMA 8.4. *If the Assumptions 2.1 and 3.1 hold, then the operators $\mathcal{S}_i$ are monotone, i.e.,*

$$(\mathcal{S}_i \eta - \mathcal{S}_i \mu, \eta - \mu)_{L^2(\Gamma)} \geq 0, \quad \text{for all } \eta, \mu \in D(\mathcal{S}_i),$$

and the operators $sI + \mathcal{S}_i : D(\mathcal{S}_i) \rightarrow L^2(\Gamma)$ are bijective for any $s > 0$.

Proof. The monotonicity follows by Lemma 7.2, as

$$\begin{aligned} (\mathcal{S}_i \eta - \mathcal{S}_i \mu, \eta - \mu)_{L^2(\Gamma)} &= (J^{-1} \mathcal{S}_i \eta - J^{-1} \mathcal{S}_i \mu, \eta - \mu)_{L^2(\Gamma)} \\ &= \langle \mathcal{S}_i \eta - \mathcal{S}_i \mu, \eta - \mu \rangle \geq 0, \quad \text{for all } \eta, \mu \in D(\mathcal{S}_i) \subseteq \Lambda_i. \end{aligned}$$

For a fixed $s > 0$ and an arbitrary $\mu \in L^2(\Gamma)$ we have, due to Corollary 7.7, that there exists a unique $\eta \in \Lambda_i$ such that $(sJ + \mathcal{S}_i)\eta = J\mu$ in Λ_i^* , i.e.,

$$\mathcal{S}_i \eta = J(\mu - s\eta) \in L^2(\Gamma)^*.$$

Hence, $\eta \in D(\mathcal{S}_i)$ and $(sI + \mathcal{S}_i)\eta = \mu$ in $L^2(\Gamma)$. The operators $sI + \mathcal{S}_i : D(\mathcal{S}_i) \rightarrow L^2(\Gamma)$ are therefore bijective. $\square$

The Peaceman–Rachford splitting on $L^2(\Gamma)$ is now given by finding $(\eta_1^n, \eta_2^n) \in D(\mathcal{S}_1) \times D(\mathcal{S}_2)$, for $n = 1, 2, \dots$, such that

$$(8.1) \quad \begin{cases} (sI + \mathcal{S}_1)\eta_1^{n+1} = (sI - \mathcal{S}_2)\eta_2^n, \\ (sI + \mathcal{S}_2)\eta_2^{n+1} = (sI - \mathcal{S}_1)\eta_1^{n+1}, \end{cases}$$

where $\eta_2^0 \in D(\mathcal{S}_2)$ is an initial guess. [Lemma 8.4](#) then directly yields the existence of the approximation.

COROLLARY 8.5. *If the [Assumptions 2.1](#) and [3.1](#) hold and $\eta_2^0 \in D(\mathcal{S}_2)$, then there exists a unique Peaceman–Rachford approximation $(\eta_1^n, \eta_2^n)_{n \geq 1} \subset D(\mathcal{S}_1) \times D(\mathcal{S}_2)$ given by (8.1) in $L^2(\Gamma)$.*

COROLLARY 8.6. *Let the [Assumptions 2.1](#) and [3.1](#) hold, $\eta_2^0 \in D(\mathcal{S}_2)$ and set $u_2^0 = F_2\eta_2^0$. The Peaceman–Rachford approximation $(\eta_1^n, \eta_2^n)_{n \geq 1} \subset D(\mathcal{S}_1) \times D(\mathcal{S}_2)$ also satisfies the weak formulation (6.3), and $(u_1^n, u_2^n)_{n \geq 1} = (F_1\eta_1^n, F_2\eta_2^n)_{n \geq 1}$ is a weak Robin–Robin approximation (5.2).*

Proof. Assume that $(\eta_1^n, \eta_2^n)_{n \geq 1} \subset D(\mathcal{S}_1) \times D(\mathcal{S}_2)$ is a Peaceman–Rachford approximation in $L^2(\Gamma)$. Then,

$$\begin{aligned} ((sI + \mathcal{S}_1)\eta_1^{n+1}, \mu)_{L^2(\Gamma)} &= ((sI - \mathcal{S}_2)\eta_2^n, \mu)_{L^2(\Gamma)}, & \text{for all } \mu \in L^2(\Gamma), \\ ((sI + \mathcal{S}_1)\eta_1^{n+1}, \mu)_{L^2(\Gamma)} &= \langle (sJ + \mathcal{S}_1)\eta_1^{n+1}, \mu \rangle, & \text{for all } \mu \in \Lambda_1, \text{ and} \\ ((sI - \mathcal{S}_2)\eta_2^n, \mu)_{L^2(\Gamma)} &= \langle (sJ - \mathcal{S}_2)\eta_2^n, \mu \rangle, & \text{for all } \mu \in \Lambda_2. \end{aligned}$$

This implies that

$$\langle (sJ + \mathcal{S}_1)\eta_1^{n+1}, \mu \rangle = \langle (sJ - \mathcal{S}_2)\eta_2^n, \mu \rangle, \quad \text{for all } \mu \in \Lambda = \Lambda_1 \cap \Lambda_2,$$

i.e., the first assertion of (6.3) holds. The same argumentation yields that the second assertion of (6.3) is valid. As $(\eta_1^n, \eta_2^n)_{n \geq 1}$ satisfies (6.3), [Lemma 6.3](#) directly implies that $(u_1^n, u_2^n)_{n \geq 1} = (F_1\eta_1^n, F_2\eta_2^n)_{n \geq 1}$ is a weak Robin–Robin approximation (5.2). $\square$

Remark 8.7. At a first glance, finding an initial guess satisfying $\eta_2^0 \in D(\mathcal{S}_2)$ might seem limiting, as the domain is not explicitly given. However, such an initial guess can, e.g., be found by solving $\langle \mathcal{S}_2\eta_2^0, \mu \rangle = 0$, for all $\mu \in \Lambda_2$.

With this $L^2(\Gamma)$ -framework the key part of the convergence proof follows by the abstract result [14, Proposition 1]. For sake of completeness we state a simplified version of the short proof in the current notation.

LEMMA 8.8. *Consider the solution of $\mathcal{S}\eta = 0$ and the Peaceman–Rachford approximation $(\eta_1^n, \eta_2^n)_{n \geq 1}$. If $\eta_2^0 \in D(\mathcal{S}_2)$ and the [Assumptions 2.1](#), [2.6](#) and [3.1](#) hold, then*

$$(8.2) \quad (\mathcal{S}_i\eta_i^n - \mathcal{S}_i\eta, \eta_i^n - \eta)_{L^2(\Gamma)} \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

for $i = 1, 2$.

Proof. By the hypotheses and [Lemma 8.2](#), we obtain that $\eta \in D(\mathcal{S}_1) \cap D(\mathcal{S}_2)$ and $\mathcal{S}_1\eta = -\mathcal{S}_2\eta$. Furthermore,

$$\eta_1^{n+1} = (sI + \mathcal{S}_1)^{-1}(sI - \mathcal{S}_2)\eta_2^n \in D(\mathcal{S}_1) \text{ and } \eta_2^{n+1} = (sI + \mathcal{S}_2)^{-1}(sI - \mathcal{S}_1)\eta_1^{n+1} \in D(\mathcal{S}_2).$$

Next, we introduce the notation

$$\mu^n = (sI + \mathcal{S}_2)\eta_2^n, \quad \mu = (sI + \mathcal{S}_2)\eta, \quad \lambda^n = (sI - \mathcal{S}_2)\eta_2^n \quad \text{and} \quad \lambda = (sI - \mathcal{S}_2)\eta,$$

which yields the representations

$$\begin{aligned} \eta &= \frac{\mu + \lambda}{2s}, & \mathcal{S}_2 \eta &= \frac{\mu - \lambda}{2}, & \mathcal{S}_1 \eta &= \frac{\lambda - \mu}{2}, \\ \eta_2^n &= \frac{\mu^n + \lambda^n}{2s}, & \mathcal{S}_2 \eta_2^n &= \frac{\mu^n - \lambda^n}{2}, \\ \eta_1^{n+1} &= \frac{\mu^{n+1} + \lambda^n}{2s}, & \mathcal{S}_1 \eta_1^{n+1} &= \frac{\lambda^n - \mu^{n+1}}{2}. \end{aligned}$$

The monotonicity of $\mathcal{S}_i$ then gives the bounds

$$\begin{aligned} 0 &\leq (\mathcal{S}_2 \eta_2^n - \mathcal{S}_2 \eta, \eta_2^n - \eta)_{L^2(\Gamma)} \\ &= \frac{1}{4s} ((\mu^n - \mu) - (\lambda^n - \lambda), (\mu^n - \mu) + (\lambda^n - \lambda))_{L^2(\Gamma)} \\ &= \frac{1}{4s} (\|\mu^n - \mu\|_{L^2(\Gamma)}^2 - \|\lambda^n - \lambda\|_{L^2(\Gamma)}^2), \end{aligned}$$

and

$$\begin{aligned} 0 &\leq (\mathcal{S}_1 \eta_1^{n+1} - \mathcal{S}_1 \eta, \eta_1^{n+1} - \eta)_{L^2(\Gamma)} \\ &= \frac{1}{4s} ((\lambda^n - \lambda) - (\mu^{n+1} - \mu), (\lambda^n - \lambda) + (\mu^{n+1} - \mu))_{L^2(\Gamma)} \\ &= \frac{1}{4s} (\|\lambda^n - \lambda\|_{L^2(\Gamma)}^2 - \|\mu^{n+1} - \mu\|_{L^2(\Gamma)}^2). \end{aligned}$$

Putting this together yields that

$$\|\mu^{n+1} - \mu\|_{L^2(\Gamma)}^2 \leq \|\lambda^n - \lambda\|_{L^2(\Gamma)}^2 \leq \|\mu^n - \mu\|_{L^2(\Gamma)}^2,$$

and we obtain the telescopic sum

$$0 \leq \sum_{n=0}^N (\|\mu^n - \mu\|_{L^2(\Gamma)}^2 - \|\mu^{n+1} - \mu\|_{L^2(\Gamma)}^2) \leq \|\mu^0 - \mu\|_{L^2(\Gamma)}^2 - \|\mu^{N+1} - \mu\|_{L^2(\Gamma)}^2,$$

i.e., $\|\mu^n - \mu\|_{L^2(\Gamma)}^2 - \|\mu^{n+1} - \mu\|_{L^2(\Gamma)}^2 \rightarrow 0$ as $n \rightarrow \infty$. The latter together with the bounds above imply the sought after limits (8.2). $\square$

THEOREM 8.9. *Consider the Peaceman–Rachford approximation $(\eta_1^n, \eta_2^n)_{n \geq 1}$, given by (8.1), of the Steklov–Poincaré equation $\mathcal{S}\eta = 0$ in $L^2(\Gamma)$, together with the corresponding Robin–Robin approximation $(u_1^n, u_2^n)_{n \geq 1} = (F_1 \eta_1^n, F_2 \eta_2^n)_{n \geq 1}$ of the weak solution $u = \{F_1 \eta \text{ on } \Omega_1; F_2 \eta \text{ on } \Omega_2\}$ to the nonlinear elliptic equation (2.1). If $\eta_2^0 \in D(\mathcal{S}_2)$ and the Assumptions 2.1, 2.6 and 3.1 hold, then*

$$(8.3) \quad \|\eta_1^n - \eta\|_{\Lambda_1} + \|\eta_2^n - \eta\|_{\Lambda_2} \rightarrow 0, \quad \text{and} \quad \|u_1^n - u\|_{W^{1,p}(\Omega_1)} + \|u_2^n - u\|_{W^{1,p}(\Omega_2)} \rightarrow 0,$$

as n tends to infinity.

Proof. By the monotonicity bound in Lemma 7.2, the property that $\eta_i^n, \eta \in D(\mathcal{S}_i)$ and Lemma 8.8, we have the limits

$$\begin{aligned} c_i (\|\nabla(F_i \eta_i^n - F_i \eta)\|_{L^p(\Omega_i)^d}^p + \|F_i \eta_i^n - F_i \eta\|_{L^r(\Omega_i)}^r) &\leq \langle \mathcal{S}_i \eta_i^n - \mathcal{S}_i \eta, \eta_i^n - \eta \rangle \\ &= (\mathcal{S}_i \eta_i^n - \mathcal{S}_i \eta, \eta_i^n - \eta)_{L^2(\Gamma)} \rightarrow 0, \quad \text{as } n \rightarrow \infty, \end{aligned}$$

for $i = 1, 2$. Hence, each of the terms $\|\nabla(F_i\eta_i^n - F_i\eta)\|_{L^p(\Omega_i)^d}$ and $\|F_i\eta_i^n - F_i\eta\|_{L^r(\Omega_i)}$ tend to zero, which yields that

$$\|\eta_i^n - \eta\|_{\Lambda_i} \leq C\|F_i\eta_i^n - F_i\eta\|_{V_i} \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

for $i = 1, 2$. The desired convergence (8.3) is then proven, as $\|\cdot\|_{V_i}$ and $\|\cdot\|_{W^{1,p}(\Omega_i)}$ are equivalent norms. $\square$

REFERENCES

- [1] V. I. AGOSHKOV AND V. I. LEBEDEV, *Variational algorithms of the domain decomposition method [translation of Preprint 54, Akad. Nauk SSSR, Otdel. Vychisl. Mat., Moscow, 1983]*, vol. 5, 1990, pp. 27–46. Soviet Journal of Numerical Analysis and Mathematical Modelling.
- [2] H. BERNINGER, R. KORNUBER, AND O. SANDER, *Convergence behaviour of Dirichlet-Neumann and Robin methods for a nonlinear transmission problem*, in Domain decomposition methods in science and engineering XIX, vol. 78 of Lect. Notes Comput. Sci. Eng., Springer, Heidelberg, 2011, pp. 87–98, https://doi.org/10.1007/978-3-642-11304-8_8, https://doi.org/10.1007/978-3-642-11304-8_8.
- [3] W. CHEN, X. XU, AND S. ZHANG, *On the optimal convergence rate of a Robin-Robin domain decomposition method*, J. Comput. Math., 32 (2014), pp. 456–475, <https://doi.org/10.4208/jcm.1403-m4391>, <https://doi-org.ludwig.lub.lu.se/10.4208/jcm.1403-m4391>.
- [4] D. L. COHN, *Measure theory*, Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks], Birkhäuser/Springer, New York, second ed., 2013, <https://doi.org/10.1007/978-1-4614-6956-8>, <https://doi.org/10.1007/978-1-4614-6956-8>.
- [5] M. DISCACCIATI, A. QUARTERONI, AND A. VALLI, *Robin-Robin domain decomposition methods for the Stokes-Darcy coupling*, SIAM J. Numer. Anal., 45 (2007), pp. 1246–1268, <https://doi.org/10.1137/06065091X>, <https://doi.org/10.1137/06065091X>.
- [6] M. DRYJA AND W. HACKBUSCH, *On the nonlinear domain decomposition method*, BIT, 37 (1997), pp. 296–311, <https://doi.org/10.1007/BF02510214>, <https://doi.org/10.1007/BF02510214>.
- [7] M. J. GANDER, L. HALPERN, AND F. NATAF, *Optimized Schwarz methods*, in Domain decomposition methods in sciences and engineering (Chiba, 1999), DDM.org, Augsburg, 2001, pp. 15–27.
- [8] V. GIRAULT AND P.-A. RAVIART, *Finite element approximation of the Navier-Stokes equations*, vol. 749 of Lecture Notes in Mathematics, Springer-Verlag, Berlin-New York, 1979.
- [9] W. GUO AND L. S. HOU, *Generalizations and accelerations of Lions' nonoverlapping domain decomposition method for linear elliptic PDE*, SIAM J. Numer. Anal., 41 (2003), pp. 2056–2080, <https://doi.org/10.1137/S0036142902407150>, <https://doi-org.ludwig.lub.lu.se/10.1137/S0036142902407150>.
- [10] A. KUFNER, O. JOHN, AND S. FUČÍK, *Function spaces*, Noordhoff International Publishing, Leyden; Academia, Prague, 1977. Monographs and Textbooks on Mechanics of Solids and Fluids; Mechanics: Analysis.
- [11] P. D. LAX, *Functional analysis*, Pure and Applied Mathematics (New York), Wiley-Interscience [John Wiley & Sons], New York, 2002.
- [12] G. M. LIEBERMAN, *Boundary regularity for solutions of degenerate elliptic equations*, Nonlinear Anal., 12 (1988), pp. 1203–1219, [https://doi.org/10.1016/0362-546X\(88\)90053-3](https://doi.org/10.1016/0362-546X(88)90053-3), [https://doi.org/10.1016/0362-546X\(88\)90053-3](https://doi.org/10.1016/0362-546X(88)90053-3).
- [13] P.-L. LIONS, *On the Schwarz alternating method. III. A variant for nonoverlapping subdomains*, in Third International Symposium on Domain Decomposition Methods for Partial Differential Equations (Houston, TX, 1989), SIAM, Philadelphia, PA, 1990, pp. 202–223.
- [14] P.-L. LIONS AND B. MERCIER, *Splitting algorithms for the sum of two nonlinear operators*, SIAM J. Numer. Anal., 16 (1979), pp. 964–979, <https://doi.org/10.1137/0716071>, <https://doi.org/10.1137/0716071>.
- [15] S. H. LUI, *A Lions non-overlapping domain decomposition method for domains with an arbitrary interface*, IMA J. Numer. Anal., 29 (2009), pp. 332–349, <https://doi.org/10.1093/imanum/drm011>, <https://doi-org.ludwig.lub.lu.se/10.1093/imanum/drm011>.
- [16] J. NAUMANN AND C. G. SIMADER, *Measure and Integration on Lipschitz-Manifolds*, Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Fakultät II, Institut für Mathematik, 2011, <https://doi.org/http://dx.doi.org/10.18452/2773>.

- [17] J. NEČAS, *Direct methods in the theory of elliptic equations*, Springer Monographs in Mathematics, Springer, Heidelberg, 2012, <https://doi.org/10.1007/978-3-642-10455-8>, <https://doi.org/10.1007/978-3-642-10455-8>. Translated from the 1967 French original by Gerard Tronel and Alois Kufner, Editorial coordination and preface by Šárka Nečasová and a contribution by Christian G. Simader.
- [18] A. QUARTERONI AND A. VALLI, *Domain decomposition methods for partial differential equations*, Numerical Mathematics and Scientific Computation, The Clarendon Press, Oxford University Press, New York, 1999. Oxford Science Publications.
- [19] T. ROUBÍČEK, *Nonlinear partial differential equations with applications*, vol. 153 of International Series of Numerical Mathematics, Birkhäuser/Springer Basel AG, Basel, second ed., 2013, <https://doi.org/10.1007/978-3-0348-0513-1>, <https://doi.org/10.1007/978-3-0348-0513-1>.
- [20] S. SCHREIBER AND R. HOCHMUTH, *On the equivalence of transmission problems in nonoverlapping domain decomposition methods for quasilinear PDEs*, Numer. Funct. Anal. Optim., 31 (2010), pp. 596–615, <https://doi.org/10.1080/01630563.2010.490625>, <https://doi.org/10.1080/01630563.2010.490625>.
- [21] X.-C. TAI AND M. ESPEDAL, *Rate of convergence of some space decomposition methods for linear and nonlinear problems*, SIAM J. Numer. Anal., 35 (1998), pp. 1558–1570, <https://doi.org/10.1137/S0036142996297461>, <https://doi.org/10.1137/S0036142996297461>.
- [22] X.-C. TAI AND J. XU, *Global and uniform convergence of subspace correction methods for some convex optimization problems*, Math. Comp., 71 (2002), pp. 105–124, <https://doi.org/10.1090/S0025-5718-01-01311-4>, <https://doi.org/10.1090/S0025-5718-01-01311-4>.
- [23] A. TOSELLI AND O. WIDLUND, *Domain decomposition methods—algorithms and theory*, vol. 34 of Springer Series in Computational Mathematics, Springer-Verlag, Berlin, 2005, <https://doi.org/10.1007/b137868>, <https://doi-org.ludwig.lub.lu.se/10.1007/b137868>.
- [24] X. XU AND L. QIN, *Spectral analysis of Dirichlet-Neumann operators and optimized Schwarz methods with Robin transmission conditions*, SIAM J. Numer. Anal., 47 (2010), pp. 4540–4568, <https://doi.org/10.1137/080713653>, <https://doi-org.ludwig.lub.lu.se/10.1137/080713653>.
- [25] E. ZEIDLER, *Nonlinear functional analysis and its applications. II/B*, Springer-Verlag, New York, 1990, <https://doi.org/10.1007/978-1-4612-0985-0>, <https://doi.org/10.1007/978-1-4612-0985-0>. Nonlinear monotone operators, Translated from the German by the author and Leo F. Boron.